UTAH GEOLOGICAL SURVEY
a division of
Utah Department of Natural Resources

Plate 1
Utah Geological Survey Map 223
Geologic Map of the Skinner Peaks Quadrangle



Base from U.S. Geological Survey, Skinner Peaks 7.5' quadrangle, 2001
Geologic data and base map in NAD 1927

Although this product represents the work of professional scientists, the Utah Department of Natural Resources, Utah Geological Survey, makes no warranty, expressed or implied, regarding its suitability for a particular use. The Utah Department of Natural Resources, Utah Geological Survey, shall not be liable under any circumstances for any direct, indirect, special, incidental, or consequential damages with respect to claims by users of this product.

For use at 1:24,000 scale only. The UGS does not guarantee accuracy or completeness of data.

SCALE 1:24,000

1 0.5 0 1 MILE
1000 0 1000 2000 3000 4000 5000 6000 7000 FEET
1 0.5 0 1 KILOMETER

CONTOUR INTERVAL 20 FEET
SUPPLEMENTARY CONTOUR INTERVAL 10-FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

12°25'
TRUE NORTH
MAGNETIC NORTH
APPROXIMATE MEAN DECLINATION, 2007

GEOLOGIC MAP OF THE SKINNER PEAKS QUADRANGLE, JUAB AND SANPETE COUNTIES, UTAH

by

Tracey J. Felger[1], Donald L. Clark[2], and Michael D. Hylland[2]

2007

[1]U.S. Geological Survey, 2255 N. Gemini Drive, Flagstaff, AZ 86001
[2]Utah Geological Survey, P.O. Box 146100, Salt Lake City, UT 84114-6100

UTAH
QUADRANGLE LOCATION

1	2	3	1. Sage Valley
4		5	2. Juab 3. Levan 4. Mills 5. Chriss Canyon 6. Scipio North 7. Hells Kitchen Canyon SW 8. Hells Kitchen Canyon SE
6	7	8	

ADJOINING 7.5' QUADRANGLE NAMES

Project Manager: Robert Biek
Field mapping by authors 1988-89, 2003-05
Digital Cartography and Photogrammetry: Darryl Greer, Basia Matyjasik, Kent Brown, Lori Douglas, Donald Clark
Cartography: James Parker
ISBN 1-55791-762-0

Utah Geological Survey, 1594 W. North Temple,
P.O. Box 146100, Salt Lake City, Utah 84114-6100
Phone: 801-537-3300; fax 801-537-3400
geology.utah.gov

UTAH GEOLOGICAL SURVEY
a division of
Utah Department of Natural Resources

Plate 2
Utah Geological Survey Map 223
Geologic Map of the Skinner Peaks Quadrangle

MAP UNIT DESCRIPTIONS

QUATERNARY

Alluvial deposits

Qal **Stream alluvium** (Holocene) – Gravel, sand, and silt with minor clay and scattered cobbles and boulders; clasts well rounded to subangular; generally stratified; deposited in modern stream channels and flood plains, and adjacent to Chicken Creek Reservoir; locally forms terraces less than 30 feet (10 m) above stream level that grade downslope into upper Holocene alluvial-fan deposits (unit Qafy); also grades to map units Qaf and Qac; may include small alluvial fans and minor colluvium along steep stream embankments; exposed thickness less than 30 feet (<10 m).

Qafy **Younger fan alluvium** (upper Holocene) – Pebble and cobble gravel, locally bouldery, in a matrix of sand, silt, and minor clay; clasts angular to subrounded; deposited by intermittent streams, debris flows, and debris floods graded to modern stream level; deposits form discrete fans, typically with bar and swale topography; local weak stage I carbonate morphology (Birkeland and others, 1991); exposed thickness less than 15 feet (<5 m).

Qafi **Intermediate-age fan alluvium** (lower Holocene) – Pebble and cobble gravel, locally bouldery, in a matrix of sand, silt, and minor clay; clasts angular to well rounded; preserved as intermediate-level remnants incised by modern streams and locally buried by younger fan alluvium (Qafy); incised by, and grades downslope into, undifferentiated fan alluvium (Qaf); stage I+ carbonate morphology (Birkeland and others, 1991); exposed thickness less than 15 feet (<5 m).

Qafo **Older fan alluvium** (upper to middle Pleistocene) – Pebble and cobble gravel, locally bouldery, in a matrix of sand, silt, and minor clay; clasts subangular to well rounded; preserved as relatively high, isolated remnants that lack geomorphologic; stage II to III+ carbonate morphology (Birkeland and others, 1991); exposed thickness less than 50 feet (<15 m).

Qaf **Undifferentiated fan alluvium** (upper Holocene to middle? Pleistocene) – Proximal facies: pebble and cobble gravel, locally bouldery, in a matrix of sand, silt, and minor clay; clasts subangular to well rounded; distal facies: silt and fine sand with minor clay; deposited by perennial and intermittent streams, debris flows, and debris floods graded to or slightly above modern stream level; includes a significant component of eolian silt; deposits form large, low-gradient fans that cover most of the floor of Juab Valley and a small area of Flat Canyon in the southeast corner of the quadrangle; includes local areas of active fan deposition (unit Qafy); total deposits of unit Qaf to small to map separately; locally includes small unmapped debris-flow deposits; maximum thickness greater than 600 feet (>180 m); Qaf represents the southernmost portion of the Juab Valley basin-fill deposits, which thicken northward.

Qap **Piedmont alluvium** (lower Holocene to upper Pleistocene) – Gently sloping mantle of clay- to boulder-size material overlying a truncated bedrock surface on the east flank of the West Hills in the northwest part of the quadrangle; exposed thickness from 0 to over 40 feet (0-12+ m); total thickness unknown.

Lacustrine deltaic deposits

Deltaic and lacustrine sediment was deposited in the Sevier River estuary in the southwestern part of the quadrangle during the transgressive phase of Lake Bonneville and during its highstand (Bonneville level). The shoreline elevation in the quadrangle is approximately 5100 feet (1555 m) (also see Chrisney, 1962, Oviatt, 1992).

Qdf **Deltaic (estuarine) fines** (upper Pleistocene) – Light-brown, unconsolidated, coarse- to fine-grained sand, silt, and mud deposited by Lake Bonneville; deposits are finely laminated and cross-laminated; soft-sediment deformation structures and ripple cross-lamination are common near the base of the exposed section; exposed thickness up to about 20 feet (0-6 m), but total thickness uncertain.

Qc **Colluvial deposits**

Colluvium (upper Holocene) – Slope-wash deposits of clay- to boulder-size, locally derived sediments deposited on upland slopes adjacent to drainages; generally less than 20 feet (<6 m) thick.

Qco **Older colluvium** (lower Holocene to upper Pleistocene) – Locally derived clast- and matrix-supported pebble and cobble gravel, locally bouldery; clasts angular; matrix consists of sand, silt, and clay; grain size generally decreases downslope; forms a dissected surface on various bedrock units around the base of Skinner Peaks; thickness from 0 to over 40 feet (0-12+ m).

Eolian deposits

Qes **Eolian sand** (Holocene) – Moderate-brown sand forming hummocky surfaces along the southern map border; derived from lacustrine or beach deposits or bedrock near Sevier Bridge Reservoir; exposed thickness from 0 to 20 feet (0-6 m); total thickness unknown.

Mixed-environment deposits

Qca **Colluvium and alluvium, undivided** (Holocene to upper Pleistocene) – Undifferentiated hillslope colluvium, stream and fan alluvium, and small landslide deposits; thickness up to approximately 50 feet (0-15 m).

Qac **Alluvium and colluvium, undivided** (Holocene) – Undifferentiated stream and fan alluvium and hillslope colluvium; may also locally include eolian silt and sand; deposited in shallow drainages associated with intermittent streams, and in small, shallow basins; thickness up to approximately 50 feet (0-15 m).

Mass-movement deposits

Qmsy **Younger landslide deposits** (historical to upper Pleistocene?) – Angular, poorly sorted debris and blocks of rock, similar to Qmso; derived from volcanic rock units, Arapien Shale, and Green River Formation; includes one slide along State Route 28 having probable historical movement; thickness highly variable.

Qmso **Older landslide deposits** (upper? to middle Pleistocene) – Angular, poorly sorted debris and blocks of rock; derived from the Green River Formation, volcanic rock units, and surficial deposits; dissected and subdued morphology distinguishes these deposits from unit Qmsy; thickness highly variable.

Human-made deposits

Qf **Fill** (historical) – Fill material along Interstate Highway 15 (I-15); fill and local earth materials that have been graded or reworked in the vicinity of the Mills interchange on I-15; local earth materials used to construct dams for stock ponds and berms to divert drainages; thickness 0 to 20 feet (0-6 m).

Qa/ Qaf **Stacked-unit deposits**

Undifferentiated alluvium over deltaic fines (Holocene to upper Pleistocene) – Fine-grained alluvial deposits forming a veneer over Lake Bonneville deltaic fines; forms flat-topped and irregular surfaces covering underlying materials; thickness 0 to 10 feet (0-3 m).

Qa/ QTaf **Undifferentiated alluvium over oldest alluvial-fan deposits** (Holocene/Pleistocene to Pliocene?) – Fine-grained alluvial deposits forming a veneer over oldest alluvial-fan deposits; forms gently sloping to irregular surfaces covering underlying materials; thickness 0 to 10 feet (0-3 m).

Qf/ QTaf **Undifferentiated human-made deposits over oldest alluvial-fan deposits** (upper Pleistocene/Holocene to Pliocene?) – Clay- to pebble-size lacustrine sediment forming a veneer over oldest alluvial-fan deposits located immediately below the Bonneville shoreline; thickness less than 20 feet (<6 m).

Qf/ QTfg **Undifferentiated human-made deposits over Tertiary fines and gravels** (upper Pleistocene/Pliocene? to Miocene?) – Clay- to pebble-size lacustrine sediments forming a veneer over Tertiary fines and gravels; deposited immediately below the Bonneville shoreline; thickness less than 20 feet (<6 m).

QUATERNARY-TERTIARY

QTaf **Oldest alluvial-fan deposits** (Pleistocene to Pliocene?) – Yellowish-gray to light-brownish-orange, coarse- to fine-grained, poorly sorted, deeply dissected alluvial, debris flow, and colluvial sediments shed from San Pitch Mountains to the east (much of it derived from the Green River Formation); typically unconsolidated, but locally consolidated; covers a large portion of the South Hills and Flat Canyon areas; clasts include brown sandstone, quartzite, including welded Fernow Quartz Latite clasts (34.83 ± 0.15 Ma; Clark, 2003), thus suggesting a northward source area; ledge-forming unit that is medium to thickly bedded with very low-angle primary stratification; locally includes small outcrops of other members too small to map separately; unit thickness from about 50 to 90 feet (15-25 m).

Tp2 **Member 2** – Grayish-pink, poorly to moderately welded rhyolitic ash-flow tuff, with phenocrysts of quartz, sanidine(?), and minor biotite, as well as pumice(?) and dark rock fragments; sample PR-16 yielded an $^{40}Ar/^{39}Ar$ age of 35.19 ± 0.14 Ma on sanidine; available data suggest that member 2 correlates with the Fernow Quartz Latite exposed in the southern East Tintic Mountains and Sage Valley; typically forms ledge-forming slopes; locally includes small outcrops of other members too small to map separately; from 40 to 100 feet (12-30 m) thick.

Tp1 **Member 1** – Upper part of unit (about 200 feet [60 m]) exposed at Painted Rocks consists of gray and light-bluish-gray volcanic conglomerate and tuffaceous sandstone; pebbles to boulder-size, dark to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

Thmp **Hornblende monzonite porphyry** (upper Oligocene) – Pinkish-gray to light-gray hornblende monzonite porphyry that weathers to a yellow and brown grusy talus with a cinder-like appearance; two small stocks, exposed just north of Little Salt Creek as dikes or sills within the Jurassic Arapien Shale near the contact with the Green River Formation; similar rock types form stocks, plugs, dikes, and sills in adjacent quadrangles to the east and north (John, 1964, 1972; Anderson, 1991; Weiss and others, 2003); the rock has an aphanitic groundmass of K-feldspar and andesine and phenocrysts of feldspar, hornblende, and biotite (also see John, 1964, 1972); isotopic and related intrusions are 23.3 to 23.8 million years (Witkind and Marvin, 1989; Auby, 1991); we obtained a new $^{40}Ar/^{39}Ar$ isochron age on hornblende of 23.97 ± 0.14 Ma from a hornblende monzonite porphyry (sample PR-15) that crops out one mile (1.6 km) northeast of the Skinner Peaks quadrangle exposures (in the Chriss Canyon quadrangle) (NMGRL and UGS, 2005; Clark, 2006); exposed width less than 30 feet (<9 m).

unconformity

TERTIARY SEDIMENTARY AND VOLCANIC ROCKS

Tv **Undifferentiated Oligocene-Eocene volcanic rocks** – Subsurface only.

Formation of Painted Rocks *(new informal name)* (lower Oligocene to middle Eocene) We recommend use of this informal unit terminology rather than applying the name Goldens Ranch Formation because the exposures are not a regional study of the volcanic rocks. Refer to Correlation of Map Units. Further clarification of volcanic stratigraphy in this region is warranted.

Tps-d, Tps-s **Members 4 and 5, undivided** – Locally mapped as combined unit in small exposures.

Tps **Member 5** – Gray to brown volcanic conglomerate, "salt and pepper" quartz sandstone, and tuffaceous sandstone; poorly consolidated, rubbly to sandy exposures; volcanic conglomerate and sandstone of Painted Rocks; clasts of dark-pink subangular to subrounded volcanic clasts and lesser quartzite and carbonate clasts; conglomerate exposures typically contain abundant Fernow Quartz Latite clasts, and South Hills exposures also locally contain clasts of tuff derived from member 4 and tuff of Little Sage Valley (Clark, 2003); sandstone forms a few ledges but typically weathers to moderate- to dark-brown sandy exposures; total thickness unknown due to poor exposures and structural complications, but possibly 1000 to 2000 feet (300-600 m) thick; locally includes small areas of other members too small to map separately; interbedded with these lahar and alluvial deposits are welded tuff and ash-flow tuff units described below.

Tpsw **Member 5, welded-tuff unit** – Grayish-pink densely welded rhyolite with phenocrysts of quartz and biotite (Clark, 2006); unit is typically fractured to the western margin of or foliation; forms small cliffs; sample PR-28 yielded an $^{40}Ar/^{39}Ar$ age of 29.81 ± 0.14 Ma (on biotite); as mapped may also include small exposures of Tp_{5d}; stratigraphic position within Tp_5 not well constrained; thickness up to approximately 50 feet (0-15 m).

Tpsa **Member 5, ash-flow-tuff unit** – Ivory, light gray, poorly to moderately welded dacite with phenocrysts of quartz, feldspar, and bronze biotite flakes (see Clark, 2006); forms slopes to small cliffs; locally present beneath welded tuff unit; as mapped may include small exposures of Tp_{5d}; stratigraphic position within Tp_5 not well constrained; thickness up to approximately 50 feet (0-15 m).

Tp1-4 **Members 1, 2, 3 and 4, undivided** – Combined unit; cross section A-A' only.

Tp2-4 **Members 2, 3 and 4, undivided** – Combined unit; cross section B-B' only.

Tp4 **Member 4** – Very pale orange to grayish-orange-pink, poorly to moderately welded rhyolitic ash-flow tuff with conspicuous pumice and rock fragments and some quartz and biotite phenocrysts (Clark, 2006); exposures vary from poor to good; weathers to honeycomb pattern and small caves; vertical fracturing and lack of layering typical; distinctive rock unit capping Painted Rocks cuesta at Yuba State Park (section 32, T. 16 S., R. 1 W. and section 5, T. 17 S., R. 1 W., Hells Kitchen Canyon SW quadrangle); sample PR-9 yielded an $^{40}Ar/^{39}Ar$ single crystal fusion sanidine age of 34.60 ± 0.13 Ma, which supercedes prior K-Ar ages from Witkind and Marvin (1989); locally includes small outcrops of other members too small to map separately; locally quarried for use as poultry grit and soil mineralizer (Pratt and Callaghan, 1970); approximately 70 to 150 feet (20-45 m) thick.

Tp2-3 **Members 2 and 3, undivided** – Locally mapped as combined unit in small exposures.

Tp3 **Member 3** – Pinkish-gray to locally moderate-reddish-orange (near Skinner Peaks) quartz-rich sandstone; unique unit containing about 60-80% coarse to very coarse sand-size, euhedral to subangular quartz grains; also locally includes rounded pebbles and cobbles of volcanic rocks and quartzite, including welded Fernow Quartz Latite clasts (34.83 ± 0.15 Ma; Clark, 2003), thus suggesting a northward source area; ledge-forming unit that is medium to thickly bedded with very low-angle primary stratification; locally includes small outcrops of other members too small to map separately; unit thickness from about 50 to 90 feet (15-25 m).

TKnh **North Horn Formation** (Paleocene? to Upper Cretaceous [Maastrichtian]) – Subsurface only (refer to cross section A-A'); includes gray to dark-gray mudstone, sandstone, and conglomerate.

unconformity?

TKu **TERTIARY-CRETACEOUS**

Tertiary-Cretaceous strata, undifferentiated – Corresponds to Colton, Flagstaff, and North Horn Formations; subsurface only (see Correlation of Map Units).

TKuu **Upper unit of undifferentiated Tertiary-Cretaceous strata** (middle to lower Eocene? to Upper Cretaceous?) – Conglomeratic outcrop belt on the northeast side of Skinner Peaks consisting of carbonate-clast conglomerate overlying red quartzite-clast conglomerate; the carbonate clast conglomerate predominantly pebbles and cobbles; gray to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

TKul **Lower unit of undifferentiated Tertiary-Cretaceous strata** (lower Eocene? to Upper Cretaceous) – One conglomeratic outcrop belt on the northeast side of Skinner Peaks consisting of carbonate-clast conglomerate overlying red quartzite-clast conglomerate; the carbonate clast conglomerate predominantly pebbles and cobbles; gray to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

Goldens Ranch Formation (upper? to middle Eocene) – Originally defined by Muessig (1953a, 1953b). Revised by Meibos (1983) to include only the lower part of Muessig's formation and divided into three members, named the upper Sage Valley Limestone, middle Hall Canyon Conglomerate, and lower Chicken Creek Tuff. We do not follow the suggested revision of the Goldens Ranch Formation by Witkind and Marvin (1989), and believe that a regional study of the volcanic rocks is needed to clarify outstanding stratigraphic issues. Refer to Correlation of Map Units. After Meibos (1983) and excluding the Sage Valley Limestone Member, which is not present in this quadrangle:

Ku (Ku, Kisp) **Hall Canyon Conglomerate Member** – Gray, poorly consolidated, cobble conglomerate and volcanic conglomerate; clast composition varies from quartzite-carbonate to volcanic; clasts are angular to subrounded pebbles to boulders in an ashy to sandy matrix; typically forms rubbly slopes; mapped in the northwest part of the quadrangle near Chicken Creek Reservoir where it directly overlies the Green River Formation; apparently laterally equivalent to lower portion of member 1 – formation of Painted Rocks; exposed thickness about 100 to 300 feet (30-90 m).

Tgc **Chicken Creek Tuff Member** – Grayish-pink to light-gray, poorly welded, vitric, dacitic ash-flow tuff with pumice lapilli, volcanic and quartzite rock fragments, and minor biotite; generally not well exposed; named for small exposure southwest of Chicken Creek Reservoir in section 19, T. 15 S., R. 1 W.; other exposures are in and near "Milky Wash"; query indicates uncertain designation; $^{40}Ar/^{39}Ar$ age of 38.61 ± 0.13 Ma from Sage Valley (Clark, 2003) supercedes K-Ar ages from Everenden and James (1964) and Witkind and Marvin (1989); source unknown; near the reservoir, located within the Hall Canyon Conglomerate Member from roughly 200 feet (60 m) or less above the top of the Green River Formation; near Painted Rocks and "Milky Wash" located stratigraphically within the lower part of member 1 (about 470 feet [145 m] above base); exposed thickness less than 100 feet (30 m), but total dip-corrected thickness in Placid Barton well is 170 feet (50 m).

Tg **Green River Formation** (middle Eocene) – West Hills exposures (see Clark, 1987, 1990; Felger, 1991): upper part is interbedded moderate-greenish-yellow and moderate-greenish-gray mudstone, and yellowish-gray, fossiliferous, platy, micritic limestone; mudstones become ashy near contact with overlying Hall Canyon Conglomerate; middle part is orange, brown, and gray conglomerate, mudstone, sandstone, and siltstone; lower part is green mudstone capped by yellowish-gray to light-olive-gray stromatolitic limestone; the formation is nearly 900 feet (275 m) thick (Clark, 1990).

Skinner Peaks area exposures: clastic lithofacies previously referred to as the "Tawny facies" (Zeller, 1949; Hardy and Zeller, 1953; Vogel, 1957; Millen, 1982; Felger, 1991; Weiss and others, 2003) composed of green, red, and variegated mudstone, and yellowish-tan coarse-grained sandstone, conglomerate, conglomeratic sandstone, and limestone; this lithofacies grades laterally eastward into the upper limestone member of the Green River Formation in the adjacent Chriss Canyon quadrangle (Millen, 1982; Weiss and others, 2003); sandstone near top of section contains reptilian(?) vertebrate fossils (Vogel, 1957; Felger, 1991); on the north side of Chriss Creek, unit has a heavy colluvial cover or locally a veneer of QTaf that has not been mapped separately; contact with overlying volcanic rocks not exposed, but present in Barton #1 well and is possibly conformable; directly overlies the Jurassic Arapien Shale at and north of Skinner Peaks; total thickness in subsurface from 1100 to 1300 feet (335-395 m).

Tf **Flagstaff Formation** (lower Eocene to upper Paleocene) – As mapped in West Hills, gravish-yellow to pale-reddish-orange calcareous mudstone, sandstone, sandy limestone, limestone, and conglomerate (also see Lambert, 1976; Vorce, 1979); base not exposed; exposed thickness 525 feet (160 m), but total thickness in West Hills is about 650 feet (200 m) (Clark, 1987, 1990); 295 feet (90 m) logged in the Skinner #1 drill hole; locally absent near Skinner Peaks or included within map unit TKuu and/or TKul.

TKnh **North Horn Formation** (Paleocene? to Upper Cretaceous [Maastrichtian]) – Subsurface only (refer to cross section A-A'); includes gray to dark-gray mudstone, sandstone, sandy limestone, limestone, and conglomerate (also see Lambert, 1976; Vorce, 1979); base not exposed; exposed thickness 525 feet (160 m), but total thickness in West Hills is about 650 feet (200 m) (Clark, 1987, 1990); 295 feet (90 m) logged in the Skinner #1 drill hole; locally absent near Skinner Peaks or included within map unit TKuu and/or TKul.

unconformity?

TKu **TERTIARY-CRETACEOUS**

Tertiary-Cretaceous strata, undifferentiated – Corresponds to Colton, Flagstaff, and North Horn Formations; subsurface only (see Correlation of Map Units).

TKuu **Upper unit of undifferentiated Tertiary-Cretaceous strata** (middle to lower Eocene? to Upper Cretaceous?) – Conglomeratic outcrop belt on the northeast side of Skinner Peaks consisting of carbonate-clast conglomerate overlying red quartzite-clast conglomerate; the carbonate clast conglomerate predominantly pebbles and cobbles; gray to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

Tp1 **Member 1** – Upper part of unit (about 200 feet [60 m]) exposed at Painted Rocks consists of gray and light-bluish-gray volcanic conglomerate and tuffaceous sandstone; pebbles to boulder-size, dark to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

Tgh **Hall Canyon Conglomerate Member** – Gray, poorly consolidated, cobble conglomerate and volcanic conglomerate; clast composition varies from quartzite-carbonate to volcanic; clasts are angular to subrounded pebbles to boulders in an ashy to sandy matrix; typically forms rubbly slopes; mapped in the northwest part of the quadrangle near Chicken Creek Reservoir where it directly overlies the Green River Formation; apparently laterally equivalent to lower portion of member 1 – formation of Painted Rocks; exposed thickness about 100 to 300 feet (30-90 m).

Tgc **Chicken Creek Tuff Member** – Grayish-pink to light-gray, poorly welded, vitric, dacitic ash-flow tuff with pumice lapilli, volcanic and quartzite rock fragments, and minor biotite; generally not well exposed; named for small exposure southwest of Chicken Creek Reservoir in section 19, T. 15 S., R. 1 W.; other exposures are in and near "Milky Wash"; query indicates uncertain designation; $^{40}Ar/^{39}Ar$ age of 38.61 ± 0.13 Ma from Sage Valley (Clark, 2003) supercedes K-Ar ages from Everenden and James (1964) and Witkind and Marvin (1989); source unknown; near the reservoir, located within the Hall Canyon Conglomerate Member from roughly 200 feet (60 m) or less above the top of the Green River Formation; near Painted Rocks and "Milky Wash" located stratigraphically within the lower part of member 1 (about 470 feet [145 m] above base); exposed thickness less than 100 feet (30 m), but total dip-corrected thickness in Placid Barton well is 170 feet (50 m).

Tg **Green River Formation** (middle Eocene) – West Hills exposures (see Clark, 1987, 1990; Felger, 1991): upper part is interbedded moderate-greenish-yellow and moderate-greenish-gray mudstone, and yellowish-gray, fossiliferous, platy, micritic limestone; mudstones become ashy near contact with overlying Hall Canyon Conglomerate; middle part is orange, brown, and gray conglomerate, mudstone, sandstone, and siltstone; lower part is green mudstone capped by yellowish-gray to light-olive-gray stromatolitic limestone; the formation is nearly 900 feet (275 m) thick (Clark, 1990).

Skinner Peaks area exposures: clastic lithofacies previously referred to as the "Tawny facies" (Zeller, 1949; Hardy and Zeller, 1953; Vogel, 1957; Millen, 1982; Felger, 1991; Weiss and others, 2003) composed of green, red, and variegated mudstone, and yellowish-tan coarse-grained sandstone, conglomerate, conglomeratic sandstone, and limestone; this lithofacies grades laterally eastward into the upper limestone member of the Green River Formation in the adjacent Chriss Canyon quadrangle (Millen, 1982; Weiss and others, 2003); sandstone near top of section contains reptilian(?) vertebrate fossils (Vogel, 1957; Felger, 1991); on the north side of Chriss Creek, unit has a heavy colluvial cover or locally a veneer of QTaf that has not been mapped separately; contact with overlying volcanic rocks not exposed, but present in Barton #1 well and is possibly conformable; directly overlies the Jurassic Arapien Shale at and north of Skinner Peaks; total thickness in subsurface from 1100 to 1300 feet (335-395 m).

Tc **Colton Formation** (lower Eocene) – Moderate-reddish-orange, moderate-reddish-brown, and pale-red mudstone, and lesser sandstone, conglomerate, and mottled carbonates (mottled pale red, reddish orange, and white with reduction spots); existence in West Hills follows Witkind and Weiss (1991), but is apparently absent in the Skinner Peaks area; conformable with overlying Green River Formation and underlying Flagstaff Formation; thickness from 0 to 300 feet (0-90 m); 414 feet (126 m) logged in the State #1 drill hole.

Tf **Flagstaff Formation** (lower Eocene to upper Paleocene) – As mapped in West Hills, gravish-yellow to pale-reddish-orange calcareous mudstone, sandstone, sandy limestone, limestone, and conglomerate (also see Lambert, 1976; Vorce, 1979); base not exposed; exposed thickness 525 feet (160 m), but total thickness in West Hills is about 650 feet (200 m) (Clark, 1987, 1990); 295 feet (90 m) logged in the Skinner #1 drill hole; locally absent near Skinner Peaks or included within map unit TKuu and/or TKul.

TKnh **North Horn Formation** (Paleocene? to Upper Cretaceous [Maastrichtian]) – Subsurface only (refer to cross section A-A'); includes gray to dark-gray mudstone, sandstone, sandy limestone, limestone, and conglomerate (also see Lambert, 1976; Vorce, 1979); base not exposed; exposed thickness 525 feet (160 m), but total thickness in West Hills is about 650 feet (200 m) (Clark, 1987, 1990); 295 feet (90 m) logged in the Skinner #1 drill hole; locally absent near Skinner Peaks or included within map unit TKuu and/or TKul.

unconformity?

TKu **TERTIARY-CRETACEOUS**

Tertiary-Cretaceous strata, undifferentiated – Corresponds to Colton, Flagstaff, and North Horn Formations; subsurface only (see Correlation of Map Units).

TKuu **Upper unit of undifferentiated Tertiary-Cretaceous strata** (middle to lower Eocene? to Upper Cretaceous?) – Conglomeratic outcrop belt on the northeast side of Skinner Peaks consisting of carbonate-clast conglomerate overlying red quartzite-clast conglomerate; the carbonate clast conglomerate predominantly pebbles and cobbles; gray to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

Tp1 **Member 1** – Upper part of unit (about 200 feet [60 m]) exposed at Painted Rocks consists of gray and light-bluish-gray volcanic conglomerate and tuffaceous sandstone; pebbles to boulder-size, dark to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuffaceous sandstone and mudstone with minor volcanic conglomerate and tuff; includes the Chicken Creek Tuff (38.61 ± 0.13 Ma; Clark, 2003) within the upper part of this unit; the Placid Oil Barton #1 well (table 1) contains an incomplete section of member 1 totaling about 1325 feet (405 m), which includes about 470 feet (143 m) of member 1 below the Chicken Creek Tuff; locally includes small outcrops of other members too small to map separately; total member 1 thickness unknown but estimated at 1700 feet (520 m).

Goldens Ranch Formation (upper? to middle Eocene) – Originally defined by Muessig (1953a, 1953b). Revised by Meibos (1983) to include only the lower part of Muessig's formation and divided into three members, named the upper Sage Valley Limestone, middle Hall Canyon Conglomerate, and lower Chicken Creek Tuff. We do not follow the suggested revision of the Goldens Ranch Formation by Witkind and Marvin (1989), and believe that a regional study of the volcanic rocks is needed to clarify outstanding stratigraphic issues. Refer to Correlation of Map Units. After Meibos (1983) and excluding the Sage Valley Limestone Member, which is not present in this quadrangle:

Ku (Ku, Kisp) **Hall Canyon Conglomerate Member** – Gray, poorly consolidated, cobble conglomerate and volcanic conglomerate; clast composition varies from quartzite-carbonate to volcanic; clasts are angular to subrounded pebbles to boulders in an ashy to sandy matrix; typically forms rubbly slopes; mapped in the northwest part of the quadrangle near Chicken Creek Reservoir where it directly overlies the Green River Formation; apparently laterally equivalent to lower portion of member 1 – formation of Painted Rocks; exposed thickness about 100 to 300 feet (30-90 m).

Tgc **Chicken Creek Tuff Member** – Grayish-pink to light-gray, poorly welded, vitric, dacitic ash-flow tuff with pumice lapilli, volcanic and quartzite rock fragments, and minor biotite; generally not well exposed; named for small exposure southwest of Chicken Creek Reservoir in section 19, T. 15 S., R. 1 W.; other exposures are in and near "Milky Wash"; query indicates uncertain designation; $^{40}Ar/^{39}Ar$ age of 38.61 ± 0.13 Ma from Sage Valley (Clark, 2003) supercedes K-Ar ages from Everenden and James (1964) and Witkind and Marvin (1989); source unknown; near the reservoir, located within the Hall Canyon Conglomerate Member from roughly 200 feet (60 m) or less above the top of the Green River Formation; near Painted Rocks and "Milky Wash" located stratigraphically within the lower part of member 1 (about 470 feet [145 m] above base); exposed thickness less than 100 feet (30 m), but total dip-corrected thickness in Placid Barton well is 170 feet (50 m).

Tg **Green River Formation** (middle Eocene) – West Hills exposures (see Clark, 1987, 1990; Felger, 1991): upper part is interbedded moderate-greenish-yellow and moderate-greenish-gray mudstone, and yellowish-gray, fossiliferous, platy, micritic limestone; mudstones become ashy near contact with overlying Hall Canyon Conglomerate; middle part is orange, brown, and gray conglomerate, mudstone, sandstone, and siltstone; lower part is green mudstone capped by yellowish-gray to light-olive-gray stromatolitic limestone; the formation is nearly 900 feet (275 m) thick (Clark, 1990).

Skinner Peaks area exposures: clastic lithofacies previously referred to as the "Tawny facies" (Zeller, 1949; Hardy and Zeller, 1953; Vogel, 1957; Millen, 1982; Felger, 1991; Weiss and others, 2003) composed of green, red, and variegated mudstone, and yellowish-tan coarse-grained sandstone, conglomerate, conglomeratic sandstone, and limestone; this lithofacies grades laterally eastward into the upper limestone member of the Green River Formation in the adjacent Chriss Canyon quadrangle (Millen, 1982; Weiss and others, 2003); sandstone near top of section contains reptilian(?) vertebrate fossils (Vogel, 1957; Felger, 1991); on the north side of Chriss Creek, unit has a heavy colluvial cover or locally a veneer of QTaf that has not been mapped separately; contact with overlying volcanic rocks not exposed, but present in Barton #1 well and is possibly conformable; directly overlies the Jurassic Arapien Shale at and north of Skinner Peaks; total thickness in subsurface from 1100 to 1300 feet (335-395 m).

Tc **Colton Formation** (lower Eocene) – Moderate-reddish-orange, moderate-reddish-brown, and pale-red mudstone, and lesser sandstone, conglomerate, and mottled carbonates (mottled pale red, reddish orange, and white with reduction spots); existence in West Hills follows Witkind and Weiss (1991), but is apparently absent in the Skinner Peaks area; conformable with overlying Green River Formation and underlying Flagstaff Formation; thickness from 0 to 300 feet (0-90 m); 414 feet (126 m) logged in the State #1 drill hole.

Tf **Flagstaff Formation** (lower Eocene to upper Paleocene) – As mapped in West Hills, gravish-yellow to pale-reddish-orange calcareous mudstone, sandstone, sandy limestone, limestone, and conglomerate (also see Lambert, 1976; Vorce, 1979); base not exposed; exposed thickness 525 feet (160 m), but total thickness in West Hills is about 650 feet (200 m) (Clark, 1987, 1990); 295 feet (90 m) logged in the Skinner #1 drill hole; locally absent near Skinner Peaks or included within map unit TKuu and/or TKul.

TKnh **North Horn Formation** (Paleocene? to Upper Cretaceous [Maastrichtian]) – Subsurface only (refer to cross section A-A'); includes gray to dark-gray mudstone, sandstone, sandy limestone, limestone, and conglomerate (also see Lambert, 1976; Vorce, 1979); base not exposed; exposed thickness 525 feet (160 m), but total thickness in West Hills is about 650 feet (200 m) (Clark, 1987, 1990); 295 feet (90 m) logged in the Skinner #1 drill hole; locally absent near Skinner Peaks or included within map unit TKuu and/or TKul.

unconformity?

TKu **TERTIARY-CRETACEOUS**

Tertiary-Cretaceous strata, undifferentiated – Corresponds to Colton, Flagstaff, and North Horn Formations; subsurface only (see Correlation of Map Units).

TKuu **Upper unit of undifferentiated Tertiary-Cretaceous strata** (middle to lower Eocene? to Upper Cretaceous?) – Conglomeratic outcrop belt on the northeast side of Skinner Peaks consisting of carbonate-clast conglomerate overlying red quartzite-clast conglomerate; the carbonate clast conglomerate predominantly pebbles and cobbles; gray to dark-pink, subangular to subrounded volcanic clasts with minor quartzite and carbonate clasts; most extensive exposures of member 1 in "Milky Wash" area of Vogel (1957) (not named on topographic base map; section 27 and 34, T. 16 S., R. 1 W.); also includes gray, brown, green, and blue tuff