APPENDIX A

Strike Valley Section
Cedar Mountain Formation measured by Douglas A. Sprinkel and Elizabeth Balls, August 22, 2000; Dakota Formation measured by Douglas A. Sprinkel and Scott K. Madsen, June 9, 2005
Along US Highway 191 between Steinaker and Red Fleet Reservoirs
N 40° 34.011' W 109° 29.930'

DESCRIPTION	THICKNESS (METERS)	
Dakota Formation	**Interval**	**Cumulative**
Unit 17 – Sandstone and mudstone. Sandstone is yellow-gray to white, fine- to-coarse-grained with lenses of pebble conglomerate, black chert, and white quartz grains, well-sorted, angular to subangular grains, thick-bedded, planar cross-bedding; interbedded mudstone, medium-gray, silty	16.6	24.4
Unit 16 – Mudstone, dark-gray, carbonaceous, contains petrified wood fragments, blocky, slightly silty; sample **SV042605-1** is composite sample collected from near base and from near the top of the mudstone unit (middle to late Albian swamp/deltaic); samples **SKM-1 through SKM-7** were collected about 1 meter apart along a trench dug the length of the unit (SKM-1 through SKM-4, middle to late Albian swamp/deltaic and SKM-6 and SKM-7, late Albian swamp deltaic)	5.8	7.8
Unit 15 – Sandstone and mudstone; sandstone is light-yellow-gray, fine-grained, white quartz grains, well-sorted, angular to subangular grains, non-calcareous, thin-bedded, low-angle cross-bedding; this unit has three subunits of lower resistant sandstone, middle friable sandstone that is orange-yellow stained near top, and upper resistant sandstone	1.8	2.0
Unit 14 – Shale, dark-gray, siliceous, fissile; sample **SV082200-2 and SV060806-2** (middle to late Albian; nearshore marine to estuarine)	0.2	0.2
Cedar Mountain Formation		
Unit 13 – Claystone, purplish-red to green, scattered sand-size grains, non-calcareous to slightly calcareous, blocky; some beds have mudcracks on bedding surface of less silty beds, slight popcorn texture in clay-rich zones, becoming more clay-rich near the top	8.9	35.7
Unit 12 – Siltstone, medium- to dark-gray, contains scattered rounded quartz and red chert sand-size grains, non-calcareous, blocky; some beds		

have mudcracks on bedding surface, slight popcorn texture in clay-rich zones, becoming more clay-rich near the top 9.8 26.8

Unit 11 – Sandstone, purplish-red and mottled yellow, calcareous. Purple sandstone is fine- to coarse-grained, poorly sorted, conglomeratic, yellow sandstone is coarse-grained, moderately sorted 0.5 17.0

Unit 10 – Mudstone that grades upward to silty mudstone and siltstone, dark-purplish-red, non-calcareous, poorly exposed 2.9 16.5

Unit 9 – Calcrete, light-gray to light-purple-gray, dense, irregular thickness 0.3 13.6

Unit 8 – Mudstone that grades upward to silty mudstone and siltstone, dark-purplish-red, non-calcareous, poorly exposed 2.2 13.3

Unit 7 – Limestone, medium-brown-gray with slight purple cast, dense, aphanitic with scattered small dendritic dark minerals (pyrolusite) some of which are oxidized to reddish brown 0.6 11.1

Unit 6 – Mudstone, dark-brownish-gray with slight purple cast, silty, non-calcareous, weathers as small thin plates, poorly exposed 3.1 10.5

Unit 5 – Calcrete (limestone), weathers out as nodules, light-gray calcareous with concentric rind about 5 mm in diameter 0.6 7.4

Unit 4 – Mudstone and siltstone, clay-rich, purple-gray; light-gray calcareous nodules about 5 mm in diameter, poorly exposed 1.8 6.8

Unit 3 – Calcrete (limestone), weathers out as nodules, light-gray calcareous with concentric rind about 5 mm in diameter 1.5 5.0

Unit 2 – Mudstone and siltstone, clay-rich, purple-gray; light-gray calcareous nodules about 5 mm in diameter, poorly exposed 2.3 3.5

Unit 1 – Calcrete, light-gray, chert, thick-bedded, forms prominent resistant ledge; unit thickens and thins as much as 2.5 m along strike 1.2 1.2

Basal Cedar Mountain Formation is an alteration zone of varicolored mudstone with chert pebbles and thin zone of chert pebble conglomerate below contact ..... not measured

Morrison Formation

Steinaker Reservoir Section
Measured by Douglas A. Sprinkel and Elizabeth Balls, August 21, 2000
Along US Highway 191 east of Steinaker Reservoir State Park entrance
N 40° 32.031' W 109° 31.204'

DESCRIPTION	THICKNESS (METERS)	
Dakota Formation	**Interval**	**Cumulative**
Unit 11 – Sandstone, yellow-gray to light-gray, fine- to coarse-grained with lenses of conglomerate, black chert, and white quartz grains, thick-bedded, cross-bedded	16.1	53.5
Unit 10 - Mudstone, dark-gray, carbonaceous	17.4	37.4
Unit 9 - Sandstone, yellow-gray to white, fine- to medium grained, well-sorted, thin-bedded, low-angle cross-bedding	4.2	20.0
Unit 8 - Mudstone, dark-gray, carbonaceous	15.0	15.8
Unit 7 - Sandstone, light-gray to yellow-gray, coarse- to medium-grained, subrounded, moderately sorted, calcareous, thin- to medium-bedded, small-scale trough cross-beds	0.8	0.8
Cedar Mountain Formation		
Unit 6 - Siltstone and mudstone; lower half is siltstone, greenish-purple-gray to varicolored beds that grade upward to mudstone, dark-gray, non-calcareous, hard	10.4	20.9
Unit 5 - Claystone (altered ash[?]), medium-gray, contains a few scattered small subhedral biotite and possibly hornblende crystals. Sample **SR-082100-1** (barren of palynomorphs) collected from the darkest gray interval within the unit	0.6	10.5
Unit 4 - Siltstone, purplish-red, calcareous, mostly covered with light-gray popcorn-textured soil. Top 0.3 m becomes grayish-purple with abundant calcareous nodules that are 15 to 30 cm in diameter	1.8	9.9
Unit 3 - Limestone and calcrete, light-gray to light-brown, nodular weathering, thick-bedded, brecciated in part by recrystallization and secondary calcite veining	3.3	8.1

Unit 2 - Claystone and mudstone, purple-gray, gypsum crystals on surface, somewhat of a popcorn texture, poorly exposed 4.0 4.8

Unit 1 – Calcrete, light-brown-gray, nodular weathering, thick-bedded, brecciated in part by recrystallization and secondary calcite veining, thickness varies along strike 0.8 0.8

Basal Cedar Mountain is an alteration zone of varicolored mudstone with chert pebbles and thin zone of chert pebble conglomerate below contact ..... not measured

Morrison Formation

Reef Quarry Section
Measured by James I. Kirkland and Gary Hunt, June 5, 6, and 11, 2010
east of pipeline and ornithopod quarry Utah paleontological locality # Un 153 on south flank of The Reef, north of Island Park road. Measured section runs from N 40° 29'53.4" W 109° 22'3.9" up gulley to north northeast to N 40° 29'58.6" W 109° 22'1.7".

DESCRIPTION	**THICKNESS (METERS)**	
Tununk Shale Member of Frontier Formation	**Interval**	**Cumulative**
Unit 37 – Shale, medium-olive-gray, laminated, sharp lower contact with Mowry Shale	not measured	
Mowry Shale		
Unit 36 – Shale, siliceous, dark-brownish-gray, finely laminated with abundant fish scales and associated fish skeletons on many horizons, numerous orange-stained partings that may represent volcanic ashes, not examined in detail, lower contact sharp and in direct contact with pebbles	27.90	138.76
Dakota Formation		
Unit 35 – Conglomeratic sandstone, 1-3 cm chert and lithic pebbles, bedded at 20-30 cm intervals, yellow-gray to light-gray, fine- to coarse-grained with lenses of conglomerate, black chert, and white quartz grains, thick-bedded, cross-bedded	1.52	110.86
Unit 34 – Sandstone, medium-yellow-brown fine- to medium-grained, low-angle cross-beds 20-30 cm thick,	7.51	109.34
Unit 33 – Interbedded sandy coal and coaly shale, dark-olive-gray to black, abundant plant debris	3.31	101.83
Unit 32 – Sandstone, pale-grayish-yellow, medium- to coarse-grained, cross-bed set 20-40 cm thick,	2.82	98.52
Unit 31 – Interbedded sandy coal and coaly shale, dark-olive-gray to black, abundant plant debris	2.07	95.70
Unit 30 – Bentonite, pale-brown to pale-yellow; zircons were extracted with 44 of 50 grains yielding U-Pb ages between 96-109 Ma and 21 grains yielding ages between 100 and 104 Ma giving a final age of **101.4 ± 0.4 Ma** at ± 2 sigma,	0.31	93.63

Unit 29 – Interbedded sandy coal and coaly shale, dark-olive-gray to black, abundant plant debris, ... 2.05 93.32

Unit 28 – Conglomeratic sandstone, light- to medium-orangish gray with dark orange fracture staining, chert and lithic pebbles, basal 20 cm poorly consolidated with pebbles 1-2 cm and 20 cm of erosional relief, abundant plant debris 1.11 91.17

Unit 27 – Interbedded shale, siltstone, and fine sandstone at 1-5 cm scale, dark-olive-gray overall with orangish-stained partings and buff-colored sandstone layers, abundant organic debris, ... 6.12 90.06

Unit 26 – Sandstone, light-orangish-brown, fine to medium grained, low-angle cross-beds 20-40 cm thick, soft; no better indurated than shalely units above and below, few organic debris rich layers, few vertical burrows, *Skolithos*? 5.25 83.94

Cedar Mountain Formation

Unit 25 – Claystone, light-olive-gray, smectitic, poorly exposed 1.65 77.79

Unit 24 – Claystone, dark-olive-gray, olive-green, and purplish, gray slightly gritty, slightly smectitic, ornithopod dinosaur skeleton (Un 153) collected 870 cm above base of unit about 100 meters to west of measured section associated with an approximately 1 meter interval of poorly developed iron stained nodules and iron stained fractures 14.64 76.14

Unit 23 – Gritty mudstone with iron-stained carbonate nodules ~ 10 cm thick by 20-50 cm across, olive-gray 1.80 61.50

Unit 22 – Calcrete in gritty mudstone, dark-purplish-red weathering to reddish-mauve; with sandstone of unit 18 forms prominent hogback near middle of Cedar Mountain Formation in this area 3.38 60.70

Unit 21 – Gritty mudstone with small carbonate nodules, light greenish-gray 1.95 57.32

Unit 20 – Sandy mudstone, light-gray, carbonate cemented, moderately resistant, tiny mm scale oncoids 0.80 54.37

Unit 19 – Gritty mudstone like unit 17 but with fewer and smaller gray pedogenetic and palustrine carbonate nodules, dark-purplish-gray 4.70 54.57

Unit 18 – Coarse sandstone, pale-brownish-gray, sublithic arenite, trough and planar cross-bed sets 40-70 cm thick, conglomeratic lenses of chert

pebbles and lithic fragments, laterally extensive across visible outcrop belt (> 1 mile) 4.05 49.87

Unit 17 – Gritty mudstone, dark-purplish-red weathering to pale-maroon, with abundant gray pedogenetic and palustrine carbonate nodules, horizons of large septarized nodules 30-50 cm across every 0.5 - 1.0 meter on average, chert replacement along septarian fractures common 18.50 45.78

Unit 16 – Sandy mudstone, dark-reddish-brown, scattered chert pebbles ~ 1 cm, few pedogenic slickensides 1.36 27.28

Unit 15 – Sandy mudstone dark-reddish-brown, with numerous gray chert laminae and chert-replaced roots, capped by 3 cm thick chert layer 0.45 25.92

Unit 14 – Interclast conglomerate, matrix medium-purplish-red mudstone with grain supported gray angular clasts <5 mm of mudstone and siltstone, lower contact irregular with as much as 10 cm of relief 0.10 25.47

Unit 13 – Pebbly mudstone, medium-purplish-red 0.49 25.37

Unit 12 – Gritty mudstone. Light-olive-gray, with rare pebbles and scattered ash laminae 0.56 24.88

Unit 11 – Mudstone, mottled medium-red and olive-gray, pebbly, rooted, with thin ash laminae at 6 cm up and at top 0.13 24.32

Unit 10 – Bentonite(?), pale-yellowish-gray, waxy 0.06 24.19

Unit 9 – Siltstone to gritty mudstone, mottled medium-red and olive-gray, contains about 1-cm-diameter chert pebbles, a bit better indurated than mudstones below, 2 cm pale-yellowish-green claystone at base 0.33 24.13

Unit 8 – Mudstone, dark-brownish-red, few scattered 1-cm-diameter chert pebbles 0.48 23.80

Unit 7 – Mudstone, medium-purplish-red, gritty, resting on waxy parting, may represent volcanic ash. 0.17 23.32

Unit 6 – Mudstone, dark-purplish-gray in lower 50 cm, becoming medium-greenish-gray to olive-gray with rare yellow mottles up section, middle 2 meters preserves common iron-stained mottling with granules of chert common from 2-3 mm and rarely to 5 mm 6.53 23.15

Morrison Formation (not completely measured)

Unit 5 – Indurated siltstone "porcellanite," similar to some altered volcanic ashes, zircons small (mostly < than 30 microns) and rounded, laser ablation for U-Pb dating revealed a suite of Upper Jurassic (153.05 ± 3.69 Ma at ± 2 sigma) and many Precambrian grains typical of the detrital zircon suites from the Morrison Formation; because of the fine grain size unit is reinterpreted as a loessite derived from wind deflation of the Morrison Formation	0.66	16.62
Unit 4 – Mudstone, medium-purplish-gray weathering to pale-mauve, some yellow staining on fractures, deeply weathered	4.5	15.96
Unit 3 – Claystone-mudstone, moderate-red with some minor yellow staining, smectitic(?), poorly exposed	5.83	11.46
Unit 2 – Conglomerate and coarse sandstone, somewhat fining upward with decreasing cross-bed set thickness from 50 to 20 cm, lithic and chert pebbles purple	1.74	5.53
Unit 1 – Smectitic claystone-mudstone, pale at base darkening to medium-dark-red, upper 10 cm bleached to pale-greenish-gray	3.79	3.79
Unit 0 – Coarse sandstone, pale-yellow-gray, trough cross-bedded, some lithic pebbles, lenticular (several meters across), forms hogback	not measured	

Pipeline Access Section
Measured by Scott K. Madsen and Dale Gray, October 24, 2006
West of pipeline access road, south of Island Park road, west nose of Split Mountain anticline; N 40° 28.360' W 109° 22.539'

DESCRIPTION	**THICKNESS (METERS)**	
Dakota Formation	**Interval**	**Cumulative**
Unit 10 – Sandstone, yellow-gray to light-gray, fine- to coarse-grained with lenses of conglomerate, black chert, and white quartz grains, thick-bedded, cross-bedded	4.9	37.2
Unit 9 – Covered and slope-forming interval; likely mudstone dominated	14.3	32.3
Unit 8 – Sandstone, yellow-gray to light-gray, fine-grained	1.0	18.0
Unit 7 – Mudstone, light- to dark-gray, carbonaceous, non-calcareous, contains root casts near base; sample **SKM102406-1** collected 15 cm above base of unit and **SKM102406-2** collected 50 cm above base of unit (both samples barren)	3.3	17.0
Unit 6 – Sandstone, light-brown (buff) to greenish-gray, fine-grained, non-calcareous	1.2	13.7
Unit 5 – Mudstone, light-brown to light-gray, silty, non-calcareous, bentonitic; section was too wet to trench but looked to contain carbonaceous material (one test hole revealed sandstone stringer)	3.7	12.5
Unit 4 – Sandstone, light-brown, light-gray to gray, weathers light-brown to pink, non-calcareous, fine-grained, bed thickens laterally to west from 0.5 to 1.0 m (this sandstone unit caps previously sampled mudstone interval)	0.3	8.8
Unit 3b – Shale, dark-gray to black, fissile, **PA060806-3** collected at top of unit (middle to late Albian nearshore marine to estuarine)	1.0	8.5
Unit 3a– Mudstone, black, highly carbonaceous; two samples collected from this unit; **PA060806-1** collected at base of unit (middle Albian to early late Albian swamp-deltaic), **PA060806-2** collected about 3 m from base of unit (middle to late Albian nearshore marine to estuarine)	5.0	7.5
Unit 2 – Sandstone, light-brown, poorly exposed (was not trenched because section was too wet), untrenched surface has pebbly and lignitic appearance	1.0	2.5

Cedar Mountain Formation (not completely measured)

Unit 1 – Mudstone, purple 1.5 1.5

Dinosaur National Monument - DNM 16 (sauropod locality)
Measured by Scott K. Madsen, October 2001
Started just west of DNM 16 on north side of hogback at highest white sandstone in Morison Formation; strike 68° W, dip 64° south

DESCRIPTION	THICKNESS (METERS)	
Dakota Formation	**Interval**	**Cumulative**
Unit 42 – Sandstone, yellow-gray to light-gray, fine- to coarse-grained with pebble conglomerate beds, well sorted, normally graded, planar cross-beds, limonitic staining is greater at the top of the unit; uppermost sandstone beds also exhibit siliceous deformation veining	12.8	30.8
Unit 41 – Mudstone, dark-gray to black, yellow surface stains, very deeply weathered slope, forms slope between unit 40 (lowest Dakota Formation bed) and main Dakota Formation ledge	14.0	18.0
Unit 40 – Sandstone, weathers yellow-gray, with light rust color, white on unweathered surface, non-calcareous, well-sorted, very fine grained, steeply inclined cross-beds; forms yellow-weathering blocky slopes and has same features as main Dakota Formation ledge above. Contact appears scoured at top and bottom	2.0	4.0
Unit 39a – Mudstone, very dark gray to black, deeply weathered, very carbonaceous with sandstone stringers; similar in lithology and stratigraphic position to unit 3 of the Pipeline Access section that yielded significant palynology; originally included in uppermost 2 m of unit 39	2.0	2.0
Cedar Mountain Formation		
Unit 39 – Siltstone and mudstone; siltstone, very light gray, light-brown (sometimes green), slope weathers pale-yellow and light-gray; bottom 2 m not trenched due to debris; lower 4.6 m is highly calcareous alternating siltstone, mudstone, and calcareous nodule beds (calcareous beds are mostly thin 1 to 5 cm thick and rare); mudstone, light-olive-gray and pale-olive, yellow surface stains and blotches; calcareous beds are light-gray; upper 4 m of slope is generally more yellow, deeply weathered, and hard to trench; yellow blotches on surface; rare gypsum on surface (not found in unit below); no "gastroliths" and few nodules; popcorn weathering	13.6	68.9

Section offset approximately 100 m east above unit 38 for better exposures.

Unit 38 – Sandstone and mudstone, interbedded; deeply weathered, yellowish subsoil, some sandstone stringers; top 2 m slightly more resistant with multiple sandstone layers (approximately 30 m west of section beds are approximately 50 cm thick), discontinuous; top bed extensively bioturbated, burrows average approximately 1 cm in diameter, some burrows penetrate beds	8.4	55.3
Unit 37 – Sandstone, light-gray (unweathered) that weathers brown, matrix dull white, dark and light grains giving "salt and pepper" appearance, minor red grains, subrounded to angular, well-sorted, some fining upward, laminae approximately 2 mm thick, cross-beds appear as climbing ripples; unit contains three prominent ledge-forming beds with many minor sandstone lenses; bottom ledge-forming sandstone bed contains bone and a base that undulates with approximately 20 cm of relief	1.4	46.9
Unit 36 – Mudstone and siltstone, brown-gray, highly calcareous, deeply weathered, numerous gray nodules (0.5 to 2 cm in diameter); includes some white calcareous nodular (up to 50 cm in diameter) irregularly shaped blocks, spongy, and weathered	4.5	45.5
Unit 35 – Clastic zone of calcrete, sandstone, and claystone (rip-up), mostly yellowish-gray (fresh surface); weathered, trench samples are mostly 1 cm in diameter or smaller fragments with nodules up to 5 cm in diameter, most nodules are white on freshly broken surface; blocky to angular, weakly bedded	2.5	41.0
Unit 34 – Calcareous zone, basal calcrete, dark-gray to black, 1 cm in diameter pebbles and calcareous nodules in matrix overlain by white to light-yellow, weakly calcareous, nodule zone; upper 40 cm is more resistant, compact calcrete, weathers light-gray with dull rust color, contains calcareous nodules (0.5 to 1 cm in diameter) and gray chert (1 cm in diameter), possible rhizoids or burrows; "gastroliths" on slope	0.9	38.5
Unit 33 – Siltstone, mostly pale-blue with mottled red and blue grading to blue at bottom, coarse at top, very calcareous, calcite veins, deeply weathered and hard to trench in. Top 1 m contains calcareous nodules and thin (5 cm thick) calcrete beds	2.2	37.6
Unit 32 – Mudstone, red, silty, deeply weathered, weakly calcareous	1.0	35.4
Unit 31 – Calcrete and calcareous nodules, some weakly calcareous; looks like sandstone altered to calcrete bed, many inclusions of clastics, includes red mudstone rip-ups, small calcareous nodules 1 to 3 cm in diameter; contains possible root casts or rhizoliths	0.5	34.4

Unit 30 – Mudstone and siltstone, red with some blue mottling, weakly calcareous, 15-cm calcareous pillows, fairly deeply weathered.....2.7 33.9

Unit 29 – Sandstone, white (weathered surface), red (unweathered surface), very calcareous, some 1-mm sized grains with white halos.....0.1 31.2

Unit 28 – Sandstone, white-red, fine-grained, very calcareous.....0.1 31.1

Unit 27 – Sandstone, brown-gray, mix of clear, dark-red, orange grains, dominantly angular to rounded, fine-grained, highly calcareous; bottom 75 cm is very coarse grained to gravelly with same color grains as above; contains green mudstone rip-ups 0.5 to 1.5 cm in diameter; top of bed has 4- to 5-sided polygons with rounded "pillow-like" surfaces with abundant burrows; lower beds contain low-angle and trough cross-beds showing east-directed flow; east and west of section this unit is mostly slope covered with abundant "gastroliths" of mostly shades of red with some gray and white, up to 2 cm in diameter; abundant calcareous nodules (0.5 to 7 cm in diameter) weathered out on slope; lowest part of unit contains 75 percent calcareous nodules, white and gray, clasts 1 to 3 cm in diameter, also minor black and red rounded pebbles; some beds sit directly on calcrete bed composed of same material as nodules in sandstone above, sauropod skull (*Abydosaurus mcintoshi*) excavated from this unit 2.4 31.0

Unit 26 – Calcrete and siltstone interbedded; calcrete, dull white, pale-gray-blue (unweathered surface), weathers dull red-brown; no spar, some nodules in beds 5 cm in diameter and possible relic bedding; calcite crystals and calcite veins found in bed 20 m west; calcrete beds approximately 20 cm thick; minor siltstone, fissile, blocky, deeply weathered, brown, drab green, calcareous; bottom 20 to 30 cm is pale blue, well indurated, fine-grained, weakly calcareous, blocky weathering.....2.5 28.6

Note: south-facing slope above contains almost no "gastroliths" or quartz-chert pebbles; north-facing slope in lower section (below massive basal calc beds) has much silcrete debris but could not find in place; some chalcedony/silcrete scattered on south-facing slope to the west.

Unit 25 – Mudstone, mottled medium-gray, light-green, and minor amount of red, calcareous, blocky to fissile, carbonate nodules 0.5 to 20 cm in diameter 2.6 26.1

Unit 24 – Calcrete, weathers tan, light-rusty-brown, mostly micrite, much white calcite crystal; no visible grains; irregular base that is pipe-shaped (dimensions of 20 x 40 cm); top is irregular with medium-grained mudstone, mottles, calcareous, possible relict bedding traces 1.7 23.5

Section offset approximately 7 m west

Unit 23 – Mudstone, dark-gray-purple, some green mottling, fissile, contains calcareous nodules up to 7 cm in diameter 0.8 21.8

Unit 22 – Mudstone, dark gray, non-calcareous 0.5 21.0

Unit 21 – Mudstone, dull green, very calcareous 0.4 20.5

Unit 20 – Calcrete, white to light-gray, with minor siltstone, weathers brown to rusty red; minor calcite crystals and calcite veins 0.6 20.1

Unit 19 – Mudstone, light-green-gray, non-calcareous; interbedded with minor sandstone, red-green, and calcrete; calcrete beds up to 8 cm thick and located at 9 cm, 144 cm, 200 cm above base of unit; carbonate nodules 0.5 to 20 cm in diameter throughout unit 2.6 19.5

Section offset approximately 3 m west to ridge top

Unit 18 – Calcrete, white to light-brown (unweathered), dense crystalline cores, sandy rinds with box-like structures, chemical weathered in relief; most calcrete beds form vertical pipe-like structures that protrude down into underlying red mudstone unit; an excavated structure in lowest part of unit extends 43 cm down into mudstone and approximately 5 cm wide; others are 32 cm wide at top and more than 36 cm deep; holds up ridge 5.4 16.9

Unit 17 – Mudstone, red, blocky to fissile, silty to sandy, minor red, subangular quartz grains, approximately 1 to 3 cm in diameter; unit contains calcrete beds (three beds 4 to 10 cm thick), pale-blue, calcareous nodules 0.5 to 7 cm in diameter throughout unit, some calcite veins, 3 mm thick; top is in-filled with protruding pipe-like structures of unit 18 ..... 2.5 11.5

Unit 16 – Calcrete, blue-gray, weathers rusty brown, lower part is very crystalline and bottom 1 to 5 cm is very platy; may thicken east; base of unit is sandstone, very minor red with blue-colored clay clasts, fine-grained; upper 25 cm is poorly cemented sandstone, bright green, very fine-grained, non-calcareous 0.5 9.0

Unit 15 – Sandstone and siltstone, maroon grading upward to pale-bluish-green, fine-grained, minor interbedded mudstone, calcareous; abundant grains (up to 1 cm in diameter), mostly red, orange, black, and brown, subangular; green clay rip-up clasts; abundant limonite especially in calcareous beds and limonite rims around nodules and grains 1.7 8.5

Unit 14 – Calcrete, white to pale-blue, mostly blocky weathering and with some nodules; flat bottom to bed, irregular top, grades upward to siltstone; fairly uniform with minor lithic grains, dull red, black, and gray, 0.5 to 5

mm in diameter, angular to rounded; some rounded crystalline infilling may be root or burrow traces; first major calcrete bed in Cedar Mountain Formation.....0.3 6.8

Strike and dip on bottom of calcareous bed, N74W, dip 63° S

Unit 13 – Mudstone, mottled drab red, green, and gray, well indurated, contains very minor quartz grains, some clays, calcareous.....0.8 6.5

Unit 12 – Mudstone, green, well indurated; minor quartz grains (red, orange, black), 0.5 mm in diameter; scattered silcrete/calcrete nodules (1 to 4 cm in diameter) and stringers 0.9 5.7

Unit 11 – Mudstone, green, with abundant quartz grains, mostly black, drab red, gray, rounded to subangular, some grains 2.5 to 3 cm in diameter.....0.2 4.8

Unit 10 – Calcrete, silcrete, and mudstone (3 alternating layers); mudstone, green, like unit 9 but contains no limonite; calcrete, rusty brown, black, and green (unweathered), top and bottom with clay contacts, irregular thickness (approximately 3 to 4 cm thick), slightly calcareous; silcrete.....0.9 4.6

Unit 9 – Mudstone and siltstone, light-green with black and red grains, abundant limonite 0.2 3.7

Unit 8 – Siltstone, pale-green, blocky; sandstone, black, red, and gray grains, rounded, up to 1 cm in diameter 0.4 3.5

Unit 7 – Mudstone, gray to pale-green, becoming green upward; interbedded siltstone; some limonite; scattered non-calcareous nodules to 8 cm in diameter; abundant clay rip-ups, green and purple; upward becoming silty with sandstone, clear, red, and gray grains, 1 to 5 mm in diameter 2.7 3.1

Unit 6 – Mudstone, green, blocky with abundant limonite; minor sandstone, non-calcareous; contact at base of "alteration zone" 0.4 0.4

Morrison Formation (Morrison was not completely measured)

Unit 5 – Mudstone, gray-green, silty to sandy 0.3 3.7

Unit 4 – Sandstone, light-green, coarse-grained, angular to subangular, white quartz and plagioclase grains, 1 to 5 mm in diameter 0.3 3.4

Unit 3 – Mudstone, light-green, silty, non-calcareous 0.2 3.1

Unit 2 – Mudstone, purple-medium green, blocky weathering	1.4	2.9
Unit 1 – Sandstone, white to pale-blue, fine-grained, grades to purple below and purple above, non-calcareous	1.5	1.5

UPbICP and Related Excel Workbooks: Software for Calculating U-Pb Zircon Ages and Presenting U-Pb Data Obtained by LA-ICP-MS

Raymond A. Donelick
Apatite to Zircon, Inc.
1075 Matson Road
Viola, Idaho 83872-9709 U.S.A.
donelick@apatite.com

1. UPbICP Data Processing Software

1.1. Background

Two useful studies that demonstrate methods of U-Pb dating of zircon based on laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS) are Chang et al. (2006) and Gehrels et al. (2008). The U-Pb data analysis methods described here and the computer program UPbICP (written using Visual Basic 6.0, Microsoft Corporation) that implements them were developed by Apatite to Zircon, Inc. (A2Z) and remain the intellectual property of A2Z. A2Z offers no guarantees that the calculations performed by UPbICP are flawless but a good faith effort continues to be made to refine and verify the accuracy of the calculations. Original output data from the LA-ICP-MS system remain the property of the party that owns the sample from which they were obtained and UPbICP does not modify the original data.

Dissemination and publication of the description provided here and the ideas and concepts contained herein are prohibited without the written permission of A2Z.

1.2. Definitions

A LA-ICP-MS session is characterized by a series of zircon ***spots*** (usually one per grain; **Figure 1**) that are ablated by a laser beam resulting in ***laser ablation pits***. During laser ablation of a single spot, the ablated material is promptly transported from the deepening laser ablation pit into a mass spectrometer for a ***scan*** of the relative abundances of each target isotope. A series of scans is performed for each spot analyzed. A spot may be ablated for a zircon ***age standard*** having a known or accepted age, or for a zircon ***unknown*** having an unknown age. Spots for age standards are strategically interspersed with spots from unknowns. For each spot, measured ***background*** intensities for each isotope are obtained from scans performed prior to unblocking the laser. After the laser is unblocked, measured ***background+signal*** intensities are obtained from the remaining scans. During data analysis, ***signal*** intensities are calculated by subtracting respective background intensities from the background+signal intensities by a process referred to as ***background correction***.

The U-Pb ages for a scan are deemed ***concordant*** if the 207Pb/235Uc (where ***235Uc*** is 235U calculated on the basis of measured 238U), 206Pb/238U, and 207Pb/206Pb ages overlap each other at the 2σ level. U-Pb ages that do not meet this criterion are deemed ***discordant***. The ***preferred age*** for a spot is the U-Pb age for that spot deemed by UPbICP to be of highest quality where first preference is given to the

most precise weighted mean U-Pb age of the concordant scans (***Pb/U wmean conc.*** for 206Pb/238U; ***Pb/Pb wmean conc.*** for 207Pb/206Pb) and second preference (used for spots exhibiting only discordant scans) is given to the most precise intercept age between the discordia and the concordia (***lower int.*** for lower intercept; ***upper int.*** for upper intercept).

1.3. Data Collection and Age Calculation

Zircon grains prepared only for LA-ICP-MS U-Pb age dating are mounted in epoxide resin, cured at ~50°C for 3-10 hours, and polished to expose internal grain surfaces using 3.0μm Al_2O_3 and 0.3μm Al_2O_3 slurries on dedicated lap wheels. The polished zircon grains are then cleaned in reagent-grade 5.5M HNO_3 for 20s at 21°C. Zircon grains prepared for both zircon fission track (ZFT) and U-Pb age dating are mounted in FEP TEFLON and polished to expose internal grain surfaces using 3.0μm Al_2O_3 and 0.3μm Al_2O_3 slurries on dedicated lap wheels. Natural zircon fission tracks are revealed for optical viewing by immersing the grain mounts in an eutectic melt of NaOH+KOH (at ca. 220°C) for 12-72 hours. The polished and etched zircon grains are then cleaned in reagent-grade 48% HF for 15 minutes at 23°C.

Details concerning the LA-ICP-MS apparatus and data acquisition parameters routinely used by A2Z are summarized in **Table 1**. Images of spot 17103b_20 are shown in **Figure 1** (the data in **Figures 2-15** except for **Figure 13b** pertain to spot 17103b_20).

A2Z typically collects data from 255 scans per spot during ~40 seconds of acquisition time. Data for the isotopes used for U-Pb age calculation are collected during the first 250 scans. Data for 28Si and 91Zr are collected during the last 5 scans to verify zircon identity and to determine whether or not the laser has either dislodged the zircon grain from its mounting medium or passed through the zircon grain into the mounting medium. **Figure 2** presents 206Pb, 207Pb, and 238U data for spot 1713b_20.

The laser is blocked from hitting the target zircon surface during the first ~ 45-55 scans and background intensities are recorded for the isotopes used to calculate U-Pb ages (background intensities are not collected for 28Si and 91Zr because background+signal intensities provide sufficient information for these isotopes). Background correction of the background+signal intensities for each isotope and scan and spot is performed by first fitting a polynomial equation to the background intensity versus scan number data (routine LFIT from Press et al., 1992):

$$B_{Fi} = \sum_{j=1}^{n+1} a_j^{j-1} i \qquad \textbf{Equation 1}$$

where:
i = background intensity scan number
B_{Fi} = fitted background value at scan i
n = order of polynomial; maximum=2
a_j = fitted coefficient j

The polynomial is regressed from a selected range of scans (scans 1-40 are often used; linear regression is often used). The two highest background intensity values from the selected range of scans are eliminated prior to polynomial fitting to avoid outlier values (which are periodically observed; note the 207Pb outliers at scans 102 and 190 in **Figure 13b**; such spurious values may occur for any isotope at any scan and any spot). Acceptable polynomials are limited to those for which B_{FNlast} (**Equation 1** where i = *Nlast*, the last background intensity scan used for background correction) represents the minimum fitted value over the range of i values.

When the laser is unblocked and begins to ablate the surface of the zircon, background+signal intensities for the isotopes used for U-Pb age calculation are recorded through scan 250. For each isotope, a sum of multiple Gaussian equations is fitted to the background+signal intensity versus scan number data from a selected range of scans (scans 90-250 are often used; initial scan 90 is chosen to avoid complications that arise during the transition between background intensity data and background+signal intensity data):

$$BS_{Fi} = \sum_{j=1}^{N_{Gauss}} g_{1j} \exp\left\{-\left[\frac{\left(i - g_{2j}\right)}{g_{3j}}\right]^2\right\} \qquad \textbf{Equation 2}$$

where:
i = background+signal intensity scan number
BS_{Fi} = fitted background+signal intensity at scan i
j = Gaussian equation number
N_{Gauss} = number of Gaussian equations summed
g_{1j} = amplitude of Gaussian equation j
g_{2j} = mean of Gaussian equation j
g_{3j} = standard deviation of Gaussian equation j.

The sum of Gaussian equations is comprised of a sufficient number of individual Gaussian equations to adequately fit most complicated background+signal intensity versus scan number data structures (N_{Gauss} = 10 is usually the preferred number for **Equation 2**; numbers from 5-10 have been tested). Fitting is performed using the Levenberg-Marquardt method and chi-squared minimization (routine MRQMIN from Press et al., 1992). The χ^2 value from the Gaussian fitting is given by:

$$\chi^2 = \sigma_{SD}^2 N_{scan} \qquad \textbf{Equation 3}$$

where:
χ^2 = chi-squared statistic
σ_{SD} = standard deviation of ($BS_{Mi} - B_{Fnlast}$) about S_{Ci} (**Equation 6a** below)
N_{scan} = number of signal intensity scans

The degrees of freedom of the fitted sum of Gaussian equations are given by:

$$\nu = N_{scans} - 3N_{Gauss}$$ **Equation 4**

where:
ν = degrees of freedom.
N_{Gauss} = number of Gaussian equations summed
N_{scan} = number of signal intensity scans

The probability of a worse fit (Q in **Equation 11** below) is calculated based on the number of degrees of freedom (**Equation 4**) and the χ^2 value from the Gaussian fitting (**Equation 3**) (routine GAMMQ from Press et al.,1992). A data filtering procedure is used by which all background+signal intensity values are first fitted and then those background+signal intensity values that fall outside of 2 standard deviations from their fitted counterparts are eliminated (e.g., note the spurious background+signal intensity values at scans 102 and 190 in **Figure 13b**). The process of fitting a sum of multiple Gaussian equations is performed a second time using the filtered background+signal intensity values

Background correction of the background+signal intensity data for an isotope is performed by subtracting the background intensity at the last background scan (B_{F40} for a scan range of 1-40; **Equation 1**) from the fitted background+signal intensity at each background+signal scan.

$$S_{Ci} = BS_{Fi} - B_{FNlast}$$ **Equation 5**

where:
i = background+signal intensity scan number; signal intensity scan number after background correction
S_{Ci} = calculated signal intensity at signal intensity scan i
BS_{Fi} = fitted background+signal intensity at scan i (**Equation 2)**
$Nlast$ = scan number of the last background intensity scan
B_{FNlast} = fitted background intensity at background intensity scan $Nlast$ (**Equation 1** where $i = Nlast$)

If S_{Ci} <0 due to this subtraction, S_{Ci} is set to 0. The standard error of the calculated signal intensity for each isotope for each spot is assumed constant for all regressed scans; this standard error is set equal to the standard error of the unfitted background+signal intensities about their fitted counterparts:

$$\sigma_{SD} = \left(\left(\frac{1}{N_{scan}} \right) \sum_{i=1}^{N_{scan}} \left(\left(BS_{Mi} - B_{FNlast} \right) - S_{Ci} \right)^2 \right)^{1/2}$$ **Equation 6a**

where:
i = background+signal intensity scan number; signal intensity scan number after background correction
σ_{SD} = standard deviation of ($BS_{Mi} - B_{Fnlast}$) about S_{Ci}
N_{scan} = number of signal intensity scans

S_{Ci} = calculated signal intensity at signal intensity scan *i* (**Equation 5**)
BS_{Mi} = measured background+signal intensity at scan *i*
Nlast = scan number of the last background intensity scan
B_{FNlast} = fitted background intensity at background intensity scan *Nlast* (**Equation 1** where *i* = *Nlast*)

$$\sigma_{SE} = \frac{\sigma_{SD}}{\nu^{1/2}} \qquad \textbf{Equation 6b}$$

where:
i = background+signal intensity scan number; signal intensity scan number after background correction
σ_{SE} = standard error of ($BS_{Mi} - B_{Fnlast}$) about S_{Ci}
σ_{SD} = standard deviation of ($BS_{Mi} - B_{Fnlast}$) about S_{Ci} (**Equation 6a**)
N_{Gauss} = number of Gaussian equations summed
N_{scan} = number of signal intensity scans

After scan 250, five scans of background+signal intensities are recorded for 28Si and 91Zr. For a given spot, the median value for each isotope from the values recorded is used to avoid spurious values. Under typical analytical conditions, zircon yields a 91Zr/28Si ratio of median background+signal intensities in the range 0.5-2.0. For a whole LA-ICP-MS session, the 91Zr/28Si ratios of median background+signal intensities exhibit a standard deviation in the range 0.1-0.2. If the laser has passed through a zircon into the mounting medium, this ratio decreases due to the faster decay rate of the 91Zr background+signal intensity relative to the decay of the 28Si background+signal intensity. A median 91Zr/28Si ratio for all primary age standard spot ratios for a session is determined and the standard deviation about this median (excluding the two minimum and two maximum values to avoid any possible problematic spots) is calculated. Only secondary age standard spots and unknown spots exhibiting 91Zr/28Si ratios within 3 standard deviations of the median of the medians for the primary age standard are considered acceptable spots for U-Pb age calculation. The mineral baddeleyite yields 91Zr/28Si ratios of median background+signal intensities in the range 20-40. Other non-zircon mineral species encountered yield 91Zr/28Si ratios less than 0.2.

The common Pb correction in UPbICP is performed as described in **Appendix 1**. This correction is not routinely invoked, however. Instead, spots are rejected from further data analysis if sufficient 204Pb is encountered in at least one scan (see section **2.2.1** below).

UPbICP calculates U-Pb ages using the following conventions (Steiger and Yäger, 1977):

235U/238U = 1 / 137.88
λ_{235U} = 0.00098485 Ma^{-1}
λ_{238U} = 0.000155125 Ma^{-1}
λ_{232Th} = 0.000049475 Ma^{-1}

Two types of age standards are used during each LA-ICP-MS session (**Table 2**). A ***primary age standard*** is used to characterize isotopic fractionation factors for all scans and all spots for which U-Pb age calculations are to be performed. A series of ***secondary age standards*** are used to correct fractionation factors from the primary age standard for the effect of relative α-radiation damage on laser penetration rate at each spot. The primary age standard, usually FC-1 obtained from the University of Minnesota student Geology Club (Paces and Miller, 1993), is analyzed in pairs at the beginning and end of each session and every 20-30 spots during a session. The isotopic ratios used to calculate fractionation factors for primary age standard FC-1 are (**Table 2**; obtained from the University of Minnesota student Geology Club; Paces and Miller, 1993):

207Pb/235U = 1.951607 (accepted)
206Pb/238U = 0.185877 (accepted)
207Pb/206Pb = 0.076149 (accepted)
208Pb/232Th = 0.055878 (assumed)

Fractionation factors for the isotopic ratios 207Pb/235Uc, 206Pb/238U, and 207Pb/206Pb are calculated for every signal scan and spot position for which primary age standard data are measured. For 206Pb/238U (similar equations can be written for 207Pb/235Uc and 207Pb/206Pb):

$$FF_{C206/238i,k} = \frac{\left(\frac{^{206}Pb}{^{238}U}\right)_{accepted}}{\left(\frac{S_{C206i,k}}{S_{C238i,k}}\right)} \qquad \textbf{Equation 7}$$

where:
i = signal intensity scan number
k = primary age standard spot position in session
$FF_{C206/238i,k}$ = 206Pb/238U fractionation factor at signal intensity scan i based on individual calculated signal intensities and spot position k
$S_{C206i,k}$ = calculated 206Pb signal intensity at signal intensity scan i and spot position k (**Equation 5**)
$S_{C238i,k}$ = calculated 238U signal intensity at signal intensity scan i and spot position k (**Equation 5**)

At fixed signal scan number i, a polynomial equation is fitted to spot number (abscissa) versus fractionation factor (ordinate) for each isotopic ratio over the range of primary age standard spot positions throughout the session. For 206Pb/238U (similar equations can be written for 207Pb/235Uc and 207Pb/206Pb):

$$FF_{F206/238i,k} = \sum_{j=1}^{n+1} f_{i,j}^{\,j-1} k \qquad \textbf{Equation 8a}$$

where:
k = spot number in session
i = fixed signal scan number
$FF_{F206/238i,k}$ = fitted 206Pb/238U fractionation value at signal scan i and spot number k
n = order of polynomial; maximum 4
$f_{i,j}$ = fitted coefficient j for signal scan i

$$\sigma_{t206/238} = \left(\left(\frac{1}{N_{PAS}} \right) \sum_{j=1}^{N_{PAS}} \left(t_{206/238j} - \frac{\sum_{j=1}^{N_{PAS}} t_{206/238j}}{N_{PAS}} \right)^2 \right)^{1/2} \qquad \textbf{Equation 8b}$$

where:
j = primary age standard number
$\sigma_{t206/238}$ = standard deviation of 206Pb/238U ages for the primary age standard
N_{PAS} = number of primary age spots
$t_{206/238j}$ = 206Pb/238U age for primary age standard j

A data filtering procedure is used whereby all points are fitted and those points that fall outside of 2 standard deviations (function STEYX; Excel 2007) are eliminated and the remaining points are re-fitted and the resulting regression used. Relative errors of the 207Pb/235Uc, 206Pb/238U, and 207Pb/206Pb fractionation factors are calculated as follows: a) the primary age standard is treated as an unknown and ages for each isotopic ratio are calculated for each spot, b) the mean preferred ages and their respective standard deviations for all spots for each isotopic ratio are calculated, c) the relative error of each fractionation factor is set equal to its respective standard deviation divided by its respective mean preferred age, and d) fractionation factor relative errors are assumed constant throughout the session.

Fractionation factors are adjusted for variable laser penetration rate (and hence variable ablation pit depth at any given scan number from spot to spot) due to variable α-radiation damage using the following scheme:

- A secondary age standard is analyzed in pairs along with the primary age standard (recent sessions use TR as this secondary age standard; see **Table 2**). An additional series of secondary age standards is analyzed at the beginning and again at the end of a LA-ICP-MS session. Recent sessions include 10 spots each (5 near the beginning and 5 near the end of each session) for four additional secondary age standards (recent sessions use FC-5z, MD, IFCan, and TEM2; see **Table 2**). Preferred ages are calculated for each secondary age standard spot using uncorrected fractionation factors from the primary age standard. Relative α-radiation damage is calculated for each spot by integrating *function dAlphaEnergy* (**Appendix 2**) over all signal scans.

- For each secondary age standard, the weighted mean of preferred ages from the youngest, concordant population of spots is calculated and the mean relative α-radiation damage for the population of spots is calculated. Weighted mean ages calculated for the secondary age standards using uncorrected fractionation factors from the primary age standard are observed to deviate linearly from their accepted ages as a function of the relative α-radiation damage, with fractionation being greater in older, more α-radiation damaged zircons (**Figure 3**). Presumably, this is due to greater penetration rates of the laser into softer, more highly α-radiation damaged zircon. To permit the correct ages of the secondary age standards to be calculated, a fractionation factor correction is applied as shown in *sub AlphaEnergyFF* in **Appendix 2** and discussed further in **Section 2.5**.

U-Pb preferred ages are calculated for a spot using the following scheme:

- For age standards, UPbICP characterizes signal intensities of each isotope at each scan and each spot for which U-Pb age calculations are to be performed. Fractionation factors from the primary age standard are corrected for α-radiation using secondary age standards.
- For unknowns, UPbICP characterizes signal intensities of each isotope at each scan and each spot for which U-Pb age calculations are to be performed. As such, each scan can be considered an unique U-Pb analysis and each spot can be considered composed of a series of downhole U-Pb analyses.
- The 207Pb/206Pb fractionation factor for the primary age standard pertaining to each scan is forced to be 1 at all scans and all spots.
- The combination of isotopic ratios for unknowns and fractionation factors from age standards at each scan and each spot permits the calculation of three ages for each scan for each unknown; these are: 1) 207Pb/235Uc age, 2) 206Pb/238U age, and 3) 207Pb/206Pb age.
- A discordia (207Pb/235Uc abscissa, 206Pb/238U ordinate) can be characterized for each spot from an unknown. Ages corresponding to the upper and lower intercepts of the discordia with the concordia are calculated by: a) linear regression of the (207Pb/235Uc, 206Pb/238U) points for the spot and b) Monte Carlo simulation of the discordia using 1000 randomly generated lines based on the regression-derived slope and intercept values and their respective errors. The quality of these ages increases with increasing number of discordant scans, and increasing range of discordance of the scans.
- *Preferred age based on the weighted mean isotopic ratios of concordant scans*: This strategy seeks the weighted mean of all isotopic ratios from concordant scans. A scan is concordant if its 207Pb/235Uc age, 206Pb/238U age, and 207Pb/206Pb age overlap each other at the 2σ level (see below). According to this strategy, if at least one scan is concordant, the weighted mean of the 207Pb/235Uc ratios from all concordant scans is used to calculate the 207Pb/235Uc age (function *Age207235* in **Appendix 3**), the weighted mean of the 206Pb/238U ratios from all concordant scans is used to calculate the 206Pb/238U age (function *Age206238* in **Appendix 3**), and the weighted mean of the 207Pb/206Pb ratios from all concordant scans is used to calculate the

207Pb/206Pb age for the spot (function *Age207206* in **Appendix 3**). The error for each of these ages is based on the weighted mean error of its respective weighted mean ratio combined with the relative error of its respective fractionation factor. For 206Pb/238U ages, the equations are:

$$R_{206/238i,k} = FF_{F206/238i,k}\left(\frac{S_{C206i,k}}{S_{C238i,k}}\right) \qquad \textbf{Equation 9a}$$

where:
k = spot number in session
i = fixed signal scan number
$R_{206/238i,k}$ = 206Pb/238U ratio based on calculated 206Pb and 238U signal intensities at signal intensity scan i and spot position k
$FF_{F206/238i,k}$ = fitted 206Pb/238U fractionation value at signal scan i and spot number k (**Equation 8a**)
$S_{C206i,k}$ = calculated 206Pb signal intensity at signal intensity scan i and spot position k (**Equation 5**)
$S_{C238i,k}$ = calculated 238U signal intensity at signal intensity scan i and spot position k (**Equation 5**)

$$t_{206/238i,k} = Age206238(R_{206/238i,k}) \qquad \textbf{Equation 9b}$$

where:
k = spot number in session
i = fixed signal scan number
*Age*206238 = function for age calculation (**Appendix 3**)
$t_{206/238i,k}$ = 206Pb/238U age at signal intensity scan i and spot position k
$R_{206/238i,k}$ = 206Pb/238U ratio based on calculated 206Pb and 238U signal intensities at signal intensity scan i and spot position k (**Equation 9a**)

$$\sigma_{R206/238i,k} = R_{206/238i,k}\left(\left(\frac{\sigma_{SE206}}{S_{C206i,k}}\right)^2 + \left(\frac{\sigma_{SE238}}{S_{C238i,k}}\right)^2 + \left(\frac{\sigma_{t206/238}}{\sum_{j=1}^{N_{PAS}} t_{206/238j}}\right)\right)^{1/2}$$

Equation 9c

where:
k = fixed spot number in session
i = signal scan number
j = primary age standard number
$\sigma_{R206/238i,k}$ = absolute error for $R_{206/238i,k}$

$R_{206/238i,k}$ = 206Pb/238U ratio based on calculated 206Pb and 238U signal intensities at signal intensity scan *i* and spot position *k* (**Equation 9a**)
σ_{SE206} = standard error of ($BS_{Mi} - B_{Fnlast}$) about S_{Ci} for 206Pb (**Equation 6b**)
σ_{SE238} = standard error of ($BS_{Mi} - B_{Fnlast}$) about S_{Ci} for 238U (**Equation 6b**)
$S_{C206i,k}$ = calculated 206Pb signal intensity at signal intensity scan *i* and spot position *k* (**Equation 5**)
$S_{C238i,k}$ = calculated 238U signal intensity at signal intensity scan *i* and spot position *k* (**Equation 5**)
$\sigma_{t206/238}$ = standard deviation of 206Pb/238U ages for the primary age standard
N_{PAS} = number of primary age standard spots
$t_{206/238j}$ = 206Pb/238U age for primary age standard *j* (**Equation 8b**)

$$\sigma_{wmcR206/238k} = \left(\sum_{i=1}^{N_{CScan}} \frac{1}{\left(\sigma_{R206/238i,k}\right)^2} \right)^{-1/2}$$ **Equation 9d**

where:
k = fixed spot number in session
i = concordant scan number *i*
$\sigma_{wmcR206/238k}$ = weighted mean error of $R_{206/238i,k}$ values for all concordant scans at fixed spot number *k*
$\sigma_{R206/238i,k}$ = absolute error for $R_{206/238i,k}$ (**Equation 9c**)
N_{CScan} = number of concordant scans at fixed spot number *k*

$$R_{wmcR206/238k} = \frac{\sum_{i=1}^{N_{CScan}} \frac{R_{206/238i,k}}{\left(\sigma_{R206/238i,k}\right)^2}}{\sum_{i=1}^{N_{CScan}} \frac{1}{\left(\sigma_{R206/238i,k}\right)^2}}$$ **Equation 9e**

where:
k = fixed spot number in session
i = concordant scan number *i*
$R_{wmc206/238k}$ = weighted mean of $R_{206/238i,k}$ values for all concordant scans at fixed spot number *k*
$\sigma_{wmcR206/238k}$ = weighted mean error of $R_{206/238i,k}$ values for all concordant scans at fixed spot number *k* (**Equation 9d**)
$R_{206/238i,k}$ = 206Pb/238U ratio based on calculated 206Pb and 238U signal intensities at signal intensity scan *i* and spot position *k* (**Equation 9a**)
$\sigma_{R206/238i,k}$ = absolute error for $R_{206/238i,k}$ (**Equation 9c**)
N_{CScan} = number of concordant scans at fixed spot number *k*

$$t_{wmc206/238k} = Age206238(R_{wmc206/238k})$$ **Equation 9f**

where:
k = spot number in session
$Age206238$ = function for age calculation (**Appendix 3**)
$t_{wmc206/238k}$ = weighted mean concordant 206Pb/238U age at spot position k
$R_{wmc206/238k}$ = weighted mean of $R_{206/238i,k}$ values for all concordant scans at fixed spot number k

$$\sigma_{twmc206/238k} = Age206238(\sigma_{wmcR206/238k})$$ **Equation 9g**

where:
k = spot number in session
$\sigma_{twmc206/238k}$ = weighted mean concordant 206Pb/238U age error at spot position k
$\sigma_{wmcR206/238k}$ = weighted mean error of $R_{206/238i,k}$ values for all concordant scans at fixed spot number k (**Equation 9d**)

The above equations apply to the 207Pb/235Uc system as well. For the 207Pb/206Pb system, the 207Pb/206Pb ratio at 0 Ma is approximately 0.046003. The weighted mean concordant 207Pb/206Pb age error is calculated as follows:

$$\sigma_{twmc207/206k} = Age207206(R_{wmcR206/238k} + \sigma_{wmcR207/206k}) - Age207206(\sigma_{wmcR207/206k})$$
Equation 9h

where:
k = spot number in session
$Age207306$ = function for age calculation (**Appendix 3**)
$\sigma_{twmc207/206k}$ = weighted mean concordant 207Pb/206Pb age error at spot position k
$R_{wmc207/206k}$ = weighted mean of $R_{207/206i,k}$ values for all concordant scans at fixed spot number k (**Equation 9e** for 207Pb/206Pb)
$\sigma_{wmcR207/206k}$ = weighted mean error of $R_{207/206i,k}$ values for all concordant scans at fixed spot number k (**Equation 9d** for 207Pb/206Pb)

The preferred age for a spot exhibiting at least one concordant scan is either the 206Pb/238U or the 207Pb/206Pb age, whichever exhibits the lowest relative error.

- *Preferred age based on discordia intercepts with the concordia*: A discordia (207Pb/235Uc abscissa, 206Pb/238U ordinate) is characterized for each spot from an unknown by linear regression of the (207Pb/235Uc, 206Pb/238U) points for the spot, giving a regression intercept and regression intercept error, and a regression slope and regression slope error (routine LFIT in Press et al.,1992). The lower intercept age is the lowest age consistent with a point of intersection between the discordia and the concordia. The upper intercept age is the highest of

two ages consistent with two points of intersection between the discordia and the concordia. Sometimes, for example when the laser ablates zircon containing a mixture of two distinct age domains, the number of concordant scans is zero. For these spots, the preferred age is set equal to either the age of the upper or lower intercept between the discordia and the concordia. The error for each of these ages is obtained from the standard deviation of the respective intercept ages derived from the Monte Carlo simulation of the discordia combined with the relative errors of the 206Pb/238U and the 207Pb/206Pb fractionation factors.

$$\sigma_{t\,\mathrm{int}} = t_{\mathrm{int}} \left(\left(\frac{\sigma_{MC\,\mathrm{int}}}{t_{\mathrm{int}}} \right)^2 + \left(\frac{\sigma_{t\,206/238}}{\sum_{j=1}^{N_{PAS}} t_{206/238\,j}} \right)^2 + \left(\frac{\sigma_{t\,207/235c}}{\sum_{j=1}^{N_{PAS}} t_{207/235cj}} \right)^2 \right)^{1/2}$$

Equation 10

where:
j = primary age standard number
σ_{tint} = intercept age error
t_{int} = intercept age
σ_{MCint} = standard deviation of Monte Carlo-derived intercept ages about t_{int}
N_{PAS} = number of primary age spots
$\sigma_{t206/238}$ = standard deviation of 206Pb/238U ages for the primary age standard
$t_{206/238j}$ = 206Pb/238U age for primary age standard j (**Equation 8b**)
$\sigma_{t207/235c}$ = standard deviation of 207Pb/235Uc ages for the primary age standard
$t_{207/235cj}$ = 207Pb/235Uc age for primary age standard j (**Equation 8b**)

The age chosen as the preferred age is the intercept age that exhibits the lowest relative error.

2. Workbook: UPbICPReport_

2.1. General

LA-ICP-MS-based U-Pb zircon analysis of a single geological sample is comprised of a series of spots analyzed on a series of zircon grains (usually one spot per grain). An Excel 97-2003 workbook **UPbICPReport_** is provided containing 7 worksheets that present summary data from UPbICP for this series of grains. It is necessary to ‘enable macros’ for this worksheet to work properly. If a crash occurs during any code execution, it is necessary to either ‘reset’ Visual Basic execution prior to trying again or to close and then reopen the workbook.

- **Worksheet: Data** provides access to and permits the setting of data quality filters; and it presents a single line of calculated ages and related data for each spot analyzed.
- **Worksheet: Calculate** presents options and results related to age population and age spectrum calculations.
- **Worksheet: Plots** presents age spectra plots, age data plots, and several plots involving U and Th concentrations.
- **Worksheet: Standard_Statistics** presents data from the primary and secondary age standards used to calculate ages.
- **Worksheet: A2Z_Standards** presents preferred ages calculated for internal age standards used at A2Z from the LA-ICP-MS session(s) indicated.
- **Worksheet: Comment** is used to verify whether or not data have been loaded into arrays prior to the initiation of various calculations in **Calculate**.
- **Worksheet: Results_Settings** presents a summary of the age data for the preferred grain age population as specified in **Calculate** and the data quality filters as specified in **Data**.

2.2. Worksheet: Data (Figure 4)

2.2.1. Data quality filters. Several data quality filters are employed for objective filtering of preferred ages. The purpose of these quality filters is to parse the data from the series of spots analyzed and distinguish zircon spots from non-zircon spots, or zircon spots that show evidence of problematic data.

- **Min Fit Quality** (cell B4)

The χ^2 value (**Equation 3**) and number of degrees of freedom (**Equation 4**) for the fit of the sum of Gaussian equations to each isotope permits the probability of worse fit to be calculated (routine GAMMQ in Press et al.,1992). If the probability of worse fit *Q* for any one of the isotopes used to calculate ages (206Pb, 207Pb, 208Pb, 232Th, and 238U) is less than this parameter (<0.0500 is commonly used; cell B4 shows the value of this data quality filter; columns Y and AH show the values for each spot), the preferred age is rejected from further analysis. This parameter can be set by the user.

$$QF_{\text{fit}} = \min(Q_{206\text{Pb}}, Q_{207\text{Pb}}, Q_{208\text{Pb}}, Q_{232\text{Th}}, Q_{238\text{U}}) \qquad \textbf{Equation 11}$$

where:
QF_{fit} = minimum *Q* value (columns Y and AH); maximum probability of worse fit value
$Q_{206\text{Pb}}$, etc. = *Q* value from GAMMQ for the sum of Gaussian equations fitted to the 206Pb background+signal intensities, etc.

- **Min Clean Quality** (cell D4)

During the fitting of the sum of Gaussian equations for an isotope, a first pass is performed and then outliers greater than 2 standard deviations from the sum of fitted Gaussian equations are rejected and the fitting process is repeated. In some cases, this filtering process causes a sequence of consecutive and potentially useful scans to be

rejected. If the inverse of the number of consecutive, rejected scans for any one of the isotopes used to calculate ages (206Pb, 207Pb, 208Pb, 232Th, and 238U) is less than this parameter (<0.3333 is commonly used; cell D4 shows the value of this data quality filter; columns Z and AI show the values for each spot), the preferred age is rejected from further analysis. This parameter can be set by the user.

$$QF_{clean} = \min\left(\left(\frac{1}{CO_{206Pb}}\right),\left(\frac{1}{CO_{207Pb}}\right),\left(\frac{1}{CO_{208Pb}}\right),\left(\frac{1}{CO_{232Th}}\right),\left(\frac{1}{CO_{238U}}\right)\right)$$

Equation 12

where:
QF_{clean} = minimum CO^{-1} value (columns Z and AI); maximum number of consecutive background+signal intensities cleaned for each isotope
CO_{206Pb}, etc. = CO value for 206Pb background+signal intensities, etc.

If this parameter is <0.3333 for a spot (a maximum of 3 consecutive rejected outliers is acceptable), the preferred age is rejected from further analysis. This parameter can be user set.

- **Min 204Pb Quality** (cell F4)

The presence of significant 204Pb indicates the presence of significant common Pb in the target zircon. For each scan, the ratio of the background corrected 204Pb signal intensity to the background 204Pb intensity is determined and the maximum value of this ratio for scans for the spot is determined. If this maximum ratio is less than the set value of this data quality parameter (<0.3333 is commonly used; cell F4 shows the value of this data quality filter; columns AA and AJ show the values for each spot), the preferred age is rejected from further analysis. This parameter can be set by the user.

$$QF_{204Pb} = \max\left(\frac{S_{C204Pbi}}{B_{F204PbNlast}}\right) \text{ for } i\text{=1 to } N_{\text{scan}}$$

Equation 13

where:
QF_{204Pb} = minimum signal intensity divided by background correction intensity (columns AA and AJ)
$S_{C204Pbi}$ = 204(Hg+Pb) signal intensity at signal intensity scan i (**Equation 5**)
$Nlast$ = scan number of the last background intensity scan
$B_{F204PbNlast}$ = fitted 204(Hg+Pb) background intensity at background intensity scan $Nlast$ (**Equation 1** where $i = Nlast$)

- **Min Uppm Quality** (cell H4)

This quality parameter was originally designed to eliminate spots in grains that were likely not zircon, using an old LA-ICP-MS data collection protocol that did not permit use of the **Mineral (or "n.a.")** filter (see next paragraph). Spots that exhibit

Uppm values (column I) < 9 are eliminated from further analysis. This parameter can be user set.

- **Mineral (or "n.a.")** (cell J4)

The current LA-ICP-MS data collection protocol used by A2Z includes the analysis of 28Si and 91Zr near the bottom of a laser ablation pit. A spot that exhibits a 91Zr/28Si ratio within 3σ of the median value obtained for all zircon spot analyses of the primary age standard during a LA-ICP-MS session are deemed to be zircon. These analyses are indicated by the extension '_Zrn' at the end of the Analysis_# in column A. Non-zircon spots, or spots for which the laser has either blasted the grain out of the mounting medium or passed through the grain into the mounting medium (the latter causing a faster decay of the 91Zr background+signal intensity relative to the 28Si background+signal intensity), do not have this extension added. This parameter can be user set to "n.a." to eliminate this filter entirely.

2.2.2. Column contents. The contents of each column are described below.

Measured Isotopic Ratios

- **Analysis_#** (column A)

Extension '_Zrn' indicates a spot that exhibits a 91Zr/28Si ratio within 3σ of the median value obtained for all zircon spot analyses of the primary age standard during a LA-ICP-MS session.

- **207Pb/235Uc*** (column B)
- **207Pb/235Uc* (±2σ)** (column C)
- **206Pb/238U** (column D)
- **206Pb/238U (±2σ)** (column E)
- **207Pb/206Pb** (column F)
- **207Pb/206Pb (±2σ)** (column G)

The ratios are weighted means of the signal ratios for all concordant scans. A scan is concordant if its 207Pb/235U_c, 206Pb/238U, and 207Pb/206Pb ages overlap each other at the 2σ level. The ratio errors represent the errors of the respective weighted mean ratios combined with the relative errors for the respective fractionation factors. A value of 'undefined' appears when there are no concordant scans. A value of 'poor data' or '-' appears when the data do not permit the calculation of realistic U-Pb ages.

U and Th

- **U (ppm)** (column I)
- **Th (ppm)** (column J)
- **U/Th** (column K)

The average U and Th concentrations for primary age standard FC-1 are assumed to be equal to the weighted mean values shown in **Table 3** (data from Paces and Miller, 1993). A median value of average 235Uc signal intensity per scan and a median value of average 232Th signal intensity per scan are determined for all FC-1 primary age standard spots. The assumed average U and Th concentrations for the primary age standard FC-1 are divided by these ratios giving Uppm*scan/235Uc signal intensity

and Thppm*scan/232Th signal intensity factors. These factors are then used to convert 235Uc signal intensity/scan and 232Th signal intensity/scan to Uppm and Thppm for all scans (age standards and unknowns). For unknown and secondary age standard spots exhibiting at least 1 concordant scan, the average Uppm and Thppm concentrations for the concordant scans is reported here. For spots exhibiting no concordant scans, the average Uppm and Thppm concentrations for the discordant scans are reported here.

Apparent Ages

- **Analysis_#** (column M)

Unlike column A, no extension is added to analysis number to indicate zircon.

- **207Pb/235Uc*** (column N)
- **207Pb/235Uc* (±2σ)** (column O)
- **206Pb/238U** (column P)
- **206Pb/238U (±2σ)** (column Q)
- **207Pb/206Pb** (column R)
- **207Pb/206Pb (±2σ)** (column S)

These ages and errors are based on the **Measured Isotopic Ratios** (columns B-G) described above.

Preferred Ages

- **Analysis_#** (column U)

Unlike column A, no extension is added to analysis number to indicate zircon.

- **Preferred Age (Ma)** (column V)

Preferred age based on the weighted mean isotopic ratios of concordant scans: This strategy seeks the weighted mean of all isotopic ratios from concordant scans. The preferred age for a spot exhibiting at least one concordant scan is either the 206Pb/238U or the 207Pb/206Pb age (based on weighted mean isotopic ratios from concordant scans), whichever exhibits the lowest relative error. *Preferred age based on discordia intercepts with the concordia*: For spots exhibiting no concordant scans, the preferred age is set equal to either the age of the upper or lower intercept between the discordia and the concordia. The error for each of these ages is obtained from the standard deviation of the respective intercept ages derived from the Monte Carlo simulation of the discordia combined with the relative errors of the 206Pb/238U and the 207Pb/206Pb fractionation factors. The age chosen as the preferred age is the intercept age that exhibits the lowest relative error. A value of ‘undefined’ appears when a discordia cannot be fitted (e.g., all points lying above the concordia). A value of ‘undefined’ appears specifically for the upper intercept age and error if there is no upper intercept between the dicordia and concordia.

- **Absolute Error (±2σ)** (column W)

The error, including fractionation factor error, of the preferred age described above.

- **Age Type** (column X)

Types include a) ‘Pb/U wmean conc.’= 206Pb/238U weighted mean concordant, b) ‘Pb/Pb wmean conc.’= 207Pb/206Pb weighted mean concordant, c) ‘lower int.’=age

based on the lower intercept between the discordia and the concordia, and d) ‘upper int.’= age based on the upper intercept between the discordia and the concordia. A value of ‘poor data’ or ‘-‘ appears when the data do not permit the calculation of realistic U-Pb ages.

- **Fit Quality 1=best** (column Y)

This value indicates the minimum probability of worse fit for all of the isotopes used for calculate ages (206Pb, 207Pb, 208Pb, 232Th, and 238U).

- **Clean Quality 1=best** (column Z)

This value indicates the maximum value of [1 / (maximum number of consecutive rejected outliers)] for all of the isotopes used to calculate ages (206Pb, 207Pb, 208Pb, 232Th, and 238U).

- **204Pb Quality 1=best** (column AA)

This value indicates the minimum value of [(204(Hg+Pb)_background / (204(Hg+Pb)_signal + 204(Hg+Pb)_background)] for all data scans from a spot.

- **2σ Conc. Scans** (column AB)

This value indicates the number of concordant scans for the spot.

Preferred Ages - Sorted

- **Analysis_#** (column AD)
- **Preferred Age (Ma)** (column AE)
- **Absolute Error (±2σ)** (column AF)
- **Age Type** (column AG)
- **Fit Quality 1=best** (column AH)
- **Clean Quality 1=best** (column AI)
- **204Pb Quality 1=best** (column AJ)
- **2σ Conc. Scans** (column AK)

These are the same data as provided in columns U-AB; however, the data here have been sorted in ascending order of the value in column AE.

Fitted Discordia

- **Age-Lower Intercept (Ma)** (column AM)

A discordia (207Pb/235Uc abscissa, 206Pb/238U ordinate) is characterized for each spot from an unknown by linear regression of the (207Pb/235Uc, 206Pb/238U) points for the spot, giving a regression intercept and regression intercept error, and a regression slope and regression slope error. The regression is performed using routine LFIT in Press et al. (1992). The lower intercept age is the lowest age consistent with a point of intersection between the discordia and the concordia. If the regression intercept is <0 then a value of ‘undefined’ appears here. If the lower intercept age is <0 Ma or >4560 Ma then a value of ‘undefined’ appears here.

- **Age-Lower Intercept (±2σ)** (column AN)

See comments for column AM above. The errors for the upper and lower intercepts of the discordia with the concordia are calculated by Monte Carlo simulation of the discordia using 1000 randomly generated lines based on the regression-derived slope and intercept values and their respective errors. If a value other than ‘undefined’

appears in column AM, the standard deviation of the Monte Carlo-derived lower intercepts is calculated about the lower intercept age, ignoring any intercepts <0 Ma or >4560 Ma. The ±2σ error on the lower intercept age is calculated based on the relative errors of the lower intercept age and the 206Pb/238U and 207Pb/206Pb fractionation factors (σ = lower_intercept_age * sqrt(relative_error_lower_intercept_age^2 + relative_error_206Pb/238U_fractionation_factor^2 + relative_error_207Pb/206Pb_fractionation_factor^2)).

- **Age-Upper Intercept (Ma)** (column AO)

See comments for column AM above. The upper intercept age is the highest of two ages consistent with points of intersection between the discordia and the concordia. If the discordia fails to intersect the concordia twice, the upper intercept age does not exist and a value of 'undefined' appears here. If the upper intercept age is >4560 Ma and zero upper intercepts <4560 Ma are encountered during the Monte Carlo simulation of the discordia, the upper intercept age does not exist and a value of 'undefined' appears here. If the upper intercept age is >4560 Ma and at least one upper intercept <4560 Ma is encountered during the Monte Carlo simulation, the upper intercept age does exist and a value of '4560' appears here.

- **Age-Upper Intercept (±2σ)** (column AP)

See comments for column AM above. The errors for the upper and lower intercepts of the discordia with the concordia are calculated by Monte Carlo simulation of the discordia using 1000 randomly generated lines based on the regression-derived slope and intercept values and their respective errors. If a value other than 'undefined' appears in column AO, the standard deviation of the Monte Carlo-derived upper intercepts is calculated about the upper intercept age, ignoring any intercepts <0 Ma or >4560 Ma. The ±2σ error on the upper intercept age is calculated based on the relative errors of the upper intercept age and the 206Pb/238U and 207Pb/206Pb fractionation factors (σ = upper_intercept_age * sqrt(relative_error_upper_intercept_age^2 + relative_error_206Pb/238U_fractionation_factor^2 + relative_error_207Pb/206Pb_fractionation_factor^2)).

- **Regression Intercept** (column AQ)

This value is the y-axis intercept of the best-fit line through all (207Pb/235Uc abscissa, 206Pb/238U ordinate) points for all scans, calculated using routine LFIT in Press et al. (1992).

- **Regression Intercept (±1σ)** (column AR)

This value is calculated using LFIT in Press et al. (1992).

- **Regression Slope** (column AS)

This value is the slope of the best-fit line through all (207Pb/235Uc abscissa, 206Pb/238U ordinate) points for all scans, calculated using routine LFIT in Press et al. (1992).

- **Regression Slope (±1σ)** (column AT)

This value is calculated using LFIT in Press et al. (1992).

Analysis Details

- **% Minimum Discordance** (column AV)

Discordance here is defined as 100*(207Pb/206Pb_age -206Pb/238Pb_age) / 207Pb/206Pb_age (%). If either 207Pb/206Pb_age or 206Pb/238Pb_age is zero, this discordance value is set to -200.00%. The minimum value of this discordance for all scans is reported here. Any value <-999.99% is arbitrarily set to -999.99%. If the value reported here is -200.00%, then at least one scan yielded either a 207Pb/206Pb_age or a 206Pb/238Pb_age value equal to zero and all other discordance values for the scans were >-200.00%.

- **% Maximum Discordance** (column AW)

The maximum value of this discordance for all scans is reported here. If the value reported here is -200.00%, then at least one scan yielded either a 207Pb/206Pb_age or a 206Pb/238Pb_age value equal to zero and all other discordance values for the scans were <-200.00%.

- **Preferred Scan #** (column AX)

The scan number that yielded the discordance value closest to zero.

- **Discordia Scans** (column AY)

The number of scans used to calculate the discordia.

- **2σ Conc. Scans** (column AZ)

This value indicates the number of concordant scans for the spot. This is the same as columns AB and AK (column AK is sorted with other data).

- **Relative α-Radiation** (column BA)

For each spot, this number is relative to the median α-radiation value for all primary age standard spots. It represents the sum of the U and Th signal values for a spot, integrated over the preferred age of the spot, using the decay constants for various isotopes. Information regarding its calculation and usage is provided in **Appendix 2**.

- **Analysis_#** (column BB)

Unlike column A, no extension is added to analysis number to indicate zircon.

2.3. Worksheet: Calculate (Figure 5)

Calculate requires the following operations be performed so that the various features work properly:

- **Step 1**: Enable macros.
- **Step 2**: Set 'xmax_plot' (cell H5), 'xbinwidth_plot' (cell H6), and 'xstep_plot' (cell H7). Copy H5:H7 and paste to I5:I7. These parameters have the following meaning: xmax_plot = maximum plot x-axis value in Ma (see **Plots**); xbinwidth_plot = histogram bin width for distribution of grain ages into bins; xstep_plot = Δx value for probability distribution function points (xmax_plot/xstep_plot must be ≤5000).
- **Step 3**: Press 'transfer data' button. This operation invokes several Visual Basic routines that read data into arrays and performs the following preliminary calculations:
    - the grain ages in column A are distributed into histogram bins (histogram x and y values are in columns N and O, respectively),

- the grain ages and their 2σ errors in columns A and B, respectively, are converted to individual Gaussian distributions and added together to form the raw data probability distribution function (probability distribution function x and y values are in columns P and Q, respectively),
- peaks and troughs are determined for the raw data probability distribution function and following parameters related to the Gaussian deconvolution are set:
    - **ngauss**: Cell H14 is the number of Gaussians into which **Calculate** prefers to deconvolve the probability distribution function. The maximum number is 20. The minimum number is 3. For a counted number of probability distribution function peaks between 1 and 16, the peaks are ranked in order of decreasing area and the largest two are split into 3 Gaussians of area equal to 1/3 of their parent Gaussian (by dividing the amplitude by 3) and positions x=mean-1σ, x=mean, x=mean+1σ. For a counted number of probability distribution function peaks >16, only the largest 16 are used. Means are specified as the peak x-position (in Ma); standard deviations are specified as equal to area/(amplitude*1.772415); areas are then normalized to the area of the raw data probability distribution function.
    - **gauss**: Cells G22:G41
    - **amplitude**: Cells H22:H41
    - **mean**: Cells I22:I41
    - **std dev**: Cells J22:J41
    - **data fraction**: Cells K22:K41

  These parameters represent the initial guesses for the Gaussians after pressing 'transfer data'.
- d) all of the grains are combined into a single, preferred population (data in *red italics*) and the following data are calculated for the preferred population:
    - **# ages**: Cell H2 contains the number of ages total; cell I2 contains the number of ages in the red preferred population.
    - **maximum row**: Cell J11 contains the maximum row for the age to be included in the preferred population. The ages in column A are ranked from youngest to oldest (and below the grains with acceptable ages are the rejected grains with the following codes: F C Pb U M; _ _ Pb _ _ indicates rejection for failure to pass the quality test for Pb, and so on). Upon pressing 'transfer data', maximum row is set to the row of the oldest, acceptable grain and all grains are included in the population.
    - **wmean**: Cell L6 contains the weighted mean age of the grains in preferred population.
    - **2σ+FFerr**: Cell L8 contains the 2σ error of the weighted mean age of the grains in the preferred population; this error also includes the fractionation factor error.

- **2σ**: Cell L10 contains the 2σ error of the weighted mean age of the grains in the preferred population; this error does not include the fractionation factor error.
- **MSWDw**: Cell L12 contains the weighted mean square of the weighted deviates of the grain ages in the preferred population.
- **mean**: Cell M6 contains the mean age of the grains in preferred population.
- **2σ stddev**: Cell M8 contains the 2 standard deviation error of the mean age of the grains in the preferred population.
- **2σ stderr**: Cell M10 contains the 2 standard error of the mean age of the grains in the preferred population.
- **P(MSWDw)**: Cell M12 contains MSWDw probability (probability of a greater MSWDw value) of the grain ages in the preferred population.
- **start_row**: Cell L14 contains the row with lowest age in the preferred population.
- **end_row**: Cell M14 contains the row with highest age in the preferred population.
- **# 1σ Absolute Errors**: Cell L16 contains the number of 1σ absolute errors used for overlap to determine concordance between different grains ages within the preferred population. The grain age errors in column B are 2σ. This parameter is very important when using button **maximum row** (see below).
- **Uppm**: Cell M22 contains the mean Uppm value (**Data** column I) for the grains in the preferred population.
- **Thppm**: Cell M23 contains the mean Thppm value (**Data** column J) for the grains in the preferred population.
- **U/Th**: Cell M24 contains the mean U/Th value (**Data** column K) for the grains in the preferred population.
- **alpha**: Cell M25 contains the mean relative α-radiation damage value (**Data** column BA) for the grains in the preferred population.

- **Step 4**: Press 'rescale plots' button. This operation rescales the plots in **Plots** to the specified xmax_plot value.

  Calculate is now ready to work. The grain age columns are described below:

- **Preferred Age (Ma)** (column A)

From **Data** column AE.

- **Absolute Error (±2σ)** (column B)

From **Data** column AF.

- **Age Type** (column C)

From **Data** column AG.

- **Analysis #** (column D)

From **Data** column AD.

- **Product of Qualities** (column E)

Product of the values from columns **Data** AH, AI, AJ.

Suppose one desires the age of the largest population for which all grains are concordant, overlapping by the # of 1σ absolute errors set in cell L16. Simply ensure that the maximum row parameter in cell J11 indicates the row number of the oldest grain and press the button 'maximum row'. All of the preferred population parameters (in *red italics*) will update.

Suppose one desires the age of the youngest population containing 2 or more grains. Set the maximum row in cell J11 to 5 and press button 'maximum row'. If the grain in row 5 is NOT part of the population in *red italics* but grains in rows 2-4 are, then this is the youngest population. If this fails, try another value of maximum row in cell J11 and press button 'maximum row'. Repeat until the data of interest appear for the preferred population.

Suppose one desires the age of the population for grains in rows 11-17, regardless of their ages (whether or not they are concordant) and regardless of the ages for grains in rows 2-10 and 18+. Change parameter start_row in cell L14 to 11 and parameter end_row in cell M14 to 17 and press button 'statistics'. The population in *red italics* will update to reflect on these grains in this row interval.

Finally, press button 'deconvolve' to deconvolve the probability distribution function into the number of Gaussians indicated by parameter ngauss in cell H14. Cells H11, H12, I12, H16 (current mean χ^2 per nodal point deconvolved), H17 (best mean χ^2 per nodal point deconvolved), and G22:K41 will update as the deconvolution progresses and finally finishes. Parameters dym_factor in cell H15 and chisqtol in cell H18 may be user set but this is not recommended. The significance of these parameters pertains to the chi-squared minimization process employed during the deconvolution and the may be ascertained directly by viewing the source code and searching for 'dymfactor' and 'chisqtol'.

The curious user can investigate the meaning of the remaining columns and cells in this worksheet.

2.4. Worksheet: Plots (Figures 6a and 6b)

The plots are self-explanatory:

- **U-Pb Age (Ma) vs. Relative Age Frequency**

U-Pb age on x-axis vs. raw data probability distribution function and histogram bins on y-axis.

- **U-Pb Age (Ma) vs. Relative Age Frequency/U/Th**

U-Pb age on x-axis vs. raw data probability distribution function and U/Th on y-axis.

- **U-Pb Age (Ma) vs. Relative Age Frequency**

U-Pb age on x-axis vs. raw data probability distribution function and Gaussian deconvolution probability distribution function on y-axis.

- **U-Pb Age (Ma) vs. U/Th**

U-Pb age on x-axis vs. U/Th on y-axis.

- **U-Pb Age (Ma) vs. Frequency**

Gaussin deconvolution peak mean x-axis vs. Gaussian deconvolution peak data fraction on y-axis.

- **U-Pb Age (Ma) vs. U ppm**

U-Pb age on x-axis vs. U on y-axis.

- **Upper-Age Intercept (Ma) vs. Lower-Age Intercept (Ma)**

U-Pb upper-age intercept on x-axis vs. U-Pb lower-age intercept on y-axis.

- **U-Pb Age (Ma) vs. Th ppm**

U-Pb age on x-axis vs. Th on y-axis.

2.5. Worksheet: Standard_Statistics (Figure 7)

This sheet contains information pertaining to the primary age standard FC-1 (indicated by age standard index 6 in UPbICP code; cell B2 here; age standard name F in UPbICP code; cell D2 here) of accepted age 1099 Ma (cell B3) and the secondary age standard TR run in pair along with the primary age standard (indicated by age standard index 2 in UPbICP code; cell B1 here; age standard name T in UPbICP code; cell D11 here) of accepted age 61.23 Ma (cell B12). Cells B5:B9 display mean preferred U-Pb ages for the primary age standard FC-1, calculated when FC-1 is treated as an unknown. Cells C5:C9 present 1σ relative errors on the ages in cells B5:B9; these relative errors are used as the relative errors of the various isotopic ratio fractionation factors for the session.

During a first-pass run of UPbICP, the weighted mean age for the secondary age standard is given as the uncorrected age in cell B13 (uncorrected meaning that the fractionation factors were not corrected for relative α-radiation damage). Parameter measAgeStdPreferred2 in **Appendix 2** subroutine AlphaEnergyFF is the accepted age of this secondary age standard in cell B12. Parameter std207235Age(nAgeStdPreferred2) (and for the other isotopic ratios) in **Appendix 2** subroutine AlphaEnergyFF is the uncorrected age of this secondary age standard in cell B13.

Cell B17 gives the median 91Zr/28Si (neither background corrected) value for all primary age standard spots and cell B18 gives the 1σ standard deviation about this median ratio. All spots during the session within 3σ of this median value are deemed zircon and the extension '_Zrn' (in cell B15) is added to the Analysis_# name in **Data** column A. The values in cells C17:D20 represent statistics calculated for non-background corrected counts for the primary age standard.

2.6. Worksheet: A2Z_Standards (Figure 8)

These columns are self-explanatory and provided for information purposes only. This is a compilation of U-Pb zircon ages (column C) and their 2σ errors (column D) for all secondary age standards in the session from which the data in worksheet **Data** were derived. A value near zero for cell I2 and a value near one for cell K2 is desired.

2.7. Worksheet: Comment (Figure 9)

This sheet is required to provide a necessary comment in the event calculations are attempted in **Calculate** prior to pressing the 'transfer data' button.

2.8. Worksheet: Results_Settings (Figure 10)

These data are described above and they are simply combined here, minus all of the other data, for easy viewing. Click on each value to see the source of the value in the status bar.

3. Workbook: UPbICPViewGrain_

3.1. General

This workbook is provided as a tool to permit the user to quickly view the raw data, sum of Gaussian equation fits, and discordia for individual spots analyzed. The workbook is composed of 5 worksheets:

- **Worksheet: Raw_Data** allows the raw data from the ICP-MS system to be viewed, along with data pertaining to the background values and fitted Gaussians for each isotope.
- **Worksheet: Raw_Data_Plots** shows plots of the raw data.
- **Worksheet: bkg_Corrected_Data_Plots** shows plots of the background correted data and several relevant background corrected isotopic ratio plots.
- **Worksheet: Discordia** shows age and discordance versus scan number plots as well as discordia plots on the concordia diagram.
- **Worksheet: Standard_Statistics** presents data from the primary and secondary age standards used to calculate ages.

3.2. Worksheet: Raw_Data (Figure 11)

Note that it is important to exit active cells containing data being set for the program to operate properly. Prior to pressing any of the buttons described below, an empty cell should be selected as the active cell. The following steps must be taken in order for this program to work:

1. Select the **Raw_Data** tab at the bottom of the workbook.
2. File location examples from A2Z session z031110:
   - Location of run details file (*.csv):
     - Folder: c:\UPbICP\UPbZrSiZ031110\
     - File: UPbZrSiZ031110.csv
   - Location of ICP-MS data files (*.txt):
     - Folder: c:\UPbICP\UPbZrSiZ031110\
     - Files: *.txt where * indicates sample root name and grain number (sample root name, cell U14, is case sensitive; cases can be verified in the run details file UPbZrSiZ031110.csv or by checking the names of the *.txt files themselves)
   - Location of UPbICP output files (*.txt):
     - Folder: c:\UPbICP\ UPbZrSiZ031110\
     - Files:

- UPbZrSiZ031110StdBestGrainAge.txt
- UPbZrSiZ031110StdBkgFit.txt
- UPbZrSiZ031110StdScanAge.txt
- UPbZrSiZ031110StdSigDelete.txt
- UPbZrSiZ031110StdSigFit.txt
- UPbZrSiZ031110StdFF.txt (not used at the time of this writing)
- UPbZrSiZ031110UnkBestGrainAge.txt
- UPbZrSiZ031110UnkBkgFit.txt
- UPbZrSiZ031110UnkScanAge.txt
- UPbZrSiZ031110UnkSigDelete.txt
- UPbZrSiZ031110UnkSigFit.txt

3. Set run details file (cell U1) and press the upper 'refresh run details file' button. Cells U2:U9 are updated by this operation. Data transfer is complete when cell U15 is made the active cell.
4. Set U11 through U17 and U20 (the spin up/down button may not work well, if this happens, set U15 manually). **Cell U14 is case sensitive**. Only 'Std' or 'std' are valid for age standard; only 'Unk' or 'unk' are valid for unknown in cell U20. Cell U17 is the maximum grain number for which an ICP-MS data file is available for the root filename in cell U14 in the specified folder location. Press 'refresh grain' button. Five files are read during data transfer (progress reported in cells U24 and U25). Data transfer and the calculations are complete when cell Q20 on worksheet **Discordia** is made the active cell.

   Brief descriptions of various cells and columns follow:

- Columns A-R are read directly from, or calculated based on, data contained in the *.txt file for the spot of interest (17103b_20.txt in cell U18 in **Figure 11**).
- Cells AA2:AI10 are updated by reading the data in *BkgFit.txt pertaining to the spot of interest.
- Cells AA15:AI112 are updated by reading the data in *SigFit.txt pertaining to the spot of interest.
- Cells W22:W31 represent the area/10000 of the deleted background-corrected, measured signal intensities. Cells X22:X31 represent the area/10000 of the background-corrected, measured signal intensities to which the sum of Gaussian equations is fitted. Cells Y22:Y31 represent the area/10000 of the sum of Gaussian equations fitted to the background corrected, measured signal intensities.
- Column pairs AM and AN, AO and AP…BC and BD contain background corrected signal data (value in AM91 equals bkg+signal value in C91 minus the bkg value in AH2; see the formula in cell AM91), and calculated signal data fitted values (value in AN91 is the value indicated by the sum of the Gaussians detailed in cells AG15:AI24 minus the bkg value in AH2), respectively.
- Columns BF and BG represent the calculated 207Pb/235Uc and 206Pb/238U ratios, respectively for the signal scan indicated by the discordia index value in column BH, based on the sum of Gaussian equations for the respective isotopes (discordia index 1 equals scan 90, the first bkg corrected signal scan).

- The number 500 in column BH indicates a scan that is not acceptable for calculating the discordia. These data come from *ScanAge.txt.
- Columns BN, BO, and CG represent bkg corrected signal data for the 207Pb/235Uc, 206Pb/238U, and 207Pb/206Pb ratios for the scans shown in column BM and CF.
- Columns BP, BQ, and CH represent the fractionation factors for the 207Pb/235Uc, 206Pb/238U, and 207Pb/206Pb ages, respectively for the scans shown in columns BM and CF. These data come from *ScanAge.txt. Note the 207Pb/206Pb fractionation factors are all set to 1. This is a fundamental feature of UPbICP.
- The number 500 in column BI indicates a scan that is not either not acceptable or not concordant. These data come from *ScanAge.txt.
- Column BS, BT, and BU represent the ages indicated for the scan shown in column BM. These data come from *ScanAge.txt.
- Column BV represents the discordance defined as defined as 100*(207Pb/206Pb_age -206Pb/238Pb_age) / 207Pb/206Pb_age (%). These data come from *ScanAge.txt.
- Cells BX2:BY3 show the minima maxima (the latter set to 1.5 x the maxima of columns BN and BO) of the measured data for plotting the concordia diagram on worksheet **Discordia**.

3.3. Worksheet: Raw_Data_Plots (Figure 12)

1. Select the **Raw_Data_Plots** tab at the bottom of the workbook.
2. The plots are self-explantory.

3.4. Worksheet: bkg_Corrected_Data_Plots (Figure 13a and Figure 13b)

1. Select the **bkg_Corrected_Data_Plots** tab at the bottom of the workbook.
2. The plots are self-explantory.

3.5. Worksheet: Discordia (Figure 14)

1. Select the **Discordia** tab at the bottom of the workbook.
2. The following plots are self-explantory.

- **207Pb/235Uc vs. 206Pb/238U discordia (small scale)**
207Pb/235Uc on x-axis vs. 206Pb/238U on y-axis.
- **207Pb/235Uc vs. 206Pb/238U discordia (large scale)**
207Pb/235Uc on x-axis vs. 206Pb/238U on y-axis.
- **Scan vs. U-Pb Age (Ma)**
Scan number on x-axis vs. 207Pb/235Uc, 206Pb/238U, and 207Pb/206Pb U-Pb ages on y-axis.
- **Scan vs. Discordance (%)**
Scan number on x-axis vs. 206Pb/238U vs. 207Pb/206Pb discordance on y-axis.
- **Scan vs. U/Th**

Scan number on x-axis vs. U/Th on y-axis.

- **Scan vs. U ppm**

Scan number on x-axis vs. U ppm on y-axis.

- **Scan vs. Th ppm**

Scan number on x-axis vs. Th ppm on y-axis.

The ‘refresh’ button does very little at this point. Ultimately, it will calculation of the weighted mean concordant age for a sub-population of user-set scans (user will set Q20 and Q21 and the ‘refresh’ button will perform calculations to update all fields in *red italics*.

3.6. Worksheet: Standard_Statistics (Figure 15)

See the discussion in Section 2.5 above.

References Cited

Black, L.P., Kamo, S.L., Allen, C.M., Davis, D.W., Aleinikoff, J.N., Valley, J.W., Mundil, R., Campbell, I.H., Korsch, R.J., Williams, I.S., and Foudoulis, C. 2004. Improved 206Pb/238U microprobe geochronology by the monitoring of trace-element-related matrix effect; SHRIMP, ID-TIMS, ELA-ICP-MS and oxygen isotope documentation for a series of zircon standards. Chemical Geology, 205, p.15-140

Chang, Z., Vervoort, J.D., McClelland, W.C., and Knaack, C., 2006. U-Pb dating of zircon by LA-ICP-MS. Geochemistry, Geophysics, Geosystems, American Geophysical Union, v. 7, n. 5, 14 p.

Chen, J.H. and Wasserburg, G.J., 1983. The least radiogenic Pb in iron meteorites. 14th Lunar and Planetary Science Conference, Abstracts, Part 1. Lunar and Planetary Institute, Houston, Texas, p. 103-104.

Excel 2007. © Microsoft Corporation.

Friedlander, G., Kennedy, J.W., Macias, E.S., and Miller, J.M., 1981. Nuclear and Radiochemistry. John Wiley and Sons, New York, 685 p.

Gehrels, G.E., Valencia, V.A., and Ruiz, J., 2008. Enhanced precision, accuracy, efficiency, and spatial resolution of U-Pb ages by laser ablation-multicollector-inductively coupled plasma-mass spectrometry. Geochemistry Geophysics Geosystems, American Geophysical Union, v. 9, 13 p.

Kuiper, K.F., Deino, A., Hilgen, P.J., Krijgsman, W., Renne, P.R., and Wijbrans, J.R., 2008. Synchronizing rock clocks of Earth history. Science, 320, p. 500-504.

Lanphere, M.A. and Baadsraard, H., 2001. Precise K-Ar, 40Ar/39Ar, Rb-Sr and U-Pb mineral ages from the 27.5 Ma Fish Canyon Tuff reference standard. Chemical Geology, v. 175, p. 653-671.

Paces, J.B. and Miller, J.D., 1993. Precise U-Pb ages of Duluth Complex and related mafic intrusions, northeastern Minnesota: Geochronological insights to physical, petrogenic, paleomagnetic, and tectonomagmatic processes associated with the 1.1 Ga Midcontinent Rift System. Journal of Geophysical Research, v. 98, no. B8, p. 13997-14013.

Renne, P.R., Swisher, C.C., Deino, A.L., Karner, D.B., Owens, T.L., and DePaolo, D.J., 1998. Intercalibration of standards, absolute ages and uncertainties in 40Ar/39Ar dating. Chemical Geology, v. 45, p. 117-152.

Press, W.H., Teukolsky, S.A., Vetterling, W.T., and Flannery, B.P., 1992. Numerical Recipes in FORTRAN. The Art of Scientific Computing. Second Edition, Cambridge University Press, 963 p.

Steiger, R.H. and Jäger, E., 1977. Subcommission on geochronology: Convention on the use of decay constants in geo- and cosmochronology. Earth and Planetary Science Letters, v. 36, p. 369-371.

Tucker, R.D., Bradley, D.C., Ver Straiten, C.A., Harris, A.G., Ebert, J.R., and McCutcheon, S.R., 1998. New U-Pb zircon ages and the duration and division of Devonian time. Earth and Planetary Science Letters, v. 158, p.175-186.

Appendix 1: Common Pb Correction

202Hg and the combination of 204Hg+204Pb are measured for purposes of common Pb correction. In the analysis below, no fractionation is assumed for the various isotopes; as such, the current A2Z preference is to avoid applying this correction until the fractionation of Hg and Pb isotopes in the apparatus used is better understood. The isotopic abundances of the Hg isotopes measured are (Friedlander et al., 1981):

202Hgfraction = 0.298
204Hgfraction = 0.069

Common Pb isotopic ratios are (Chen and Wasserburg, 1983):

206Pb/204Pbcommon = 9.3066
207Pb/204Pbcommon = 10.293
208Pb/204Pbcommon = 29.475

Common 204Pb is calculated as follows:

204Pbcommon = (204Hg204Pb) – [(204Hgfraction/202Hgfraction) * 202Hg] (if <0 for a scan then set to 0)

Common Pb corrected isotopic ratios are calculated as follows:

206Pb = 206Pb_background_corrected – (204Pbcommon * 206Pb/204Pbcommon) (if <0 for a scan then set to 0)
207Pb = 207Pb_background_corrected – (204Pbcommon * 207Pb/204Pbcommon) (if <0 for a scan then set to 0)
208Pb = 208Pb_background_corrected – (204Pbcommon * 208Pb/204Pbcommon) (if <0 for a scan then set to 0)

Appendix 2: Primary Age Standard Fractionation Factor Correction for α-Radiation Damage

Function dAlphaEnergy (below) is used to calculate the relative α-radiation damage. Th232, U235c, and U238 represent the signal intensities for these isotopes for the current scan. Parameters ageTh232, ageU235c, and ageU238 represent the respective UPb ages for the current scan. Values of dAlphaEnergy are summed for all scans for a spot.

```
Function dAlphaEnergy(U235c, U238, Th232, ageU235c, ageU238, ageTh232, ffage)
    'dim ageU235c, ageU238, ageTh232, ffage(5) implicit
    'U235c
    dAlphaEnergy = (7 * 5.94 * U235c / ffage(2)) * (Exp(lamdaU235 * ageU235c) - 1)
    'U238
    dAlphaEnergy = dAlphaEnergy + (8 * 5.36 * U238 / ffage(3)) * (Exp(lamdaU238 * ageU238) - 1)
    'Th232
    dAlphaEnergy = dAlphaEnergy + (6 * 5.83 * Th232 / ffage(5)) * (Exp(lamdaTh232 * ageTh232) - 1)
End Function
```

Corrections of the fractionation factors for the primary age standard are performed using the following subroutine. Several important parameters have the following meanings:

nAgeStdPreferred1=index for the primary age standard (FC-1 in most cases)
nAgeStdPreferred2=index for the secondary age standard (TR in most cases)
StdAlphaEnergy(nAgeStdPreferred1)=relative α-radiation damage for the primary age standard
StdAlphaEnergy(nAgeStdPreferred2)=relative α-radiation damage for the secondary age standard
SAlphaEnergy=average StdAlphaEnergy(nAgeStdPreferred1) per scan multiplied by the scan number i
UAlphaEnergy=sum of dAlphaEnergy values (function dAlphaEnergy) for scans 1 through i
std207235Age(nAgeStdPreferred2)=accepted 207Pb/235Uc age for the secondary age standard (same for std208238… and std207206…)
measAgeStdPreferred2=measured age for the secondary age standard when calculated *without* fractionation factor correction

Several conditional statements limit the amount of fractionation factor corrections that are permitted. a) If UAlphaEnergy≥SAlphaEnergy then no correction is performed. b) No fractionation factor is permitted to be set less than the minimum value observed for the respective isotopic ratio for the primary age standard.

```
Sub AlphaEnergyFF(SAlphaEnergy, UAlphaEnergy, ffage, minff)
    'dim ffage(5),minff(5) implicit
    Dim x1, x2, y1, y21, y22, y2235, y2238, y2232, m, b
    If UAlphaEnergy < SAlphaEnergy Then
        'calibration for acceptable 207/206
        x1 = 1 'FC-1/FC-1
        x2 = (StdAlphaEnergy(nAgeStdPreferred2) / StdAlphaEnergy(nAgeStdPreferred1)) 'IFCT/FC-1
        y1 = 1 'FC-1/FC-1
        '235Uc
        y2235 = std207235Age(nAgeStdPreferred2) / measAgeStdPreferred2
        m = (y2235 - y1) / (x2 - x1)
        b = y2235 - x2 * m
        ffage(2) = ffage(2) * (m * UAlphaEnergy / SAlphaEnergy + b)
        If ffage(2) < minff(2) Then
            ffage(2) = minff(2)
        End If
        '238U
```

```
        y2238 = std206238Age(nAgeStdPreferred2) / measAgeStdPreferred2
        m = (y2238 - y1) / (x2 - x1)
        b = y2238 - x2 * m
        ffage(3) = ffage(3) * (m * UAlphaEnergy / SAlphaEnergy + b)
        If ffage(3) < minff(3) Then
            ffage(3) = minff(3)
        End If
        '232Th
        y2232 = std208232Age(nAgeStdPreferred2) / measAgeStdPreferred2
        m = (y2232 - y1) / (x2 - x1)
        b = y2232 - x2 * m
        ffage(5) = ffage(5) * (m * UAlphaEnergy / SAlphaEnergy + b)
        If ffage(5) < minff(5) Then
            ffage(5) = minff(5)
        End If
    End If
End Sub
```

Appendix 3: Functions for Calculating U-Pb Ages

The following functions are used by UPbICP to calculate U-Pb ages for their respective isotopic ratios (input parameter *ratio*).

```
Function Age207235(ratio)
    'dim ratio,Age207235 implicit
    If ratio < 0 Then
        Age207235 = 0
    Else
        Age207235 = (1 / lamdaU235) * Log(ratio + 1)
    End If
End Function

Function Age206238(ratio)
    'dim ratio,Age206238 implicit
    If ratio < 0 Then
        Age206238 = 0
    Else
        Age206238 = (1 / lamdaU238) * Log(ratio + 1)
    End If
End Function
```

For 207Pb/206Pb ratios, U-Pb ages must be calculated by iteration. To avoid this, UPbICP reads an array *ratio207206_data* indexed from i = 1 to 45600 and containing 207Pb/206Pb ratios corresponding to i x 10^5 years for each index i.

```
Function Age207206(ratio)
    'dim ratio,Age207206 implicit
    Dim i, m, b
    If ratio <= Pb207206init Or ratio > ratio207206_data(45600) Then
        Age207206 = 0
    Else
        For i = 1 To 45600
            If ratio < ratio207206_data(i) Then
                m = 0.1 / (ratio207206_data(i) - ratio207206_data(i - 1))
                b = CSng(i - 1) * 0.1 - ratio207206_data(i - 1) * m
                Age207206 = m * ratio + b
                Exit Function
            End If
        Next i
    End If
End Function
```

Table 1. ICP-MS and laser ablations system operating conditions and data acquisition parameters.

ICP-MS: operating conditions	
Instrument	Finnigan Element II Magnetic Sector ICP-MS
Forward power	1.25 kW
Reflected power	<5 W
Plasma gas	Ar
Coolant flow	15 l/min
Carrier flow	1.0 l/min (Ar) 0.8 l/min (He)
Auxiliary flow	1.0 l/min
ICP-MS: data acquisition parameters	
Dwell time	24 milliseconds per peak point
Points per peak	3
Mass window	5%
Scans	255
Data acquisition time	38.5 sec
Data acquisition mode	E scanning
Isotopes measured	^{202}Hg, ^{204}Pb, ^{206}Pb, ^{207}Pb, ^{208}Pb, ^{232}Th, ^{235}U, ^{238}U (scans ~60-250) ^{28}Si, ^{91}Zr (scans 251-255)
Laser ablation system: operating conditions	
Laser type	New Wave Neodymium: YAG
Wavelength	213 nm
Laser mode	Q switched
Laser output power	10 J/cm^2
Laser warm up time	6 sec
Shot repetition rate	5 Hz
Sampling scheme	spot (20 μm)

Table 2. Age standards used at A2Z.

Standard	Standard	U-Pb age (±2σ)	Reference
FC-1	Duluth complex	1099.0 ± 0.6 Ma	Paces and Miller, 1993
FC-5z	Duluth complex	1099.0 ± 0.6 Ma (assumed equal to FC-1)	Paces and Miller, 1993
TR	Tardree Rhyolite	61.23 ± 0.11 Ma	Dave Chew, personal communication
IFCan	Fish Canyon Tuff	28.201 ± 0.012 Ma	Kuiper et al., 2008
R33	Braintree complex	418.9 ± 0.4 Ma	Black et al., 2004
TEM2	Temora 2, Middledale gabbroic diorite	416.78 ± 0.33 Ma	Black et al., 2004
MD	Mount Dromedary	99.12 ± 0.14 Ma	Renne et al., 1998
200-01	Tioga Bed K bentonite	391.4 ± 1.8 Ma	Tucker et al., 1998

Table 3. Weighted mean U and Th ppm values assumed for FC-1 (Paces and Miller, 1993)

mg	Uppm	Thppm	mg*Uppm	mg*Thppm
125	420	258	52500	32250
120	1510	905	181200	108600
155	299	170	46345	26350
118	218	115	25724	13570
133	292	176	38836	23408
sum mg			mg*Uppm/sum mg	mg*Thppm/sum mg
651		weighted mean	529.35	313.64



Figure 1a. Transmitted and reflected light image of spot 1713b_20 at 625x magnification. The plane of focus is on the polished and etched zircon surface.



Figure 1b. Transmitted light image of spot 1713b_20 at 625x magnification. The plane of focus is on the lower prismatic crystal surface of the zircon grain.



Figure 1c. Transmitted light image of spot 1713b_20 at 625x magnification. The plane of focus is on the interior of the zircon grain and it shows the boundary between the old core and young rim.



Figure 2a. 206Pb data for spot 1713b_20. Left-hand yellow curve represents the polynomial fitted to the background intensities. Right-hand yellow curve is represents the sum of Gaussian equations fitted to the signal intensities.



Figure 2b. 207Pb data for spot 1713b_20.



Figure 2c. 238U data for spot 1713b_20.



	A	B	C	D	E	F	G	H	I	J	K
1	Standard	Uncorrect	Corrected	Uncorrect	Corrected	Accepted	Uncorrect	Corrected	Averages	regressior	regression
2	FC-1	1099	1100.37	1.0942	1.0921	1099.00	1.000000	1.001247	0.978219	m	3.47E-02
3	TR	59.08	60.67	0.0390	0.0376	61.23	0.964886	0.990854	1.000774	b	9.67E-01
4	FC-5z	1096.69	1115.09	0.5068	0.5088	1099.00	0.997898	1.014641		UPbICP co	59.31
5	IFCan	27.53	28.28	0.0185	0.0173	28.20	0.976207	1.002801			
6	MD	94.53	97.84	0.0879	0.0854	99.12	0.953692	0.987086			
7	R33					418.90					
8	TEM2	407.04	420.12	0.1402	0.1376	416.78	0.976630	1.008014			

Figure 3. Age data for the secondary age standards before and after correction for relative α-radiation damage in session z031110.



A2Z Report ICP_NEW

U-Pb zircon analytical results

Min Fit Quality	0.0500	Min Clean Quality	0.3333	Min 204Pb Quality	0.3333	Min Uppm Quality	9	Mineral (or "n.a.")	Zrn	Ctrl+r reserved

La Sal Mtns. 17103b 2_031110

La Sal Mtns. 17103b 2_031110

	Measured Isotopic Ratios						U and Th			
Analysis_#	207Pb/235Uc*	207Pb/235Uc* (±2σ)	206Pb/238U	206Pb/238U (±2σ)	207Pb/206Pb	207Pb/206Pb (±2σ)	U (ppm)	Th (ppm)	U/Th	Analysis_#	207Pb/2
17103azl_1_Zrn	4.1957	0.1570	0.2950	0.0166	0.1032	0.0048	687.07	979.73	0.701	17103azl_1	1673
17103azl_10_Zrn	3.8135	0.1445	0.2715	0.0157	0.1017	0.0049	3636.43	1934.22	1.880	17103azl_10	1595
17103azl_11_Zrn	3.5099	0.1321	0.2541	0.0144	0.1002	0.0047	1334.59	188.74	7.071	17103azl_11	1529
17103azl_12_Zrn	2.6644	0.1003	0.2271	0.0128	0.0849	0.0040	31.81	12.90	2.466	17103azl_12	1318
17103azl_13_Zrn	3.1785	0.1189	0.2476	0.0140	0.0943	0.0044	631.51	220.15	2.869	17103azl_13	1451
17103azl_14_Zrn	2.5864	0.0969	0.2156	0.0122	0.0870	0.0041	447.13	186.40	2.399	17103azl_14	1296
17103azl_15_Zrn	3.8122	0.1429	0.2683	0.0152	0.1030	0.0048	2289.18	2674.21	0.856	17103azl_15	1595
17103azl_16_Zrn	3.8980	0.1459	0.2768	0.0156	0.1021	0.0048	2048.37	1132.71	1.808	17103azl_16	1613
17103azl_17_Zrn	undefined	undefined	undefined	undefined	undefined	undefined	1671.23	1529.26	1.093	17103azl_17	undef
17103azl_18_Zrn	undefined	undefined	undefined	undefined	undefined	undefined	206.48	82.27	2.510	17103azl_18	undef
17103azl_19_Zrn	0.0275	0.0010	0.0041	0.0002	0.0480	0.0023	2572.45	2283.89	1.126	17103azl_19	27.
17103azl_2_Zrn	3.9475	0.1479	0.2872	0.0162	0.1004	0.0047	134.88	48.54	2.778	17103azl_2	1623
17103azl_20_Zrn	3.8664	0.1454	0.2721	0.0154	0.1032	0.0049	962.57	124.55	7.729	17103azl_20	1606
17103azl_21_Zrn	2.7952	0.1102	0.2109	0.0124	0.0923	0.0045	419.82	30.60	13.718	17103azl_21	1354
17103azl_22_Zrn	2.7901	0.1048	0.2255	0.0127	0.0896	0.0042	592.72	115.56	5.129	17103azl_22	1352
17103azl_23_Zrn	0.0284	0.0012	0.0042	0.0002	0.0494	0.0025	869.56	357.59	2.432	17103azl_23	28.
17103azl_24_Zrn	undefined	undefined	undefined	undefined	undefined	undefined	111.72	22.63	4.937	17103azl_24	undef
17103azl_25_Zrn	3.3625	0.1266	0.2562	0.0145	0.0991	0.0047	259.50	32.63	7.952	17103azl_25	1495
17103azl_3_Zrn	3.8796	0.1458	0.2757	0.0156	0.1022	0.0048	1976.81	106.54	18.554	17103azl_3	1609
17103azl_4_Zrn	4.4894	0.1680	0.3058	0.0173	0.1063	0.0050	148.12	96.79	1.530	17103azl_4	1729
17103azl_5_Zrn	4.4558	0.1664	0.2994	0.0169	0.1080	0.0051	674.55	170.70	3.952	17103azl_5	1722

Figure 4. Worksheet: Data



UPbICPReport_17103b_2_031110 [Compatibility Mode] - Microsoft Excel non-commercial use

J9: not ready

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
1	Preferred Age (Ma)	Absolute Error (±2σ)	Age Type	Analysis #	Product of Qualities		workspace	workspace	workspace	workspace	workspace	workspace	workspace	x hist age
2	21.27	1.90	lower int.	17103azs_45	0.8004		# ages	72	28					0.00
3	21.74	1.24	Pb/U wmean conc.	17103azs_27	0.7766		xmin_data	18.42		transfer data		rescale plots		0.00
4	23.38	1.33	Pb/U wmean conc.	17103azl_9	0.4314		xmax_data	2060.58	preferred					20.00
5	23.43	1.76	lower int.	17103azs_36	0.2769	set	xmax_plot	2000.00	2000.00			wmean	mean	20.00
6	23.71	1.82	lower int.	17103b_3	0.5676	set	xbinwidth_	20.00	20.00			26.35	26.27	20.00
7	23.72	1.57	Pb/U wmean conc.	17103b_14	0.8145	set	xstep_plot	0.5	0.5			2σ+FFerr	2σ stddev	40.00
8	24.00	2.04	lower int.	17103azs_32	0.6877					Progress		1.27	2.65	40.00
9	24.25	2.34	lower int.	17103azs_41	0.5054					not ready		2σ	2σ stderr	40.00
10	25.15	1.66	Pb/U wmean conc.	17103b_9	0.8431		area histog	1440.00		Maximum Row		0.31	0.50	60.00
11	25.60	1.45	Pb/U wmean conc.	17103b_23	0.5022		area pdf	1440.01		73		MSWDw	P(MSWDw)	60.00
12	26.06	1.49	Pb/U wmean conc.	17103azs_49	0.8162		area gauss	1430.29	1430.26			2.12	0.004676	60.00
13	26.10	1.57	Pb/U wmean conc.	17103azs_43	0.5089							start_row	end_row	80.00
14	26.23	1.52	Pb/U wmean conc.	17103b_24	0.9797		ngauss	12		maximum row	set	5	32	80.00
15	26.32	2.06	lower int.	17103azs_35	0.5219	set	dym_factor	2.25				# of 1σ Absolute Errors		80.00
16	26.42	1.68	Pb/U wmean conc.	17103b_8	0.9977		chisqm	0.0035316			set	3.00		100.00
17	26.44	1.51	Pb/U wmean conc.	17103b_2	0.8305		chisqbestm	0.0035316						100.00
18	26.44	1.52	Pb/U wmean conc.	17103b_5	0.6612	set	chisqtol	0.1000		deconvolve		statistics		100.00
19	26.60	1.51	Pb/U wmean conc.	17103azl_19	0.6268									120.00
20	26.78	1.86	lower int.	17103b_15	0.7152									120.00
21	26.79	1.54	Pb/U wmean conc.	17103azl_23	0.5228		gauss	amplitude	mean	std dev	data fraction		mean	120.00
22	26.86	1.53	Pb/U wmean conc.	17103b_18	0.7283		1	27.132	21.590	0.837	2.81%	Uppm	631.77	140.00
23	26.92	1.53	Pb/U wmean conc.	17103azs_28	0.8270		2	64.663	23.589	0.830	6.65%	Thppm	467.29	140.00
24	27.00	1.54	Pb/U wmean conc.	17103azs_33	0.7205		3	30.524	25.489	1.506	5.70%	U/Th	1.975	140.00
25	27.01	1.54	Pb/U wmean conc.	17103azs_29	0.8514		4	176.130	26.810	1.114	24.32%	alpha	0.0254	160.00
26	27.15	1.56	Pb/U wmean conc.	17103azs_37	0.7874		5	42.509	29.318	2.251	11.86%		Q Filters	160.00
27	27.48	1.57	Pb/U wmean conc.	17103b_4	0.9975		6	7.091	44.150	1.591	1.40%	Fit	0.0500	160.00
28	27.54	1.57	Pb/U wmean conc.	17103b_6	0.6853		7	0.263	981.700	42.878	1.40%	Clean	0.3333	180.00
29	27.59	1.58	Pb/U wmean conc.	17103b_12	0.8529		8	1.226	1290.650	56.656	8.61%	204Pb	0.3333	180.00
30	27.63	1.62	Pb/U wmean conc.	17103b_13	0.5923		9	0.723	1465.360	99.559	8.92%	U ppm	9	180.00
31	27.82	1.60	Pb/U wmean conc.	17103b_25	0.3160		10	0.447	1615.032	30.051	1.66%	Mineral	Zrn	200.00

Figure 5. Worksheet: Calculate



Figure 6a. Worksheet: Plots



Figure 6b. Worksheet: Plots



	A	B	C	D
1	**Primary Standard**			
2	UPbICP Standard Index	6	Standard	F
3	Accepted Age	1099		
4	**Isotopic Ratio**	**Mean Age**	**FF Relative Error**	
5	207Pb/235U	1096.29	0.0117	
6	207Pb/235Uc	1097.07	0.0186	
7	206Pb/238U	1096.32	0.0281	
8	207Pb/206Pb	1099.00	0.0233	
9	208Pb/232Th	1097.88	0.0679	
10	**Secondary Standard**			
11	UPbICP Standard Index	2	Standard	T
12	Accepted Age	61.23		
13	Uncorrected Age	59.05		
14	**28Si 91Zr All Standards**			
15	Mineral	Zrn		
16	**Statistics**	**91Zr/28Si**	**28Si**	**91Zr**
17	median	0.669	4064708	2720961
18	standard deviation	0.071	225898	245894
19	minimum	not applicable	3588061	2220840
20	maximum	not applicable	4391908	3083139

Figure 7. Worksheet: Standard_Statistics



Session	Grain	Preferred (Measured) Age (Ma)	Absolute Error (±2σ)	Age Type	Accepted Age (Ma)	Absolute Error (±2σ)	(Accepted - Preferred) / Accepted	Average (perfect=0)	Accepted / Measured	Average (perfect=1)
z031110	TR_11	54.00	4.15	lower int.	61.23	0.11	0.1181	-0.0004	1.1339	1.0006
z031110	TR_15	57.30	3.41	Pb/U wmean conc.	61.23	0.11	0.0641		1.0685	
z031110	TR_10	59.01	3.37	Pb/U wmean conc.	61.23	0.11	0.0362		1.0376	
z031110	TR_12	59.10	3.41	Pb/U wmean conc.	61.23	0.11	0.0347		1.0360	
z031110	TR_16	59.65	3.41	Pb/U wmean conc.	61.23	0.11	0.0259		1.0266	
z031110	TR_18	60.07	3.43	Pb/U wmean conc.	61.23	0.11	0.0190		1.0194	
z031110	TR_21	60.10	3.43	Pb/U wmean conc.	61.23	0.11	0.0185		1.0188	
z031110	TR_22	60.13	3.43	Pb/U wmean conc.	61.23	0.11	0.0180		1.0184	
z031110	TR_19	60.51	3.46	Pb/U wmean conc.	61.23	0.11	0.0117		1.0119	
z031110	TR_8	60.68	3.46	Pb/U wmean conc.	61.23	0.11	0.0090		1.0090	
z031110	TR_13	60.88	3.47	Pb/U wmean conc.	61.23	0.11	0.0057		1.0057	
z031110	TR_6	60.91	3.54	Pb/U wmean conc.	61.23	0.11	0.0053		1.0053	
z031110	TR_17	60.95	3.48	Pb/U wmean conc.	61.23	0.11	0.0046		1.0046	
z031110	TR_20	61.06	4.12	Pb/U wmean conc.	61.23	0.11	0.0028		1.0028	
z031110	TR_7	61.33	3.50	Pb/U wmean conc.	61.23	0.11	-0.0017		0.9983	
z031110	TR_3	62.09	3.61	Pb/U wmean conc.	61.23	0.11	-0.0140		0.9862	
z031110	TR_9	62.45	3.58	Pb/U wmean conc.	61.23	0.11	-0.0199		0.9805	
z031110	TR_1	62.50	4.49	lower int.	61.23	0.11	-0.0207		0.9797	
z031110	TR_5	63.25	3.63	Pb/U wmean conc.	61.23	0.11	-0.0330		0.9681	
z031110	TR_4	63.86	3.67	Pb/U wmean conc.	61.23	0.11	-0.0430		0.9588	
z031110	TR_14	64.72	3.69	Pb/U wmean conc.	61.23	0.11	-0.0570		0.9461	
z031110	FC5_8	1078.60	66.15	Pb/U wmean conc.	1099.00	0.60	0.0186		1.0189	
z031110	FC5_9	1087.72	66.69	Pb/U wmean conc.	1099.00	0.60	0.0103		1.0104	
z031110	FC5_6	1096.77	67.27	Pb/U wmean conc.	1099.00	0.60	0.0020		1.0020	
z031110	FC5_7	1105.31	67.84	Pb/U wmean conc.	1099.00	0.60	-0.0057		0.9943	
z031110	FC5_1	1129.00	69.40	Pb/U wmean conc.	1099.00	0.60	-0.0273		0.9734	
z031110	FC5_2	1132.16	69.61	Pb/U wmean conc.	1099.00	0.60	-0.0302		0.9707	
z031110	FC5_3	1132.50	69.66	Pb/U wmean conc.	1099.00	0.60	-0.0305		0.9704	
z031110	FC5_4	1137.66	69.98	Pb/U wmean conc.	1099.00	0.60	-0.0352		0.9660	

Figure 8. Worksheet: A2Z_Standards



Figure 9. Worksheet: Comment



B4 =Calculate!L6

	A	B	C	D
1	**Results**			
2	# ages total	72		
3	# ages in population	28		
4	weighted mean age ± 2σ (Ma; incl. fractionation error)	26.35	±	1.27
5	MSWDw	2.12		
6	P(MSWDw)	0.004676		
7				
8				
9				
10	**Settings**			
11	Min Fit Quality	0.0500		
12	Min Clean Quality	0.3333		
13	Min 204Pb Quality	0.3333		
14	Min Uppm Quality	9		
15	Mineral	Zrn		
16	# of 1σ Absolute Errors	3.00		

Figure 10. Worksheet: Results_Settings



	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG	AH	AI	AJ
1	set	run file	c:\UPbICP\UPbZrSiZ031110.csv					bkg_Fit	isotope	coefficien	coefficien	coefficien	1σ coefficie	1σ coefficie	1σ coefficie	value at n	raw standard devi	
2		file type	2						1	6834.43	-14.5273	0	345.5884	14.5238	1	6253.336	1040.544	
3		bkg order	1		refresh				2	1438.32	0	0	48.73386	1	1	1438.32	296.4365	
4		std sig order	0						3	381.6464	-7.66632	0	80.51627	3.351477	1	74.99352	239.106	
5		session order	3						4	80.02804	-1.19234	0	29.18907	1.234741	1	32.33433	86.39104	
6		unk sig order	5						5	205.6142	-2.30106	0	49.15406	2.112682	1	113.5719	152.4879	
7		bkg begin	1						6	175.2195	-0.1235	0	49.41291	2.05779	1	170.2793	144.6663	
8		bkg end	40						7	0	0	0	0	0	0	0	1	
9		sig begin	90						8	789.8837	-16.3902	0	160.8436	7.093056	1	134.2737	492.2503	
10		sig end	250						9	5.728761	-0.11887	0	1.166546	0.051444	1	0.973854	3.570136	
11																		
12	set	disk:\folder	c:\UPbICP\															
13	set	session	PbZrSiZ031110					bkg+signa	isotope	degrees c	chisq	Q	1σ (stdde	Gaussian	amplitude	mean (sca	standard deviatio	
14	set	root (CASE!)	17103b															
15	set/scroll	grain number	20		refresh			202Hg	1	121	102.6774	0.884875	93.03654	1	3650.319	70.62784	49.08384	
16	set	extension	.txt											2	3522.134	85.35723	58.29498	
17	set	grains	25											3	2622.897	122.9659	38.15323	
18		filename	c:\UPbICP\UPbZrSiZ031110\17103b_20.txt											4	2188.742	134.2201	21.56248	
19		scans	250											5	2257.01	166.2178	24.68312	
20	set	Std/Unk	Unk											6	2579.289	176.3416	29.32299	
21					deleted	measured	fit							7	2190.113	203.6227	26.52433	
22		title	17103b_20	202Hg	0.65	133.23	133.23							8	3085.601	228.7843	71.82386	
23				204Hg+204Pb	0.97	31.28	31.28							9	3330.49	268.7222	54.38105	
24		reading file	6	206Pb	0.00	6397.83	6397.80							10	3219.355	271.0428	69.00866	
25		status	ready	207Pb	1.82	664.27	664.27											
26				208Pb	0.00	155.85	155.85	204Hg+20	2	125	102.6614	0.928427	48.17649	1	518.6675	57.24311	78.13519	
27				232Th	0.00	2430.60	2430.63							2	442.5729	88.18353	42.61623	
28				235U	0.00	198.36	198.36							3	702.3949	95.76696	42.05452	
29				238U	0.00	27217.30	27217.62							4	541.595	138.892	80.49998	
30				235Uc	0.00	197.40	197.39							5	462.0818	167.9482	35.20547	
31				sum	n.a.	37426.13	37426.45							6	456.0786	162.751	35.11963	
32														7	425.6167	217.004	79.70284	

Figure 11. Worksheet: Raw_Data



Figure 12. Worksheet: Raw_Data_Plots



Figure 13a. Worksheet: bkg_Corrected_Data_Plots for spot 17103b_20.



Figure 13b. Worksheet: bkg_Corrected_Data_Plots for spot 17103b_8.



Figure 14a. Worksheet: Discordia



Figure 14b. Worksheet: Discordia



	A	B	C	D
1	**Primary Standard**			
2	UPbICP Standard Index	6	Standard	F
3	Accepted Age	1099		
4	**Isotopic Ratio**	**Mean Age**	**FF Relative Error**	
5	207Pb/235Uc	1096.40	0.0112	
6	207Pb/235Uc*	1096.83	0.0185	
7	206Pb/238U	1095.96	0.0279	
8	207Pb/206Pb	1099.00	0.0232	
9	208Pb/232Th	1097.62	0.0679	
10	**Secondary Standard**			
11	UPbICP Standard Index	2	Standard	T
12	Accepted Age	61.23		
13	Uncorrected Age	59.31		
14	**28Si 91Zr All Standards**			
15	Mineral	Zrn		
16	**Statistics**	**91Zr/28Si**	**28Si**	**91Zr**
17	median	0.669	4064708	2720961
18	standard deviation	0.071	225898	245894
19	minimum	not applicable	3588061	2220840
20	maximum	not applicable	4391908	3083139

Figure 15. Worksheet: Standard_Statistics

APPENDIX C

A2Z Report 1131-02
U-Pb zircon analytical results

Min Fit Quality: 0.0500 | Min Clean Quality: 0.3333 | Min 204Pb Quality: 0.3333 | Min Upper Quality: 3 | Mineral (or "+.+.") | 2σ | Other reserved

Morrison/Cedar Mtn

Measured Isotopic Ratios

Analysis_#	207Pb/235Uc* (±2σ)	207Pb/235Uc* (±2σ)	206Pb/238U (±2σ)	206Pb/238U (±2σ)	207Pb/206Pb (±2σ)	207Pb/206Pb (±2σ)
113102_1_2m	4.6218	0.2934	0.2934	0.0095	0.0995	0.0013
113102_2_2m	0.6417	0.0171	0.0814	0.0022	0.0571	0.0013
113102_3_2m	0.5071	0.0092	0.0675	0.0018	0.0544	0.0012
113102_4_2m	4.8146	0.0866	0.3257	0.0086	0.1094	0.0024
113102_5_2m	2.2635	0.0399	0.2064	0.0054	0.0796	0.0018
113102_6_2m	0.5753	0.0108	0.0737	0.0020	0.0565	0.0013
113102_7_2m	1.8051	0.0330	0.1807	0.0048	0.0751	0.0017
113102_8_2m	0.3109	0.0060	0.0457	0.0012	0.0494	0.0011
113102_9_2m	0.7442	0.0140	0.0992	0.0026	0.0598	0.0014
113102_10_2m	0.9299	0.0171	0.1106	0.0029	0.0617	0.0014
113102_11_2m	0.1702	0.0036	0.0243	0.0006	0.0507	0.0013
113102_12_2m	2.0683	0.0365	0.1917	0.0051	0.0783	0.0017
113102_13_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_14_2m	0.8614	0.0154	0.1023	0.0027	0.0610	0.0014
113102_15_2m	0.8940	0.0176	0.1141	0.0030	0.0632	0.0014
113102_16_2m	0.5207	0.0099	0.0663	0.0018	0.0569	0.0013
113102_17_2m	0.7396	0.0131	0.0914	0.0024	0.0587	0.0013
113102_18_2m	0.1560	0.0031	0.0225	0.0006	0.0482	0.0012
113102_19_2m	0.5001	0.0094	0.0666	0.0018	0.0544	0.0013
113102_20_2m	0.1773	0.0036	0.0256	0.0007	0.0502	0.0012
113102_21_2m	0.3758	0.0139	0.0241	0.0006	0.0529	0.0043
113102_22_2m	0.1757	0.0032	0.0229	0.0007	0.0495	0.0011
113102_23_2m	3.8753	0.0695	0.2796	0.0074	0.1007	0.0022
113102_24_2m	0.1500	0.0031	0.0215	0.0006	0.0506	0.0012
113102_25_2m	1.7361	0.0305	0.1712	0.0045	0.0736	0.0016
113102_26_2m	0.1266	0.0024	0.0188	0.0005	0.0488	0.0011
113102_27_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_28_2m	0.1590	0.0030	0.0232	0.0006	0.0497	0.0011
113102_29_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_30_2m	0.7513	0.0134	0.0957	0.0024	0.0584	0.0013
113102_31_2m	0.9002	0.0161	0.1066	0.0028	0.0613	0.0014
113102_32_2m	1.5460	0.0287	0.1582	0.0042	0.0709	0.0016
113102_33_2m	3.2183	0.0567	0.2538	0.0067	0.0921	0.0020
113102_34_2m	0.6092	0.0131	0.0830	0.0022	0.0575	0.0014
113102_35_2m	4.5768	0.0811	0.3060	0.0082	0.1078	0.0024
113102_36_2m	0.4631	0.0086	0.0635	0.0017	0.0529	0.0012
113102_37_2m	0.4631	0.0084	0.0618	0.0016	0.0542	0.0012
113102_38_2m	2.1153	0.0407	0.2078	0.0055	0.0808	0.0018
113102_39_2m	0.9542	0.0170	0.1120	0.0030	0.0618	0.0014
113102_40_2m	0.1729	0.0034	0.0246	0.0007	0.0510	0.0012
113102_41_2m	0.1500	0.0029	0.0214	0.0006	0.0508	0.0012
113102_42_2m	0.1505	0.0030	0.0206	0.0005	0.0529	0.0013
113102_43_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_44_2m	2.0832	0.0367	0.1924	0.0051	0.0785	0.0017
113102_45_2m	0.4882	0.0088	0.0652	0.0017	0.0544	0.0012
113102_46_2m	13.1415	0.2309	0.5159	0.0136	0.1848	0.0041
113102_47_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_48_2m	0.1500	0.0029	0.0215	0.0006	0.0506	0.0011
113102_49_2m	0.1800	0.0029	0.0235	0.0006	0.0484	0.0011
113102_50_2m	0.7856	0.0139	0.0961	0.0025	0.0593	0.0013

Morrison/Cedar Mtn

U and Th

Analysis_#	U (ppm)	Th (ppm)	U/Th
113102_1_2m	680.59	417.81	1.654
113102_2_2m	387.06	466.56	0.830
113102_3_2m	401.32	105.46	3.806
113102_4_2m	987.47	414.66	2.381
113102_5_2m	668.02	336.90	1.983
113102_6_2m	293.71	157.39	1.866
113102_7_2m	391.58	127.47	3.072
113102_8_2m	1.00.00	1.00.00	1.000
113102_9_2m	95.25	71.12	1.339
113102_10_2m	246.89	253.77	0.973
113102_11_2m	380.37	259.82	1.487
113102_12_2m	638.53	425.48	1.501
113102_13_2m	1815.55	1158.21	1.576
113102_14_2m	421.57	184.24	2.289
113102_15_2m	462.91	405.41	1.142
113102_16_2m	1825.55	1158.01	1.576
113102_17_2m	421.55	140.74	0.379
113102_18_2m	566.71	461.61	1.228
113102_19_2m	964.30	242.84	3.971
113102_20_2m	1.00.00	1.00.00	1.000
113102_21_2m	383.26	763.14	0.502
113102_22_2m	430.58	298.86	1.441
113102_23_2m	374.29	2611.78	1.292
113102_24_2m	549.13	592.83	0.926
113102_25_2m	2336.55	6113.09	0.381
113102_26_2m	165.65	132.49	1.250
113102_27_2m	1.00.00	1.00.00	1.000
113102_28_2m	665.23	238.22	2.792
113102_29_2m	993.40	1308.32	0.759
113102_30_2m	2727.74	535.52	5.094
113102_31_2m	934.75	874.89	1.069
113102_32_2m	1.00.00	1.00.00	1.000
113102_33_2m	331.58	325.70	1.137
113102_34_2m	366.74	407.73	0.899
113102_35_2m	79.64	30.15	2.343
113102_36_2m	298.81	207.28	1.442
113102_37_2m	74.50	66.07	1.128
113102_38_2m	429.62	341.52	1.258
113102_39_2m	1.00.00	1.00.00	1.000
113102_40_2m	421.40	609.27	0.521
113102_41_2m	730.71	227.30	3.215
113102_42_2m	542.68	772.03	0.703
113102_43_2m	570.10	525.22	1.085
113102_44_2m	1.00.00	1.00.00	1.000
113102_45_2m	320.08	88.45	3.619
113102_46_2m	5028.91	1964.13	2.562
113102_47_2m	undefined	undefined	undefined
113102_48_2m	621.56	7.70	80.704
113102_49_2m	136.37	296.34	0.460
113102_50_2m	719.99	721.83	0.997

Morrison/Cedar Mtn

Apparent Ages

Analysis_#	207Pb/235Uc* (±2σ)	207Pb/235Uc* (±2σ)	206Pb/238U (±2σ)	206Pb/238U (±2σ)	207Pb/206Pb (±2σ)	207Pb/206Pb (±2σ)
113102_1_2m	1639.61	1658.43	1658.43	13.86	1614.08	28.04
113102_2_2m	503.34	11.64	504.53	13.86	497.04	49.44
113102_3_2m	416.51	9.34	420.88	11.55	388.82	50.13
113102_4_2m	1704.77	86.10	1817.36	56.32	1788.97	39.84
113102_5_2m	1200.99	40.15	1209.52	35.07	1186.70	43.22
113102_6_2m	461.43	10.97	458.23	12.64	472.22	50.69
113102_7_2m	1068.79	33.26	1070.82	30.77	1069.90	44.09
113102_8_2m	1.00.00	1.00.00	1.000	1.000	1.000.00	1.000
113102_9_2m	564.85	14.17	558.94	15.35	595.62	49.51
113102_10_2m	672.84	17.32	675.98	18.89	663.57	47.95
113102_11_2m	158.56	3.61	154.46	4.16	225.57	58.93
113102_12_2m	1138.36	36.77	1130.42	32.55	1153.90	43.46
113102_13_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_14_2m	630.84	15.57	628.02	17.43	640.62	47.48
113102_15_2m	700.74	17.76	696.58	19.38	713.68	46.52
113102_16_2m	425.63	10.07	413.73	11.30	489.09	50.90
113102_17_2m	562.16	13.22	563.59	15.53	557.32	47.72
113102_18_2m	141.87	3.10	143.70	3.87	119.14	57.10
113102_19_2m	411.77	9.56	415.75	11.46	388.56	51.58
113102_20_2m	160.75	3.63	162.87	4.41	205.31	55.85
113102_21_2m	164.46	14.07	153.47	4.19	324.96	177.50
113102_22_2m	165.20	3.21	164.66	4.42	189.94	51.55
113102_23_2m	1608.56	69.39	1589.17	47.78	1637.62	41.14
113102_24_2m	136.07	3.10	137.61	3.87	102.83	57.10
113102_25_2m	1022.00	30.72	1018.89	28.84	1029.43	44.09
113102_26_2m	121.07	2.40	120.22	3.21	139.04	53.35
113102_27_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_28_2m	149.82	3.00	147.80	3.97	181.56	52.83
113102_29_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_30_2m	568.99	13.55	590.65	15.59	580.86	47.80
113102_31_2m	651.81	16.25	652.69	18.14	648.88	47.36
113102_32_2m	948.90	28.88	946.73	26.95	955.21	46.33
113102_33_2m	1461.57	56.75	1457.89	43.05	1469.66	41.50
113102_34_2m	514.15	13.25	513.99	14.19	511.89	52.17
113102_35_2m	1745.05	80.69	1730.93	52.49	1763.29	40.18
113102_36_2m	398.41	8.93	386.71	10.54	378.41	50.20
113102_37_2m	399.41	8.93	386.71	10.54	378.41	50.20
113102_38_2m	1214.98	40.95	1217.17	35.29	1217.01	42.95
113102_39_2m	680.27	17.15	684.46	19.07	668.17	47.10
113102_40_2m	161.90	3.45	156.44	4.23	238.08	54.35
113102_41_2m	137.61	2.90	136.69	3.73	161.42	53.54
113102_42_2m	419.45	9.50	409.84	15.19	461.82	51.54
113102_43_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_44_2m	1143.29	36.87	1134.52	32.73	1158.75	43.36
113102_45_2m	403.67	8.90	406.88	11.12	386.31	49.81
113102_46_2m	2689.87	221.88	2681.71	87.35	2696.38	36.06
113102_47_2m	undefined	undefined	undefined	undefined	undefined	undefined
113102_48_2m	137.61	2.90	136.69	3.73	161.42	53.54
113102_49_2m	150.71	2.92	149.69	4.08	164.55	51.75
113102_50_2m	588.66	14.09	591.65	16.36	577.84	47.67

Morrison/Cedar Mtn

Preferred Ages

Analysis_#	Preferred Age (Ma) (±2σ)	Absolute Error (±2σ)	Age Type	Fit Quality 1=best	Clean Quality 1=best	204Pb Quality 1=best	2σ Conc. Scans
113102_1_2m	1639.61	28.04	Pb/Pb w/mean conc.	0.0000	1.0000	1.0000	88
113102_2_2m	504.53	13.86	Pb/U w/mean conc.	0.6726	0.5000	1.0000	111
113102_3_2m	420.99	11.55	Pb/U w/mean conc.	0.8407	1.0000	1.0000	110
113102_4_2m	1789.97	39.84	Pb/Pb w/mean conc.	0.9605	1.0000	1.0000	133
113102_5_2m	1209.52	35.07	Pb/U w/mean conc.	0.8872	1.0000	1.0000	125
113102_6_2m	458.23	12.64	Pb/U w/mean conc.	0.7505	1.0000	1.0000	55
113102_7_2m	1070.82	30.77	Pb/U w/mean conc.	0.9018	1.0000	1.0000	95
113102_8_2m	1.00.00	1.000	Pb/U w/mean conc.	0.0000	1.0000	1.0000	1
113102_9_2m	558.94	15.35	Pb/U w/mean conc.	0.6383	1.0000	1.0000	103
113102_10_2m	675.98	18.89	Pb/U w/mean conc.	0.6723	1.0000	1.0000	55
113102_11_2m	154.46	4.16	Pb/U w/mean conc.	0.6767	0.5000	1.0000	54
113102_12_2m	1130.42	32.55	Pb/U w/mean conc.	0.9639	1.0000	1.0000	97
113102_13_2m	undefined	undefined	undefined	0.9447	1.0000	1.0000	0
113102_14_2m	628.02	17.43	Pb/U w/mean conc.	0.9519	1.0000	1.0000	91
113102_15_2m	696.58	19.38	Pb/U w/mean conc.	0.9400	1.0000	1.0000	115
113102_16_2m	413.73	11.30	Pb/U w/mean conc.	0.3943	1.0000	1.0000	71
113102_17_2m	563.59	15.53	Pb/U w/mean conc.	0.9964	1.0000	1.0000	106
113102_18_2m	143.70	3.87	Pb/U w/mean conc.	0.9702	1.0000	1.0000	55
113102_19_2m	415.75	11.46	Pb/U w/mean conc.	0.7212	1.0000	1.0000	37
113102_20_2m	162.87	4.41	Pb/U w/mean conc.	0.8682	1.0000	1.0000	10
113102_21_2m	153.47	4.19	Pb/U w/mean conc.	1.0000	1.0000	0.8526	66
113102_22_2m	164.66	4.42	Pb/U w/mean conc.	0.9866	1.0000	1.0000	91
113102_23_2m	1637.62	41.14	Pb/Pb w/mean conc.	0.9861	1.0000	1.0000	52
113102_24_2m	137.61	3.87	Pb/U w/mean conc.	0.9702	1.0000	1.0000	55
113102_25_2m	1018.89	29.84	Pb/U w/mean conc.	0.8187	1.0000	1.0000	146
113102_26_2m	120.22	3.21	Pb/U w/mean conc.	0.6122	1.0000	1.0000	77
113102_27_2m	1207.40	112.05	upper int.	0.9999	1.0000	1.0000	0
113102_28_2m	147.80	3.97	Pb/U w/mean conc.	0.7967	1.0000	1.0000	57
113102_29_2m	undefined	undefined	undefined	0.9447	1.0000	1.0000	0
113102_30_2m	569.65	15.59	Pb/U w/mean conc.	0.9324	1.0000	1.0000	126
113102_31_2m	652.69	18.14	Pb/U w/mean conc.	0.9703	1.0000	1.0000	98
113102_32_2m	946.73	26.95	Pb/U w/mean conc.	0.8077	1.0000	1.0000	90
113102_33_2m	1469.66	41.50	Pb/Pb w/mean conc.	0.7780	1.0000	1.0000	100
113102_34_2m	513.99	14.19	Pb/U w/mean conc.	0.7925	1.0000	1.0000	83
113102_35_2m	1762.29	40.18	Pb/Pb w/mean conc.	0.8677	1.0000	1.0000	57
113102_36_2m	378.41	10.54	Pb/U w/mean conc.	0.7751	1.0000	1.0000	73
113102_37_2m	378.41	10.54	Pb/U w/mean conc.	0.7751	1.0000	1.0000	73
113102_38_2m	1217.17	35.29	Pb/U w/mean conc.	0.8533	1.0000	1.0000	102
113102_39_2m	684.46	19.07	Pb/U w/mean conc.	0.9995	1.0000	1.0000	114
113102_40_2m	158.44	4.23	Pb/U w/mean conc.	0.9745	1.0000	1.0000	56
113102_41_2m	136.69	3.73	Pb/U w/mean conc.	0.8187	1.0000	1.0000	52
113102_42_2m	409.84	15.19	Pb/U w/mean conc.	0.2893	1.0000	1.0000	96
113102_43_2m	undefined	undefined	undefined	0.9964	0.5000	1.0000	0
113102_44_2m	1134.52	32.73	Pb/U w/mean conc.	0.9999	1.0000	1.0000	102
113102_45_2m	406.88	11.12	Pb/U w/mean conc.	0.8300	1.0000	1.0000	69
113102_46_2m	2696.38	36.06	Pb/Pb w/mean conc.	0.9577	1.0000	1.0000	112
113102_47_2m	2252.00	594.52	upper int.	1.0000	1.0000	1.0000	0
113102_48_2m	136.69	3.73	Pb/U w/mean conc.	0.8187	1.0000	1.0000	52
113102_49_2m	149.69	4.08	Pb/U w/mean conc.	0.6358	1.0000	1.0000	82
113102_50_2m	591.65	16.36	Pb/U w/mean conc.	0.9878	1.0000	1.0000	91

Morrison/Cedar Mtn

Preferred Ages – Sorted

Analysis_#	Preferred Age (Ma) (±2σ)	Absolute Error (±2σ)	Age Type	Fit Quality 1=best	Clean Quality 1=best	204Pb Quality 1=best	2σ Conc. Scans
113102_29	undefined	undefined	undefined	0.9447	1.0000	1.0000	0
113102_18	143.70	3.87	Pb/U w/mean conc.	0.9702	1.0000	1.0000	55
113102_28	147.80	3.97	Pb/U w/mean conc.	0.7967	1.0000	1.0000	57
113102_49	149.69	4.00	Pb/U w/mean conc.	0.6358	1.0000	1.0000	82
113102_21	153.47	4.19	Pb/U w/mean conc.	1.0000	1.0000	0.8526	66
113102_11	154.46	4.16	Pb/U w/mean conc.	0.6767	0.5000	1.0000	54
113102_20	162.87	4.41	Pb/U w/mean conc.	0.8682	1.0000	1.0000	10
113102_40	158.44	4.23	Pb/U w/mean conc.	0.9745	1.0000	1.0000	56
113102_48	136.61	3.64	Pb/U w/mean conc.	0.7501	1.0000	1.0000	53
113102_37	386.71	10.54	Pb/U w/mean conc.	0.7761	1.0000	1.0000	73
113102_36	386.71	10.54	Pb/U w/mean conc.	0.7751	1.0000	1.0000	73
113102_42	409.66	11.19	Pb/U w/mean conc.	0.8957	1.0000	1.0000	96
113102_12	1130.42	32.55	Pb/U w/mean conc.	0.9639	1.0000	1.0000	97
113102_19	415.75	11.46	Pb/U w/mean conc.	0.7212	1.0000	1.0000	37
113102_6	458.23	12.64	Pb/U w/mean conc.	0.7505	1.0000	1.0000	55
113102_14	628.02	17.43	Pb/U w/mean conc.	0.9519	1.0000	1.0000	91
113102_3	420.99	11.55	Pb/U w/mean conc.	0.8407	1.0000	1.0000	110
113102_16	413.73	11.30	Pb/U w/mean conc.	0.3943	1.0000	1.0000	71
113102_34	513.99	14.19	Pb/U w/mean conc.	0.7925	1.0000	1.0000	83
113102_2	504.53	13.86	Pb/U w/mean conc.	0.6726	0.5000	1.0000	111
113102_17	563.59	15.53	Pb/U w/mean conc.	0.9964	1.0000	1.0000	106
113102_30	569.65	15.59	Pb/U w/mean conc.	0.9324	1.0000	1.0000	126
113102_50	591.65	16.36	Pb/U w/mean conc.	0.9878	1.0000	1.0000	91
113102_24	626.02	17.43	Pb/U w/mean conc.	0.9519	1.0000	1.0000	91
113102_45	406.88	11.12	Pb/U w/mean conc.	0.8300	1.0000	1.0000	69
113102_10	675.98	18.89	Pb/U w/mean conc.	0.6723	1.0000	1.0000	55
113102_39	684.46	19.07	Pb/U w/mean conc.	0.9995	1.0000	1.0000	114
113102_9	558.94	15.35	Pb/U w/mean conc.	0.6383	1.0000	1.0000	103
113102_35	1762.29	40.18	Pb/Pb w/mean conc.	0.8677	1.0000	1.0000	57
113102_44	1134.52	32.73	Pb/U w/mean conc.	0.9999	1.0000	1.0000	102
113102_23	1637.62	41.14	Pb/Pb w/mean conc.	0.9861	1.0000	1.0000	52
113102_13	1130.42	32.55	Pb/U w/mean conc.	0.9639	1.0000	1.0000	97
113102_1	1639.61	28.04	Pb/Pb w/mean conc.	0.0000	1.0000	1.0000	88
113102_4	1789.97	39.84	Pb/Pb w/mean conc.	0.9605	1.0000	1.0000	133
113102_5	1209.52	35.07	Pb/U w/mean conc.	0.8872	1.0000	1.0000	125
113102_7	1070.82	30.77	Pb/U w/mean conc.	0.9018	1.0000	1.0000	95
113102_8	1.00.00	1.000	Pb/U w/mean conc.	0.0000	1.0000	1.0000	1
113102_15	696.58	19.38	Pb/U w/mean conc.	0.9400	1.0000	1.0000	115
113102_22	164.66	4.42	Pb/U w/mean conc.	0.9866	1.0000	1.0000	91
113102_26	120.22	3.21	Pb/U w/mean conc.	0.6122	1.0000	1.0000	77
113102_27	1207.40	112.05	upper int.	0.9999	1.0000	1.0000	0
113102_32	946.73	26.95	Pb/U w/mean conc.	0.8077	1.0000	1.0000	90
113102_33	1469.66	41.50	Pb/Pb w/mean conc.	0.7780	1.0000	1.0000	100
113102_41	136.69	3.73	Pb/U w/mean conc.	0.8187	1.0000	1.0000	52
113102_47	2252.00	594.52	upper int.	1.0000	1.0000	1.0000	0
113102_46	2696.38	36.06	Pb/Pb w/mean conc.	0.9577	1.0000	1.0000	112
113102_36	378.41	10.54	Pb/U w/mean conc.	0.7751	1.0000	1.0000	73
113102_37	378.41	10.54	Pb/U w/mean conc.	0.7751	1.0000	1.0000	73
113102_38	1217.17	35.29	Pb/U w/mean conc.	0.8533	1.0000	1.0000	102
113102_39	684.46	19.07	Pb/U w/mean conc.	0.9995	1.0000	1.0000	114
113102_40	158.44	4.23	Pb/U w/mean conc.	0.9745	1.0000	1.0000	56
113102_41	136.69	3.73	Pb/U w/mean conc.	0.8187	1.0000	1.0000	52
113102_42	409.84	15.19	Pb/U w/mean conc.	0.2893	1.0000	1.0000	96
113102_43	undefined	undefined	undefined	0.9964	0.5000	1.0000	0
113102_44	1134.52	32.73	Pb/U w/mean conc.	0.9999	1.0000	1.0000	102
113102_45	406.88	11.12	Pb/U w/mean conc.	0.8300	1.0000	1.0000	69
113102_46	2696.38	36.06	Pb/Pb w/mean conc.	0.9577	1.0000	1.0000	112
113102_47	2252.00	594.52	upper int.	1.0000	1.0000	1.0000	0
113102_48	136.69	3.73	Pb/U w/mean conc.	0.8187	1.0000	1.0000	52
113102_49	149.69	4.08	Pb/U w/mean conc.	0.6358	1.0000	1.0000	82
113102_50	591.65	16.36	Pb/U w/mean conc.	0.9878	1.0000	1.0000	91

Morrison/Cedar Mtn

Fitted Discordia

Analysis_#	Age - Lower Intercept (Ma) (±2σ)	Age - Lower Intercept (±2σ)	Age - Upper Intercept (Ma) (±2σ)	Age - Upper Intercept (±2σ)	Regression Intercept (±2σ)	Regression Intercept (±1σ)	Regression Slope (±2σ)	Regression Slope (±1σ)
113102_29	undefined	undefined	undefined	undefined	undefined	undefined	undefined	undefined
113102_18	499.40	20.97	2302.67	664.14	0.004510	0.004425	0.041920	0.006993
113102_28	147.00	3.97	undefined	undefined	0.070420	0.002777	-0.006404	0.005492
113102_49	149.69	4.00	undefined	undefined	-0.013771	0.012152	0.006464	0.002538
113102_21	153.47	4.19	1705.50	425.01	0.105666	0.006455	0.045190	0.003721
113102_11	154.46	4.16	undefined	undefined	0.073582	0.003027	-0.000831	0.004640
113102_20	162.97	4.41	2927.73	1549.23	0.128269	0.012151	0.026465	0.006348
113102_40	158.44	4.23	4560.00	900.72	0.082655	0.003269	0.010601	0.003747
113102_48	136.61	5.04	3045.09	427.08	0.090332	0.002429	0.028035	0.003232
113102_37	386.71	10.54	935.44	0.097286	0.004397	0.012709	0.004519	-62.75%
113102_36	386.71	10.54	4560.00	574.03	0.022415	0.000630	0.009242	0.003326
113102_42	409.66	11.19	4560.00	325.03	0.118850	0.006483	0.034378	0.002607
113102_12	1130.42	32.55	undefined	undefined	0.000000	0.000000	0.129249	0.003978
113102_19	415.75	11.46	4560.00	900.72	0.092655	0.003269	0.010601	0.003747
113102_6	458.23	12.64	4560.00	544.08	0.080577	0.004015	0.028993	0.004038
113102_14	628.02	17.43	4560.00	900.72	0.092655	0.003269	0.010601	0.003747
113102_3	420.99	11.55	2927.49	544.08	0.080577	0.004015	0.028993	0.004038
113102_16	413.73	11.30	undefined	undefined	0.081221	0.005081	0.009389	0.005607
113102_2	504.53	13.86	2524.30	729.88	0.063804	0.004721	0.037792	0.006311
113102_34	513.99	14.19	2513.04	546.68	0.013956	0.001334	0.053323	0.007485
113102_30	569.65	15.59	undefined	undefined	0.022159	0.000131	0.009724	0.000191
113102_50	591.65	16.36	2506.19	491.05	0.018210	0.001057	0.042296	0.005889
113102_24	626.02	17.43	1984.50	199.67	0.109451	0.010476	0.041298	0.002689
113102_45	406.88	11.12	2927.49	544.08	0.080577	0.004015	0.028993	0.004038
113102_10	675.98	18.89	2927.49	324.61	0.090759	0.005256	0.046166	0.003514
113102_39	684.46	19.07	2927.49	562.29	0.014914	0.000619	0.030200	0.004721
113102_9	558.94	15.35	2506.19	1207.40	0.019533	0.000256	0.001829	0.000567
113102_32	946.73	26.95	1207.40	112.05	0.019533	0.000697	0.019828	0.004161
113102_33	1469.66	41.50	1207.40	112.05	0.019533	0.000697	0.019828	0.004161
113102_41	1038.97	29.73	1207.40	112.05	0.019533	0.000697	0.019828	0.004161
113102_47	1038.97	29.73	1207.40	112.05	0.019533	0.000697	0.019828	0.004161
113102_46	1038.97	29.73	1207.40	112.05	0.019533	0.000697	0.019828	0.004161
113102_7	1070.82	30.77	4560.00	900.38	0.087196	0.003357	0.015270	0.003676
113102_12	1130.42	32.55	4560.00	447.07	0.122051	0.004447	0.023684	0.002710
113102_44	1134.52	32.73

Preferred Age (Ma)	Absolute Error (±2σ)	Age Type	Analysis #	Product of Qualities
120.22	3.21	Pb/U wmean conc.	113102_26	0.6122
143.70	3.87	Pb/U wmean conc.	113102_18	0.9762
147.80	3.97	Pb/U wmean conc.	113102_28	0.7907
149.69	4.00	Pb/U wmean conc.	113102_49	0.6358
153.47	4.19	Pb/U wmean conc.	113102_21	0.8526
154.46	4.16	Pb/U wmean conc.	113102_11	0.3384
162.97	4.61	Pb/U wmean conc.	113102_20	0.8682
164.66	4.42	Pb/U wmean conc.	113102_22	0.9966
186.61	5.04	Pb/U wmean conc.	113102_48	0.7501
386.71	10.54	Pb/U wmean conc.	113102_37	0.7761
406.88	11.12	Pb/U wmean conc.	113102_45	0.8300
409.66	11.19	Pb/U wmean conc.	113102_42	0.8957
413.73	11.30	Pb/U wmean conc.	113102_16	0.9943
415.75	11.46	Pb/U wmean conc.	113102_19	0.7212
420.99	11.55	Pb/U wmean conc.	113102_3	0.8407
458.23	12.64	Pb/U wmean conc.	113102_6	0.7926
504.53	13.86	Pb/U wmean conc.	113102_2	0.3363
513.98	14.19	Pb/U wmean conc.	113102_34	0.7935
556.94	15.35	Pb/U wmean conc.	113102_9	0.6383
563.59	15.53	Pb/U wmean conc.	113102_17	0.9954
565.65	15.59	Pb/U wmean conc.	113102_30	0.9324
591.65	16.36	Pb/U wmean conc.	113102_50	0.9878
628.02	17.43	Pb/U wmean conc.	113102_14	0.9519
652.69	18.14	Pb/U wmean conc.	113102_31	0.9703
675.98	18.89	Pb/U wmean conc.	113102_10	0.6723
684.46	19.07	Pb/U wmean conc.	113102_39	0.9995
696.58	19.38	Pb/U wmean conc.	113102_15	0.9400
946.73	26.95	Pb/U wmean conc.	113102_32	0.8077
1018.89	29.04	Pb/U wmean conc.	113102_25	0.8167
1036.97	29.73	Pb/U wmean conc.	113102_41	0.7916
1070.82	30.77	Pb/U wmean conc.	113102_7	0.9218
1130.42	32.55	Pb/U wmean conc.	113102_12	0.9939
1134.52	32.73	Pb/U wmean conc.	113102_44	1.0000
1207.40	112.05	upper int.	113102_27	0.3926
1209.52	35.07	Pb/U wmean conc.	113102_5	0.9872
1217.17	35.29	Pb/U wmean conc.	113102_38	0.8533
1469.66	41.50	Pb/U wmean conc.	113102_33	0.7780
1614.08	40.71	Pb/Pb wmean conc.	113102_1	0.9834
1637.62	41.14	Pb/Pb wmean conc.	113102_23	0.9861
1737.51	39.94	Pb/Pb wmean conc.	113102_24	0.4644
1762.29	40.18	Pb/Pb wmean conc.	113102_35	0.8677
1789.97	39.84	Pb/Pb wmean conc.	113102_4	0.9605
2019.70	38.83	Pb/Pb wmean conc.	113102_36	0.9996
2252.00	594.52	upper int.	113102_47	1.0000
2576.34	270.50	upper int.	113102_29	0.9974
2684.20	36.00	Pb/Pb wmean conc.	113102_8	0.9962
2696.38	36.06	Pb/Pb wmean conc.	113102_46	0.9577
rej 156	_ _ _ _ _ M	Pb/U wmean conc.	113102_40	0.9745
undefined	undefined	undefined	113102_13	0.9447
undefined	undefined	undefined	113102_43	0.4992
0.00	0.00	0.00	0.00	0.0000
#REF!	#REF!	#REF!	#REF!	#REF!
#REF!	#REF!	#REF!	#REF!	#REF!
#REF!	#REF!	#REF!	#REF!	#REF!

378
379.5
381
382.5
384
385.5
387
388.5
390
391.5
393
394.5
396
397.5
399
400.5
402
403.5
405
406.5
408
409.5
411
412.5
414
415.5
417
418.5
420
421.5
423
424.5
426
427.5
429
430.5
432
433.5
435
436.5
438
439.5
441
442.5
444
445.5
447
448.5
450
451.5
453
454.5
456
457.5
459
460.5
462
463.5
465
466.5
468
469.5
471
472.5
474
475.5
477
478.5
480
481.5
483
484.5
486
487.5
489
490.5
492
493.5
495
496.5
498
499.5
501
502.5
504
505.5
507
508.5
510
511.5
513
514.5
516
517.5
519
520.5
522
523.5
525
526.5
528
529.5
531
532.5
534
535.5
537
538.5
540
541.5
543
544.5
546
547.5
549
550.5
552
553.5
555
556.5
558
559.5
561
562.5
564
565.5
567
568.5
570
571.5
573
574.5
576
577.5
579
580.5
582
583.5
585
586.5
588
589.5
591
592.5
594
595.5
597
598.5
600
601.5
603
604.5
606
607.5
609
610.5
612
613.5
615
616.5
618
619.5
621
622.5
624
625.5
627
628.5
630
631.5
633
634.5
636
637.5
639
640.5
642
643.5
645
646.5
648
649.5
651
652.5
654
655.5
657
658.5
660
661.5
663
664.5
666
667.5
669
670.5
672
673.5
675
676.5
678
679.5
681
682.5
684
685.5
687
688.5
690
691.5
693
694.5
696
697.5
699
700.5
702
703.5
705
706.5
708
709.5
711
712.5
714
715.5
717
718.5
720
721.5
723
724.5
726
727.5
729
730.5
732
733.5
735
736.5
738
739.5
741
742.5
744
745.5
747
748.5
750
751.5
753
754.5
756

1516.5
1518
1519.5
1521
1522.5
1524
1525.5
1527
1528.5
1530
1531.5
1533
1534.5
1536
1537.5
1539
1540.5
1542
1543.5
1545
1546.5
1548
1549.5
1551
1552.5
1554
1555.5
1557
1558.5
1560
1561.5
1563
1564.5
1566
1567.5
1569
1570.5
1572
1573.5
1575
1576.5
1578
1579.5
1581
1582.5
1584
1585.5
1587
1588.5
1590
1591.5
1593
1594.5
1596
1597.5
1599
1600.5
1602
1603.5
1605
1606.5
1608
1609.5
1611
1612.5
1614
1615.5
1617
1618.5
1620
1621.5
1623
1624.5
1626
1627.5
1629
1630.5
1632
1633.5
1635
1636.5
1638
1639.5
1641
1642.5
1644
1645.5
1647
1648.5
1650
1651.5
1653
1654.5
1656
1657.5
1659
1660.5
1662
1663.5
1665
1666.5
1668
1669.5
1671
1672.5
1674
1675.5
1677
1678.5
1680
1681.5
1683
1684.5
1686
1687.5
1689
1690.5
1692
1693.5
1695
1696.5
1698
1699.5
1701
1702.5
1704
1705.5
1707
1708.5
1710
1711.5
1713
1714.5
1716
1717.5
1719
1720.5
1722
1723.5
1725
1726.5
1728
1729.5
1731
1732.5
1734
1735.5
1737
1738.5
1740
1741.5
1743
1744.5
1746
1747.5
1749
1750.5
1752
1753.5
1755
1756.5
1758
1759.5
1761
1762.5
1764
1765.5
1767
1768.5
1770
1771.5
1773
1774.5
1776
1777.5
1779
1780.5
1782
1783.5
1785
1786.5
1788
1789.5
1791
1792.5
1794
1795.5
1797
1798.5
1800
1801.5
1803
1804.5
1806
1807.5
1809
1810.5
1812
1813.5
1815
1816.5
1818
1819.5
1821
1822.5
1824
1825.5
1827
1828.5
1830
1831.5
1833
1834.5
1836
1837.5
1839
1840.5
1842
1843.5
1845
1846.5
1848
1849.5
1851
1852.5
1854
1855.5
1857
1858.5
1860
1861.5
1863
1864.5
1866
1867.5
1869
1870.5
1872
1873.5
1875
1876.5
1878
1879.5
1881
1882.5
1884
1885.5
1887
1888.5
1890
1891.5
1893
1894.5

1896
1897.5
1902
1906.5
1908
1909.5
1910
1910.5
1911
1913.5
1914
1915.5
1916
1917
1918.5
1919
1920.5
1922
1923.5
1926
1927.5
1929
1931.5
1934
1935.5
1936
1937.5
1938
1939.5
1940
1941.5
1942
1943.5
1944
1945.5
1946
1947.5
1948
1949.5
1950
1951.5
1954
1955.5
1956
1957.5
1958
1959.5
1960
1961.5
1962
1963.5
1964
1965.5
1966
1967.5
1968
1969.5
1970
1971.5
1972
1973.5
1974
1975.5
1976
1977.5
1978
1979.5
1980
1981.5
1982
1983.5
1984
1985.5
1986
1987.5
1988
1989.5
1990
1991.5
1992
1993.5
1994
1995.5
1996
1997.5
1998
1999.5
2000
2001.5
2002
2003.5
2004
2005.5
2006
2007.5
2008
2009.5
2010
2011.5
2012
2013.5
2014
2015.5
2016
2017.5
2018
2019.5
2020
2021.5
2022
2023.5
2024
2025.5
2026
2027.5
2028
2029.5
2030
2031.5
2032
2033.5
2034
2035.5
2036
2037.5
2038
2039.5
2040
2041.5
2042
2043.5
2044
2045.5
2046
2047.5
2048
2049.5
2050
2051.5
2052
2053.5
2054
2055.5
2056
2057.5
2058
2059.5
2060
2061.5
2062
2063.5
2064
2065.5
2066
2067.5
2068
2069.5
2070
2071.5
2072
2073.5
2074
2075.5
2076
2077.5
2078
2079.5
2080
2081.5
2082
2083.5
2084
2085.5
2086
2087.5
2088
2089.5
2090
2091.5
2092
2093.5
2094
2095.5
2096
2097.5
2098
2099.5
2100
2101.5
2102
2103.5
2104
2105.5
2106
2107.5
2108
2109.5
2110
2111.5
2112
2113.5
2114
2115.5
2116
2117.5
2118
2119.5
2120
2121.5
2122
2123.5
2124
2125.5
2126
2127.5
2128
2129.5
2130
2131.5
2132
2133.5
2134
2135.5
2136
2137.5
2138
2139.5
2140
2141.5
2142
2143.5
2144
2145.5
2146
2147.5
2148
2149.5
2150
2151.5
2152
2153.5
2154
2155.5
2156
2157.5
2158
2159.5
2160
2161.5
2162
2163.5
2164
2165.5
2166
2167.5
2168
2169.5
2170
2171.5
2172
2173.5
2174
2175.5
2176
2177.5
2178
2179.5
2180
2181.5
2182
2183.5
2184
2185.5
2186
2187.5
2188
2189.5
2190
2191.5
2192
2193.5
2194
2195.5
2196
2197.5
2198
2199.5
2200
2201.5
2202
2203.5
2204
2205.5
2206
2207.5
2208
2209.5
2210
2211.5
2212
2213.5
2214
2215.5
2216
2217.5
2218
2219.5
2220
2221.5
2222
2223.5
2224
2225.5
2226
2227.5
2228
2229.5
2230
2231.5
2232
2233.5
2234
2235.5
2236
2237.5
2238
2239.5
2240
2241.5
2242
2243.5
2244
2245.5
2246
2247.5
2248
2249.5
2250
2251.5
2252
2253.5
2254
2255.5
2256
2257.5
2258
2259.5
2260
2261.5
2262
2263.5
2264
2265.5
2266
2267.5
2268
2269.5
2270
2271.5
2272
2273.5
2274

2275.5
2277
2278.5
2279.5
2280
2281.5
2282
2283.5
2284.5
2286
2287.5
2289
2290.5
2292
2293.5
2295
2296.5
2298
2299.5
2301
2302.5
2304
2305.5
2307
2308.5
2310
2311.5
2313
2314.5
2316
2317.5
2319
2320.5
2322
2323.5
2325
2326.5
2328
2329.5
2331
2332.5
2334
2335.5
2337
2338.5
2340
2341.5
2343
2344.5
2346
2347.5
2349
2350.5
2352
2353.5
2355
2356.5
2358
2359.5
2361
2362.5
2364
2365.5
2367
2368.5
2370
2371.5
2373
2374.5
2376
2377.5
2379
2380.5
2382
2383.5
2385
2386.5
2388
2389.5
2391
2392.5
2394
2395.5
2397
2398.5
2400
2401.5
2403
2404.5
2406
2407.5
2409
2410.5
2412
2413.5
2415
2416.5
2418
2419.5
2421
2422.5
2424
2425.5
2427
2428.5
2430
2431.5
2433
2434.5
2436
2437.5
2439
2440.5
2442
2443.5
2445
2446.5
2448
2449.5
2451
2452.5
2454
2455.5
2457
2458.5
2460
2461.5
2463
2464.5
2466
2467.5
2469
2470.5
2472
2473.5
2475
2476.5
2478
2479.5
2481
2482.5
2484
2485.5
2487
2488.5
2490
2491.5
2493
2494.5
2496
2497.5
2499
2500.5
2502
2503.5
2505
2506.5
2508
2509.5
2511
2512.5
2514
2515.5
2517
2518.5
2520
2521.5
2523
2524.5
2526
2527.5
2529
2530.5
2532
2533.5
2535
2536.5
2538
2539.5
2541
2542.5
2544
2545.5
2547
2548.5
2550
2551.5
2553
2554.5
2556
2557.5
2559
2560.5
2562
2563.5
2565
2566.5
2568
2569.5
2571
2572.5
2574
2575.5
2577
2578.5
2580
2581.5
2583
2584.5
2586
2587.5
2589
2590.5
2592
2593.5
2595
2596.5
2598
2599.5
2601
2602.5
2604
2605.5
2607
2608.5
2610
2611.5
2613
2614.5
2616
2617.5
2619
2620.5
2622
2623.5
2625
2626.5
2628
2629.5
2631
2632.5
2634
2635.5
2637
2638.5
2640
2641.5
2643
2644.5
2646
2647.5
2649
2650.5
2652
2653.5

2655
2659.5
2662
2661
2664
2665.5
2666.5
2668.5
2671
2673
2675.5
2676
2677.5
2679
2680.5
2682
2683.5
2685
2686.5
2688.5
2690.5
2692.5
2694
2695.5
2697
2699
2701.5
2703
2705.5
2706
2707.5
2709
2710.5
2713.5
2715
2716.5
2718
2719.5
2721
2722.5
2723.5
2724
2725.5
2727
2728.5
2729.5
2731.5
2733
2734.5
2736
2737.5
2739
2740.5
2742.5
2744.5
2745.5
2747.5
2748
2749.5
2751
2752.5
2754
2755.5
2757
2758.5
2760
2761.5
2763
2764.5
2766
2767.5
2769
2770.5
2772
2773.5
2775
2776.5
2778
2779.5
2781
2782.5
2784
2785.5
2787
2788.5
2790
2791.5
2793
2794.5
2796
2797.5
2799
2800.5
2802
2803.5
2805
2806.5
2808
2809.5
2811
2812.5
2814
2815.5
2817
2818.5
2820
2821.5
2823
2824.5
2826
2827.5
2829
2830.5
2832
2833.5
2835
2836.5
2838
2839.5
2841
2842.5
2844
2845.5
2847
2848.5
2850
2851.5
2853
2854.5
2856
2857.5
2859
2860.5
2862
2863.5
2865
2866.5
2868
2869.5
2871
2872.5
2874
2875.5
2877
2878.5
2880
2881.5
2883
2884.5
2886
2887.5
2889
2890.5
2892
2893.5
2895
2896.5
2898
2899.5
2901
2902.5
2904
2905.5
2907
2908.5
2910
2911.5
2913
2914.5
2916
2917.5
2919
2920.5
2922
2923.5
2925
2926.5
2928
2929.5
2931
2932.5
2934
2935.5
2937
2938.5
2940
2941.5
2943
2944.5
2946
2947.5
2949
2950.5
2952
2953.5
2955
2956.5
2958
2959.5
2961
2962.5
2964
2965.5
2967
2968.5
2970
2971.5
2973
2974.5
2976
2977.5
2979
2980.5
2982
2983.5
2985
2986.5
2988
2989.5
2991
2992.5
2994
2995.5
2997
2998.5
3000
3001.5
3003
3004.5
3006
3007.5
3009
3010.5
3012
3013.5
3015
3016.5
3018
3019.5
3021
3022.5
3024
3025.5
3027
3028.5
3030
3031.5
3033

3034.5
3036
3037.5
3039
3040.5
3042
3043.5
3045
3046.5
3048
3049.5
3051
3052.5
3054
3055.5
3057
3058.5
3060
3061.5
3063
3064.5
3066
3067.5
3069
3070.5
3072
3073.5
3075
3076.5
3078
3079.5
3081
3082.5
3084
3085.5
3087
3088.5
3090
3091.5
3093
3094.5
3096
3097.5
3099
3100.5
3102
3103.5
3105
3106.5
3108
3109.5
3111
3112.5
3114
3115.5
3117
3118.5
3120
3121.5
3123
3124.5
3126
3127.5
3129
3130.5
3132
3133.5
3135
3136.5
3138
3139.5
3141
3142.5
3144
3145.5
3147
3148.5
3150
3151.5
3153
3154.5
3156
3157.5
3159
3160.5
3162
3163.5
3165
3166.5
3168
3169.5
3171
3172.5
3174
3175.5
3177
3178.5
3180
3181.5
3183
3184.5
3186
3187.5
3189
3190.5
3192
3193.5
3195
3196.5
3198
3199.5
3201
3202.5
3204
3205.5
3207
3208.5
3210
3211.5
3213
3214.5
3216
3217.5
3219
3220.5
3222
3223.5
3225
3226.5
3228
3229.5
3231
3232.5
3234
3235.5
3237
3238.5
3240
3241.5
3243
3244.5
3246
3247.5
3249
3250.5
3252
3253.5
3255
3256.5
3258
3259.5
3261
3262.5
3264
3265.5
3267
3268.5
3270
3271.5
3273
3274.5
3276
3277.5
3279
3280.5
3282
3283.5
3285
3286.5
3288
3289.5
3291
3292.5
3294
3295.5
3297
3298.5
3300
3301.5
3303
3304.5
3306
3307.5
3309
3310.5
3312
3313.5
3315
3316.5
3318
3319.5
3321
3322.5
3324
3325.5
3327
3328.5
3330
3331.5
3333
3334.5
3336
3337.5
3339
3340.5
3342
3343.5
3345
3346.5
3348
3349.5
3351
3352.5
3354
3355.5
3357
3358.5
3360
3361.5
3363
3364.5
3366
3367.5
3369
3370.5
3372
3373.5
3375
3376.5
3378
3379.5
3381
3382.5
3384
3385.5
3387
3388.5
3390
3391.5
3393
3394.5
3396
3397.5
3399
3400.5
3402
3403.5
3405
3406.5
3408
3409.5
3411
3412.5

3414
3415
3416
3417
3418
3419
3420
3421
3422
3423
3424
3425
3426
3427
3428
3429
3430
3431
3432
3433
3434
3435
3436
3437
3438
3439
3440
3441
3442
3443
3444
3445
3446
3447
3448
3449
3450
3451
3452
3453
3454
3455
3456
3457
3458
3459
3460
3461
3462
3463
3464
3465
3466
3467
3468
3469
3470
3471
3472
3473
3474
3475
3476
3477
3478
3479
3480
3481
3482
3483
3484
3485
3486
3487
3488
3489
3490
3491
3492
3493
3494
3495
3496
3497
3498
3499
3500
3501
3502
3503
3504
3505
3506
3507
3508
3509
3510
3511
3512
3513
3514
3515
3516
3517
3518
3519
3520
3521
3522
3523
3524
3525
3526
3527
3528
3529
3530
3531
3532
3533
3534
3535
3536
3537
3538
3539
3540
3541
3542
3543
3544
3545
3546
3547
3548
3549
3550
3551
3552
3553
3554
3555
3556
3557
3558
3559
3560
3561
3562
3563
3564
3565
3566
3567
3568
3569
3570
3571
3572
3573
3574
3575
3576
3577
3578
3579
3580
3581
3582
3583
3584
3585
3586
3587
3588
3589
3590
3591
3592
3593
3594
3595
3596
3597
3598
3599
3600
3601
3602
3603
3604
3605
3606
3607
3608
3609
3610
3611
3612
3613
3614
3615
3616
3617
3618
3619
3620
3621
3622
3623
3624
3625
3626
3627
3628
3629
3630
3631
3632
3633
3634
3635
3636
3637
3638
3639
3640
3641
3642
3643
3644
3645
3646
3647
3648
3649
3650
3651
3652
3653
3654
3655
3656
3657
3658
3659
3660
3661
3662
3663
3664
3665
3666
3667
3668
3669
3670
3671
3672
3673
3674
3675
3676
3677
3678
3679
3680
3681
3682
3683
3684
3685
3686
3687
3688
3689
3690
3691
3692
3693
3694
3695
3696
3697
3698
3699
3700
3701
3702
3703
3704
3705
3706
3707
3708
3709
3710
3711
3712
3713
3714
3715
3716
3717
3718
3719
3720
3721
3722
3723
3724
3725
3726
3727
3728
3729
3730
3731
3732
3733
3734
3735
3736
3737
3738
3739
3740
3741
3742
3743
3744
3745
3746
3747
3748
3749
3750
3751
3752
3753
3754
3755
3756
3757
3758
3759
3760
3761
3762
3763
3764
3765

1264.5
1265
1265.5
1266
1266.5
1267
1267.5
1268
1268.5
1269
1269.5
1270
1270.5
1271
1271.5
1272
1272.5
1273
1273.5
1274
1274.5
1275
1275.5
1276
1276.5
1277
1277.5
1278
1278.5
1279
1279.5
1280
1280.5
1281
1281.5
1282
1282.5
1283
1283.5
1284
1284.5
1285
1285.5
1286
1286.5
1287
1287.5
1288
1288.5
1289
1289.5
1290
1290.5
1291
1291.5
1292
1292.5
1293
1293.5
1294
1294.5
1295
1295.5
1296
1296.5
1297
1297.5
1298
1298.5
1299
1299.5
1300
1300.5
1301
1301.5
1302
1302.5
1303
1303.5
1304
1304.5
1305
1305.5
1306
1306.5
1307
1307.5
1308
1308.5
1309
1309.5
1310
1310.5
1311
1311.5
1312
1312.5
1313
1313.5
1314
1314.5
1315
1315.5
1316
1316.5
1317
1317.5
1318
1318.5
1319
1319.5
1320
1320.5
1321
1321.5
1322
1322.5
1323
1323.5
1324
1324.5
1325
1325.5
1326
1326.5
1327
1327.5
1328
1328.5
1329
1329.5
1330
1330.5
1331
1331.5
1332
1332.5
1333
1333.5
1334
1334.5
1335
1335.5
1336
1336.5
1337
1337.5
1338
1338.5
1339
1339.5
1340
1340.5
1341
1341.5
1342
1342.5
1343
1343.5
1344
1344.5
1345
1345.5
1346
1346.5
1347
1347.5
1348
1348.5
1349
1349.5
1350
1350.5
1351
1351.5
1352
1352.5
1353
1353.5
1354
1354.5
1355
1355.5
1356
1356.5
1357
1357.5
1358
1358.5
1359
1359.5
1360
1360.5
1361
1361.5
1362
1362.5
1363
1363.5
1364
1364.5
1365
1365.5
1366
1366.5
1367
1367.5
1368
1368.5
1369
1369.5
1370
1370.5
1371
1371.5
1372
1372.5
1373
1373.5
1374
1374.5
1375
1375.5
1376
1376.5
1377
1377.5
1378
1378.5
1379
1379.5
1380
1380.5
1381
1381.5
1382
1382.5
1383
1383.5
1384
1384.5
1385
1385.5
1386
1386.5
1387
1387.5
1388
1388.5
1389
1389.5
1390
1390.5

1391
1391.5
1392
1392.5
1393
1393.5
1394
1394.5
1395
1395.5
1396
1396.5
1397
1397.5
1398
1398.5
1399
1399.5
1400
1400.5
1401
1401.5
1402
1402.5
1403
1403.5
1404
1404.5
1405
1405.5
1406
1406.5
1407
1407.5
1408
1408.5
1409
1409.5
1410
1410.5
1411
1411.5
1412
1412.5
1413
1413.5
1414
1414.5
1415
1415.5
1416
1416.5
1417
1417.5
1418
1418.5
1419
1419.5
1420
1420.5
1421
1421.5
1422
1422.5
1423
1423.5
1424
1424.5
1425
1425.5
1426
1426.5
1427
1427.5
1428
1428.5
1429
1429.5
1430
1430.5
1431
1431.5
1432
1432.5
1433
1433.5
1434
1434.5
1435
1435.5
1436
1436.5
1437
1437.5
1438
1438.5
1439
1439.5
1440
1440.5
1441
1441.5
1442
1442.5
1443
1443.5
1444
1444.5
1445
1445.5
1446
1446.5
1447
1447.5
1448
1448.5
1449
1449.5
1450
1450.5
1451
1451.5
1452
1452.5
1453
1453.5
1454
1454.5
1455
1455.5
1456
1456.5
1457
1457.5
1458
1458.5
1459
1459.5
1460
1460.5
1461
1461.5
1462
1462.5
1463
1463.5
1464
1464.5
1465
1465.5
1466
1466.5
1467
1467.5
1468
1468.5
1469
1469.5
1470
1470.5
1471
1471.5
1472
1472.5
1473
1473.5
1474
1474.5
1475
1475.5
1476
1476.5
1477
1477.5
1478
1478.5
1479
1479.5
1480
1480.5
1481
1481.5
1482
1482.5
1483
1483.5
1484
1484.5
1485
1485.5
1486
1486.5
1487
1487.5
1488
1488.5
1489
1489.5
1490
1490.5
1491
1491.5
1492
1492.5
1493
1493.5
1494
1494.5
1495
1495.5
1496
1496.5
1497
1497.5
1498
1498.5
1499
1499.5
1500
1500.5
1501
1501.5
1502
1502.5
1503
1503.5
1504
1504.5
1505
1505.5
1506
1506.5
1507
1507.5
1508
1508.5
1509
1509.5
1510
1510.5
1511
1511.5
1512
1512.5
1513
1513.5
1514
1514.5
1515
1515.5
1516
1516.5
1517

1517.5
1518
1518.5
1519
1519.5
1520
1520.5
1521
1521.5
1522
1522.5
1523
1523.5
1524
1524.5
1525
1525.5
1526
1526.5
1527
1527.5
1528
1528.5
1529
1529.5
1530
1530.5
1531
1531.5
1532
1532.5
1533
1533.5
1534
1534.5
1535
1535.5
1536
1536.5
1537
1537.5
1538
1538.5
1539
1539.5
1540
1540.5
1541
1541.5
1542
1542.5
1543
1543.5
1544
1544.5
1545
1545.5
1546
1546.5
1547
1547.5
1548
1548.5
1549
1549.5
1550
1550.5
1551
1551.5
1552
1552.5
1553
1553.5
1554
1554.5
1555
1555.5
1556
1556.5
1557
1557.5
1558
1558.5
1559
1559.5
1560
1560.5
1561
1561.5
1562
1562.5
1563
1563.5
1564
1564.5
1565
1565.5
1566
1566.5
1567
1567.5
1568
1568.5
1569
1569.5
1570
1570.5
1571
1571.5
1572
1572.5
1573
1573.5
1574
1574.5
1575
1575.5
1576
1576.5
1577
1577.5
1578
1578.5
1579
1579.5
1580
1580.5
1581
1581.5
1582
1582.5
1583
1583.5
1584
1584.5
1585
1585.5
1586
1586.5
1587
1587.5
1588
1588.5
1589
1589.5
1590
1590.5
1591
1591.5
1592
1592.5
1593
1593.5
1594
1594.5
1595
1595.5
1596
1596.5
1597
1597.5
1598
1598.5
1599
1599.5
1600
1600.5
1601
1601.5
1602
1602.5
1603
1603.5
1604
1604.5
1605
1605.5
1606
1606.5
1607
1607.5
1608
1608.5
1609
1609.5
1610
1610.5
1611
1611.5
1612
1612.5
1613
1613.5
1614
1614.5
1615
1615.5
1616
1616.5
1617
1617.5
1618
1618.5
1619
1619.5
1620
1620.5
1621
1621.5
1622
1622.5
1623
1623.5
1624
1624.5
1625
1625.5
1626
1626.5
1627
1627.5
1628
1628.5
1629
1629.5
1630
1630.5
1631
1631.5
1632
1632.5
1633
1633.5
1634
1634.5
1635
1635.5
1636
1636.5
1637
1637.5
1638
1638.5
1639
1639.5
1640
1640.5
1641
1641.5
1642
1642.5
1643
1643.5

1644
1644.5
1645
1645.5
1646
1646.5
1647
1647.5
1648
1648.5
1649
1649.5
1650
1650.5
1651
1651.5
1652
1652.5
1653
1653.5
1654
1654.5
1655
1655.5
1656
1656.5
1657
1657.5
1658
1658.5
1659
1659.5
1660
1660.5
1661
1661.5
1662
1662.5
1663
1663.5
1664
1664.5
1665
1665.5
1666
1666.5
1667
1667.5
1668
1668.5
1669
1669.5
1670
1670.5
1671
1671.5
1672
1672.5
1673
1673.5
1674
1674.5
1675
1675.5
1676
1676.5
1677
1677.5
1678
1678.5
1679
1679.5
1680
1680.5
1681
1681.5
1682
1682.5
1683
1683.5
1684
1684.5
1685
1685.5
1686
1686.5
1687
1687.5
1688
1688.5
1689
1689.5
1690
1690.5
1691
1691.5
1692
1692.5
1693
1693.5
1694
1694.5
1695
1695.5
1696
1696.5
1697
1697.5
1698
1698.5
1699
1699.5
1700
1700.5
1701
1701.5
1702
1702.5
1703
1703.5
1704
1704.5
1705
1705.5
1706
1706.5
1707
1707.5
1708
1708.5
1709
1709.5
1710
1710.5
1711
1711.5
1712
1712.5
1713
1713.5
1714
1714.5
1715
1715.5
1716
1716.5
1717
1717.5
1718
1718.5
1719
1719.5
1720
1720.5
1721
1721.5
1722
1722.5
1723
1723.5
1724
1724.5
1725
1725.5
1726
1726.5
1727
1727.5
1728
1728.5
1729
1729.5
1730
1730.5
1731
1731.5
1732
1732.5
1733
1733.5
1734
1734.5
1735
1735.5
1736
1736.5
1737
1737.5
1738
1738.5
1739
1739.5
1740
1740.5
1741
1741.5
1742
1742.5
1743
1743.5
1744
1744.5
1745
1745.5
1746
1746.5
1747
1747.5
1748
1748.5
1749
1749.5
1750
1750.5
1751
1751.5
1752
1752.5
1753
1753.5
1754
1754.5
1755
1755.5
1756
1756.5
1757
1757.5
1758
1758.5
1759
1759.5
1760
1760.5
1761
1761.5
1762
1762.5
1763
1763.5
1764
1764.5
1765
1765.5
1766
1766.5
1767
1767.5
1768
1768.5
1769
1769.5
1770
1770.5

1770.5
1771
1771.5
1772
1772.5
1773
1773.5
1774
1774.5
1775
1775.5
1776
1776.5
1777
1777.5
1778
1778.5
1779
1779.5
1780
1780.5
1781
1781.5
1782
1782.5
1783
1783.5
1784
1784.5
1785
1785.5
1786
1786.5
1787
1787.5
1788
1788.5
1789
1789.5
1790
1790.5
1791
1791.5
1792
1792.5
1793
1793.5
1794
1794.5
1795
1795.5
1796
1796.5
1797
1797.5
1798
1798.5
1799
1799.5
1800
1800.5
1801
1801.5
1802
1802.5
1803
1803.5
1804
1804.5
1805
1805.5
1806
1806.5
1807
1807.5
1808
1808.5
1809
1809.5
1810
1810.5
1811
1811.5
1812
1812.5
1813
1813.5
1814
1814.5
1815
1815.5
1816
1816.5
1817
1817.5
1818
1818.5
1819
1819.5
1820
1820.5
1821
1821.5
1822
1822.5
1823
1823.5
1824
1824.5
1825
1825.5
1826
1826.5
1827
1827.5
1828
1828.5
1829
1829.5
1830
1830.5
1831
1831.5
1832
1832.5
1833
1833.5
1834
1834.5
1835
1835.5
1836
1836.5
1837
1837.5
1838
1838.5
1839
1839.5
1840
1840.5
1841
1841.5
1842
1842.5
1843
1843.5
1844
1844.5
1845
1845.5
1846
1846.5
1847
1847.5
1848
1848.5
1849
1849.5
1850
1850.5
1851
1851.5
1852
1852.5
1853
1853.5
1854
1854.5
1855
1855.5
1856
1856.5
1857
1857.5
1858
1858.5
1859
1859.5
1860
1860.5
1861
1861.5
1862
1862.5
1863
1863.5
1864
1864.5
1865
1865.5
1866
1866.5
1867
1867.5
1868
1868.5
1869
1869.5
1870
1870.5
1871
1871.5
1872
1872.5
1873
1873.5
1874
1874.5
1875
1875.5
1876
1876.5
1877
1877.5
1878
1878.5
1879
1879.5
1880
1880.5
1881
1881.5
1882
1882.5
1883
1883.5
1884
1884.5
1885
1885.5
1886
1886.5
1887
1887.5
1888
1888.5
1889
1889.5
1890
1890.5
1891
1891.5
1892
1892.5
1893
1893.5
1894
1894.5
1895
1895.5
1896
1896.5


Morrison/Cedar Mtn
Relative Age Frequency
U-Pb Age (Ma)


Morrison/Cedar Mtn
Relative Age Frequency (red curve)
U/Th (blue points)
U-Pb Age (Ma)


Morrison/Cedar Mtn
Relative Age Frequency
U-Pb Age (Ma)


Morrison/Cedar Mtn
U/Th
U-Pb Age (Ma)


Morrison/Cedar Mtn
Frequency
U-Pb Age (Ma)


Morrison/Cedar Mtn
U ppm
U-Pb Age (Ma)


Morrison/Cedar Mtn
Lower-Age Intercept (Ma)
Upper-Age Intercept (Ma)


Morrison/Cedar Mtn
Th ppm
U-Pb Age (Ma)

Primary Standard

UPbICP Standard Index	6	Standard	F
Accepted Age	1099		
Isotopic Ratio	**Mean Age**	**FF Relative Error**	
207Pb/235U	1094.49	0.0087	
207Pb/235Uc	1098.24	0.0087	
206Pb/238U	1097.90	0.0131	
207Pb/206Pb	1099.00	0.0109	
208Pb/232Th	1096.60	0.0182	

Secondary Standard

UPbICP Standard Index	2	Standard	T
Accepted Age	61.23		
Uncorrected Age	59.94		

28Si 91Zr All Standards

Mineral	Zrn		
Statistics	**91Zr/28Si**	**28Si**	**91Zr**
median	0.759	2415364	1834113
standard deviation	0.114	238144	206892
minimum	not applicable	1972684	1558450
maximum	not applicable	2879354	2343284

Session	Grain	Preferred (Measured) Age (Ma)	Absolute Error (±2σ)	Age Type	Accepted Age (Ma)	Absolute Error (±2σ)	(Accepted - Preferred) / Accepted	Average (perfect=0)	Accepted / Measured	Average (perfect=1)
z100110	TR_1	60.90	1.62	Pb/U wmean conc.	61.23	0.11	0.0054	-0.0037	1.0054	0.9972
z100110	TR_2	61.92	1.66	Pb/U wmean conc.	61.23	0.11	-0.0112		0.9889	
z100110	TR_3	61.38	1.65	Pb/U wmean conc.	61.23	0.11	-0.0024		0.9976	
z100110	TR_5	58.69	1.58	Pb/U wmean conc.	61.23	0.11	0.0415		1.0433	
z100110	TR_6	60.84	1.64	Pb/U wmean conc.	61.23	0.11	0.0064		1.0065	
z100110	TR_7	60.45	1.64	Pb/U wmean conc.	61.23	0.11	0.0128		1.0130	
z100110	TR_8	62.47	1.71	Pb/U wmean conc.	61.23	0.11	-0.0203		0.9801	
z100110	TR_9	62.96	1.70	Pb/U wmean conc.	61.23	0.11	-0.0282		0.9726	
z100110	TR_11	64.25	1.83	Pb/U wmean conc.	61.23	0.11	-0.0493		0.9531	
z100110	TR_12	61.35	1.67	Pb/U wmean conc.	61.23	0.11	-0.0020		0.9980	
z100110	TR_13	61.62	1.68	Pb/U wmean conc.	61.23	0.11	-0.0064		0.9937	
z100110	TR_14	60.45	1.63	Pb/U wmean conc.	61.23	0.11	0.0127		1.0128	
z100110	FC5_1	1096.93	31.49	Pb/U wmean conc.	1099.00	0.60	0.0019		1.0019	
z100110	FC5_10	1095.36	31.62	Pb/U wmean conc.	1099.00	0.60	0.0033		1.0033	
z100110	FC5_2	1112.47	31.94	Pb/U wmean conc.	1099.00	0.60	-0.0123		0.9879	
z100110	FC5_3	1095.72	31.46	Pb/U wmean conc.	1099.00	0.60	0.0030		1.0030	
z100110	FC5_4	1100.93	31.65	Pb/U wmean conc.	1099.00	0.60	-0.0018		0.9982	
z100110	FC5_5	1120.66	32.23	Pb/U wmean conc.	1099.00	0.60	-0.0197		0.9807	
z100110	FC5_6	1112.76	31.98	Pb/U wmean conc.	1099.00	0.60	-0.0125		0.9876	
z100110	FC5_7	1098.72	31.56	Pb/U wmean conc.	1099.00	0.60	0.0003		1.0003	
z100110	FC5_8	1110.61	31.91	Pb/U wmean conc.	1099.00	0.60	-0.0106		0.9896	
z100110	FC5_9	1102.62	31.66	Pb/U wmean conc.	1099.00	0.60	-0.0033		0.9967	
z100110	IFCan_1	28.62	0.77	Pb/U wmean conc.	28.20	0.01	-0.0149		0.9854	
z100110	IFCan_10	28.49	0.76	Pb/U wmean conc.	28.20	0.01	-0.0104		0.9897	
z100110	IFCan_2	26.52	0.76	Pb/U wmean conc.	28.20	0.01	0.0595		1.0633	
z100110	IFCan_3	31.84	0.86	Pb/U wmean conc.	28.20	0.01	-0.1289		0.8858	
z100110	IFCan_4	29.34	0.79	Pb/U wmean conc.	28.20	0.01	-0.0402		0.9613	
z100110	IFCan_5	29.97	0.82	Pb/U wmean conc.	28.20	0.01	-0.0627		0.9410	
z100110	IFCan_6	26.65	0.86	Pb/U wmean conc.	28.20	0.01	0.0549		1.0581	
z100110	IFCan_7	27.64	0.73	Pb/U wmean conc.	28.20	0.01	0.0199		1.0203	
z100110	IFCan_8	27.26	0.73	Pb/U wmean conc.	28.20	0.01	0.0334		1.0346	
z100110	IFCan_9	29.37	0.96	Pb/U wmean conc.	28.20	0.01	-0.0416		0.9601	
z100110	MD_10	98.20	2.68	Pb/U wmean conc.	99.12	0.14	0.0092		1.0093	
z100110	MD_2	99.40	2.66	Pb/U wmean conc.	99.12	0.14	-0.0028		0.9972	
z100110	MD_3	96.35	2.72	Pb/U wmean conc.	99.12	0.14	0.0279		1.0287	
z100110	MD_4	97.24	2.59	Pb/U wmean conc.	99.12	0.14	0.0190		1.0193	
z100110	MD_5	98.32	2.63	Pb/U wmean conc.	99.12	0.14	0.0080		1.0081	
z100110	MD_6	99.50	2.65	Pb/U wmean conc.	99.12	0.14	-0.0039		0.9961	
z100110	MD_8	103.16	2.78	Pb/U wmean conc.	99.12	0.14	-0.0408		0.9608	
z100110	MD_9	99.56	2.67	Pb/U wmean conc.	99.12	0.14	-0.0044		0.9956	
z100110	TEM2_1	410.23	11.40	Pb/U wmean conc.	416.78	0.33	0.0157		1.0160	
z100110	TEM2_10	422.09	12.28	Pb/U wmean conc.	416.78	0.33	-0.0127		0.9874	
z100110	TEM2_2	418.88	11.75	Pb/U wmean conc.	416.78	0.33	-0.0050		0.9950	
z100110	TEM2_3	413.82	11.53	Pb/U wmean conc.	416.78	0.33	0.0071		1.0071	
z100110	TEM2_4	417.75	11.86	Pb/U wmean conc.	416.78	0.33	-0.0023		0.9977	
z100110	TEM2_5	427.23	11.73	Pb/U wmean conc.	416.78	0.33	-0.0251		0.9755	
z100110	TEM2_6	399.15	10.99	Pb/U wmean conc.	416.78	0.33	0.0423		1.0442	
z100110	TEM2_8	410.70	11.45	Pb/U wmean conc.	416.78	0.33	0.0146		1.0148	
z100110	TEM2_9	419.14	11.51	Pb/U wmean conc.	416.78	0.33	-0.0057		0.9944	

Results			
# ages total	47		
# ages in population	*7*		
weighted mean age ± 2σ (Ma; incl. fractionation error)	*153.05*	±	*3.69*
MSWDw	*12.93*		
P(MSWDw)	*0.000000*		

Settings	
Min Fit Quality	0.0500
Min Clean Quality	0.3333
Min 204Pb Quality	0.3333
Min Uppm Quality	9
Mineral	Zrn
# of 1σ Absolute Errors	*3.00*

APPENDIX D

A2Z Report 1131-01
U-Pb zircon analytical results

Min Fit Quality: 0.0500 | Min Clean Quality: 0.3333 | Min 204Pb Quality: 0.3333 | Min Upper Quality: 0 | Mineral (or "n.a."): Zrn | Ctrl=r reserved

Dakota Fm

Measured Isotopic Ratios

Analysis_#	207Pb/235Uc*	207Pb/235Uc* (±2σ)	206Pb/238U	206Pb/238U (±2σ)	207Pb/206Pb	207Pb/206Pb (±2σ)
131001_1_Zrn	0.1130	0.0032	0.0167	0.0004	0.0488	0.0011
131001_2_Zrn	0.1065	0.0021	0.0158	0.0004	0.0488	0.0011
131001_3_Zrn	0.1066	0.0021	0.0157	0.0004	0.0490	0.0012
131001_4_Zrn	0.1142	0.0023	0.0167	0.0004	0.0496	0.0011
131001_5_Zrn	0.5711	0.0059	0.3077	0.0061	0.1377	0.0024
131001_6_Zrn	0.1142	0.0022	0.0167	0.0005	0.0496	0.0012
131001_7_Zrn	0.1045	0.0020	0.0155	0.0004	0.0487	0.0011
131001_8_Zrn	0.1094	0.0021	0.0162	0.0004	0.0489	0.0011
131001_9_Zrn	0.1066	0.0021	0.0160	0.0004	0.0483	0.0011
131001_10_Zrn	0.1170	0.0023	0.0171	0.0005	0.0494	0.0012
131001_11_Zrn	0.1080	0.0022	0.0160	0.0004	0.0488	0.0011
131001_12_Zrn	0.1087	0.0021	0.0160	0.0004	0.0492	0.0011
131001_13_Zrn	0.1132	0.0027	0.0164	0.0005	0.0500	0.0013
131001_14_Zrn	0.1348	0.0028	0.0158	0.0004	0.0481	0.0011
131001_15_Zrn	0.1069	0.0023	0.0157	0.0004	0.0493	0.0012
131001_16_Zrn	0.1055	0.0021	0.0159	0.0004	0.0483	0.0011
131001_17_Zrn	0.1080	0.0021	0.0159	0.0004	0.0493	0.0011
131001_18_Zrn	0.1001	0.0020	0.0150	0.0004	0.0485	0.0012
131001_19_Zrn	0.1113	0.0028	0.0161	0.0005	0.0503	0.0014
131001_20_Zrn	undefined	undefined	undefined	undefined	undefined	undefined
131001_21_Zrn	0.1082	0.0022	0.0161	0.0004	0.0487	0.0012
131001_22_Zrn	0.1054	0.0020	0.0156	0.0004	0.0489	0.0011
131001_23_Zrn	0.1080	0.0022	0.0160	0.0004	0.0490	0.0012
131001_24_Zrn	0.1005	0.0020	0.0152	0.0004	0.0483	0.0012
131001_25_Zrn	0.1078	0.0023	0.0163	0.0005	0.0477	0.0012
131001_26_Zrn	0.1157	0.0025	0.0167	0.0005	0.0500	0.0013
131001_27_Zrn	0.1032	0.0023	0.0153	0.0004	0.0488	0.0013
131001_28_Zrn	0.1040	0.0021	0.0153	0.0004	0.0491	0.0012
131001_29_Zrn	0.1076	0.0021	0.0162	0.0004	0.0481	0.0011
131001_30_Zrn	0.1079	0.0022	0.0161	0.0004	0.0485	0.0012
131001_31_Zrn	0.1075	0.0025	0.0157	0.0004	0.0495	0.0013
131001_32_Zrn	0.1060	0.0021	0.0158	0.0004	0.0488	0.0011
131001_33	0.1076	0.0021	0.0161	0.0004	0.0484	0.0011
131001_34_Zrn	0.1111	0.0024	0.0164	0.0005	0.0492	0.0013
131001_35_Zrn	0.1060	0.0021	0.0158	0.0004	0.0487	0.0011
131001_36_Zrn	0.1045	0.0020	0.0158	0.0004	0.0479	0.0011
131001_37_Zrn	0.1045	0.0021	0.0156	0.0004	0.0486	0.0012
131001_38_Zrn	0.1082	0.0022	0.0159	0.0004	0.0493	0.0012
131001_39_Zrn	0.1273	0.0028	0.0177	0.0005	0.0521	0.0013
131001_40_Zrn	0.1097	0.0025	0.0165	0.0005	0.0501	0.0013
131001_41_Zrn	0.4993	0.0109	0.0659	0.0019	0.0550	0.0014
131001_42_Zrn	0.1066	0.0021	0.0160	0.0004	0.0485	0.0011
131001_43_Zrn	0.1058	0.0021	0.0158	0.0004	0.0485	0.0012
131001_44_Zrn	0.1135	0.0023	0.0165	0.0004	0.0500	0.0012
131001_45_Zrn	0.1018	0.0019	0.0151	0.0004	0.0487	0.0011
131001_46_Zrn	0.1052	0.0020	0.0154	0.0004	0.0496	0.0011
131001_47_Zrn	0.1066	0.0021	0.0158	0.0004	0.0490	0.0011
131001_48_Zrn	0.1046	0.0021	0.0155	0.0004	0.0490	0.0012
131001_49_Zrn	0.1049	0.0021	0.0152	0.0004	0.0499	0.0012
131001_50_Zrn	0.1063	0.0022	0.0155	0.0004	0.0496	0.0012

Dakota Fm

U and Th

U (ppm)	Th (ppm)	U/Th
412.91	445.67	0.927
853.73	815.39	1.047
690.82	1295.05	0.533
1147.11	1147.11	1.000
682.25	683.17	0.999
2485.42	1747.20	1.420
737.61	1090.21	0.677
969.01	1511.52	0.641
929.40	753.09	1.233
1856.87	1094.64	1.696
1 1.00	1 1.00	1.000
833.86	1162.69	0.717
561.24	783.08	0.717
1215.03	1585.16	0.766
1141.08	1393.75	0.819
475.48	706.27	0.673
1180.16	1547.12	0.763
511.79	582.26	0.879
657.72	390.17	1.686
340.35	214.75	1.585
538.64	707.02	0.761
1215.53	1719.76	0.707
1 1.00	1 1.00	1.000
716.05	724.80	0.988
723.99	786.38	0.921
356.11	352.53	1.010
354.64	192.16	1.585
618.82	740.38	0.836
1037.30	1019.78	1.018
1 1.00	1 1.00	1.000
263.99	439.47	0.600
910.86	467.15	1.950
1582.78	1597.21	0.979
587.02	620.83	0.946
1 1.00	1 1.00	1.000
1073.68	697.24	1.540
759.50	486.04	1.563
686.42	479.43	1.432
1552.16	615.64	2.521
638.02	601.93	1.060
106.16	47.74	2.224
1 1.00	1 1.00	1.000
1379.74	1797.08	0.768
461.50	264.65	1.744
679.17	675.31	1.004
877.51	739.89	1.186
1 1.00	1 1.00	1.000
725.74	533.21	1.362
530.27	715.55	0.741
566.44	418.59	1.353

Dakota Fm

Apparent Ages

Analysis_#	207Pb/235Uc*	207Pb/235Uc* (±2σ)	206Pb/238U	206Pb/238U (±2σ)	207Pb/206Pb	207Pb/206Pb (±2σ)
131001_1	108.74	2.23	106.99	2.89	140.95	54.76
131001_2	102.73	2.08	101.21	2.74	136.73	54.80
131001_3	102.88	2.15	100.64	2.71	147.43	55.80
131001_4	109.91	2.21	106.82	2.88	171.00	55.51
131001_5	1744.61	80.43	1729.39	52.33	1768.68	40.05
131001_6	109.76	2.22	106.50	2.92	177.80	54.16
131001_7	100.93	2.02	99.19	2.67	135.62	54.34
131001_8	106.39	2.13	103.67	2.79	144.92	54.25
131001_9	102.81	2.11	102.07	2.76	114.30	55.14
131001_10	112.25	2.31	109.52	3.04	168.17	54.99
131001_11	104.07	2.14	102.29	2.74	147.21	54.80
131001_12	104.76	2.11	102.28	2.74	158.33	54.11
131001_13	108.88	2.72	105.01	3.13	193.56	61.92
131001_14	161.16	1.99	100.88	2.73	106.10	53.71
131001_15	103.16	2.29	100.60	2.81	162.83	57.31
131001_16	101.85	2.14	101.59	2.80	96.45	56.65
131001_17	103.33	2.09	100.33	2.69	158.56	55.03
131001_18	96.90	2.11	96.12	2.62	124.61	57.54
131001_19	107.19	2.80	102.73	2.93	207.75	65.13
131001_20	undefined	undefined	undefined	undefined	undefined	undefined
131001_21	104.28	2.26	102.78	2.77	134.52	57.20
131001_22	101.72	2.02	99.83	2.69	148.98	53.81
131001_23	104.07	2.14	102.29	2.74	147.21	54.80
131001_24	99.88	2.04	97.17	2.60	161.04	54.44
131001_25	103.91	2.34	104.34	2.93	85.74	59.25
131001_26	111.14	2.53	106.81	2.97	196.89	58.17
131001_27	99.70	2.38	97.98	2.65	138.64	61.66
131001_28	100.42	2.15	99.13	2.64	151.05	56.59
131001_29	103.97	2.17	103.61	2.84	102.88	56.62
131001_30	104.21	2.21	101.53	2.80	161.48	56.81
131001_31	103.69	2.55	100.62	2.75	169.25	62.43
131001_32	undefined	undefined	undefined	undefined	undefined	undefined
131001_33	103.76	2.02	102.93	2.77	113.11	53.24
131001_34	106.98	2.48	104.93	2.93	157.75	59.19
131001_35	102.21	2.09	101.19	2.77	134.52	55.47
131001_36	100.88	2.06	101.19	2.74	94.70	55.49
131001_37	100.84	2.12	99.80	2.71	126.28	56.23
131001_38	104.29	2.26	101.53	2.86	163.98	56.06
131001_39	121.69	2.83	113.27	3.24	288.83	58.28
131001_40	105.70	2.58	101.49	2.95	197.38	61.23
131001_41	411.26	11.08	411.56	12.40	412.46	57.99
131001_42	102.73	2.08	101.21	2.74	136.73	54.80
131001_43	102.15	2.17	100.97	2.79	126.00	56.80
131001_44	109.16	2.36	105.23	2.87	193.27	55.94
131001_45	98.48	1.96	96.66	2.59	131.47	54.87
131001_46	101.55	2.03	98.46	2.63	174.87	53.66
131001_47	102.88	2.11	100.64	2.71	147.43	54.80
131001_48	101.02	2.12	99.11	2.67	145.52	56.00
131001_49	101.25	2.17	97.44	2.65	189.32	56.27
131001_50	102.56	2.21	99.46	2.71	177.45	56.36

Dakota Fm

Preferred Ages

Analysis_#	Preferred Age (Ma)	Absolute Error (±2σ)	Age Type	Fit Quality 1=best	Clean Quality 1=best	204Pb Quality 1=best	2σ Conc. Scans
131001_1	106.99	2.89	PbU w/mean conc.	0.9084	0.5000	1.0000	112
131001_2	101.21	2.74	PbU w/mean conc.	0.8877	1.0000	1.0000	65
131001_3	100.64	2.71	PbU w/mean conc.	0.8082	1.0000	1.0000	51
131001_4	106.82	2.88	PbU w/mean conc.	0.9000	1.0000	1.0000	130
131001_5	1760.68	40.05	PbPb w/mean conc.	0.9994	1.0000	1.0000	130
131001_6	106.50	2.92	PbU w/mean conc.	0.9881	1.0000	1.0000	77
131001_7	99.19	2.67	PbU w/mean conc.	0.6544	1.0000	1.0000	92
131001_8	103.67	2.79	PbU w/mean conc.	0.8037	1.0000	1.0000	79
131001_9	102.07	2.76	PbU w/mean conc.	0.7604	1.0000	1.0000	63
131001_10	109.52	3.04	PbU w/mean conc.	0.7806	1.0000	1.0000	79
131001_11	102.29	2.74	PbU w/mean conc.	0.8023	1.0000	1.0000	112
131001_12	102.29	2.74	PbU w/mean conc.	0.8023	1.0000	1.0000	112
131001_13	105.01	3.13	PbU w/mean conc.	0.8915	1.0000	1.0000	74
131001_14	100.88	2.73	PbU w/mean conc.	0.8615	1.0000	1.0000	88
131001_15	100.60	2.81	PbU w/mean conc.	0.7629	0.5000	1.0000	37
131001_16	101.59	2.80	PbU w/mean conc.	0.7229	1.0000	1.0000	59
131001_17	100.33	2.69	PbU w/mean conc.	0.8100	1.0000	1.0000	99
131001_18	96.12	2.62	PbU w/mean conc.	0.8547	1.0000	1.0000	97
131001_19	102.73	2.93	PbU w/mean conc.	0.9413	1.0000	1.0000	20
131001_20	1810.07	118.00	upper int.	0.9783	1.0000	1.0000	0
131001_21	102.78	2.77	PbU w/mean conc.	0.8516	1.0000	1.0000	55
131001_22	99.83	2.69	PbU w/mean conc.	0.8950	1.0000	1.0000	68
131001_23	102.29	2.74	PbU w/mean conc.	0.8023	1.0000	1.0000	112
131001_24	97.17	2.60	PbU w/mean conc.	0.9697	1.0000	1.0000	101
131001_25	104.34	2.93	PbU w/mean conc.	0.8316	1.0000	1.0000	52
131001_26	106.81	2.97	PbU w/mean conc.	0.9200	0.5000	1.0000	86
131001_27	97.98	2.65	PbU w/mean conc.	0.9346	1.0000	1.0000	60
131001_28	99.13	2.64	PbU w/mean conc.	0.8702	1.0000	1.0000	65
131001_29	103.61	2.84	PbU w/mean conc.	0.8988	1.0000	1.0000	53
131001_30	101.53	2.80	PbU w/mean conc.	0.8988	1.0000	1.0000	53
131001_31	100.62	2.70	PbU w/mean conc.	0.7278	1.0000	1.0000	62
131001_32	100.77	4.47	lower int.	0.9998	1.0000	1.0000	0
131001_33	102.93	2.77	PbU w/mean conc.	0.9991	1.0000	1.0000	108
131001_34	104.93	2.93	PbU w/mean conc.	0.9983	1.0000	1.0000	67
131001_35	101.19	2.77	PbU w/mean conc.	0.9731	1.0000	1.0000	44
131001_36	101.19	2.74	PbU w/mean conc.	0.9731	1.0000	1.0000	44
131001_37	99.80	2.71	PbU w/mean conc.	0.5888	1.0000	1.0000	60
131001_38	101.53	2.86	PbU w/mean conc.	0.9661	1.0000	1.0000	97
131001_39	113.27	3.24	PbU w/mean conc.	1.0000	1.0000	1.0000	47
131001_40	101.49	2.95	PbU w/mean conc.	0.8899	1.0000	1.0000	75
131001_41	411.56	12.40	PbU w/mean conc.	0.9150	1.0000	1.0000	117
131001_42	101.21	2.74	PbU w/mean conc.	0.8877	1.0000	1.0000	65
131001_43	100.97	2.79	PbU w/mean conc.	0.9747	1.0000	1.0000	46
131001_44	105.23	2.87	PbU w/mean conc.	0.9015	0.5000	1.0000	94
131001_45	96.66	2.59	PbU w/mean conc.	0.6949	1.0000	1.0000	104
131001_46	98.46	2.63	PbU w/mean conc.	0.9523	1.0000	1.0000	117
131001_47	100.64	2.71	PbU w/mean conc.	0.8082	1.0000	1.0000	51
131001_48	99.11	2.67	PbU w/mean conc.	0.9113	1.0000	1.0000	50
131001_49	97.44	2.65	PbU w/mean conc.	0.9803	1.0000	1.0000	77
131001_50	99.46	2.71	PbU w/mean conc.	0.9868	1.0000	1.0000	76

Dakota Fm

Preferred Ages - Sorted

Analysis_#	Preferred Age (Ma)	Absolute Error (±2σ)	Age Type	Fit Quality 1=best	Clean Quality 1=best	204Pb Quality 1=best	2σ Conc. Scans
131001_18	96.15	2.62	PbU w/mean conc.	0.8547	1.0000	1.0000	97
131001_45	96.66	2.59	PbU w/mean conc.	0.6949	1.0000	1.0000	104
131001_37	97.17	2.60	PbU w/mean conc.	0.9697	1.0000	1.0000	101
131001_24	97.17	2.60	PbU w/mean conc.	0.9697	1.0000	1.0000	101
131001_27	97.98	2.65	PbU w/mean conc.	0.9346	1.0000	1.0000	60
131001_26	98.13	2.64	PbU w/mean conc.	0.8782	1.0000	1.0000	60
131001_46	98.46	2.63	PbU w/mean conc.	0.9523	1.0000	1.0000	117
131001_47	99.11	2.67	PbU w/mean conc.	0.9113	1.0000	1.0000	50
131001_7	99.19	2.67	PbU w/mean conc.	0.6544	1.0000	1.0000	92
131001_48	99.11	2.67	PbU w/mean conc.	0.9113	1.0000	1.0000	50
131001_29	99.13	2.64	PbU w/mean conc.	0.8702	1.0000	1.0000	65
131001_30	99.13	2.64	PbU w/mean conc.	0.8988	1.0000	1.0000	53
131001_49	97.44	2.65	PbU w/mean conc.	0.9803	1.0000	1.0000	77
131001_50	99.46	2.71	PbU w/mean conc.	0.9868	1.0000	1.0000	76
131001_16	101.59	2.80	PbU w/mean conc.	0.7229	1.0000	1.0000	59
131001_15	100.60	2.81	PbU w/mean conc.	0.7629	0.5000	1.0000	37
131001_17	100.33	2.69	PbU w/mean conc.	0.8100	1.0000	1.0000	99
131001_8	103.67	2.79	PbU w/mean conc.	0.8037	1.0000	1.0000	79
131001_9	102.07	2.76	PbU w/mean conc.	0.7604	1.0000	1.0000	63
131001_10	109.52	3.04	PbU w/mean conc.	0.7806	1.0000	1.0000	79
131001_11	102.29	2.74	PbU w/mean conc.	0.8023	1.0000	1.0000	112
131001_12	102.29	2.74	PbU w/mean conc.	0.8023	1.0000	1.0000	112
131001_13	105.01	3.13	PbU w/mean conc.	0.8915	1.0000	1.0000	74
131001_14	100.88	2.73	PbU w/mean conc.	0.8615	1.0000	1.0000	88
131001_42	101.21	2.74	PbU w/mean conc.	0.8877	1.0000	1.0000	65
131001_2	101.21	2.74	PbU w/mean conc.	0.8877	1.0000	1.0000	65
131001_3	100.64	2.71	PbU w/mean conc.	0.8082	1.0000	1.0000	51
131001_47	100.64	2.71	PbU w/mean conc.	0.8082	1.0000	1.0000	51
131001_21	102.78	2.77	PbU w/mean conc.	0.8516	1.0000	1.0000	55
131001_22	99.83	2.69	PbU w/mean conc.	0.8950	1.0000	1.0000	68
131001_23	102.29	2.74	PbU w/mean conc.	0.8023	1.0000	1.0000	112
131001_44	105.23	2.87	PbU w/mean conc.	0.9015	0.5000	1.0000	94
131001_31	100.62	2.70	PbU w/mean conc.	0.7278	1.0000	1.0000	62
131001_4	106.82	2.88	PbU w/mean conc.	0.9000	1.0000	1.0000	130
131001_6	106.50	2.92	PbU w/mean conc.	0.9881	1.0000	1.0000	77
131001_35	101.19	2.77	PbU w/mean conc.	0.9731	1.0000	1.0000	44
131001_36	101.19	2.74	PbU w/mean conc.	0.9731	1.0000	1.0000	44
131001_38	101.53	2.86	PbU w/mean conc.	0.9661	1.0000	1.0000	97
131001_25	104.34	2.93	PbU w/mean conc.	0.8316	1.0000	1.0000	52
131001_29	103.61	2.84	PbU w/mean conc.	0.8988	1.0000	1.0000	53
131001_30	101.53	2.80	PbU w/mean conc.	0.8988	1.0000	1.0000	53
131001_39	113.27	3.24	PbU w/mean conc.	1.0000	1.0000	1.0000	47
131001_40	101.49	2.95	PbU w/mean conc.	0.8899	1.0000	1.0000	75
131001_41	411.56	12.40	PbU w/mean conc.	0.9150	1.0000	1.0000	117
131001_43	100.97	2.79	PbU w/mean conc.	0.9747	1.0000	1.0000	46
131001_33	102.93	2.77	PbU w/mean conc.	0.9991	1.0000	1.0000	108
131001_34	104.93	2.93	PbU w/mean conc.	0.9983	1.0000	1.0000	67
131001_1	106.99	2.89	PbU w/mean conc.	0.9084	0.5000	1.0000	112
131001_20	1810.07	118.00	upper int.	0.9783	1.0000	1.0000	0
131001_5	1760.68	40.05	PbPb w/mean conc.	0.9994	1.0000	1.0000	130
131001_32	100.77	4.47	lower int.	0.9998	1.0000	1.0000	0
131001_19	102.73	2.93	PbU w/mean conc.	0.9413	1.0000	1.0000	20

Dakota Fm

Fitted Discordia

Age - Lower Intercept (Ma)	Age - Lower Intercept (±2σ)	Age - Upper Intercept (Ma)	Age - Upper Intercept (±2σ)	Regression Intercept	Regression Intercept (±1σ)	Regression Slope	Regression Slope (±1σ)
106.58	4.41	3192.13	510.10	0.813543	0.000490	0.028269	0.004986
102.43	3.78	undefined	undefined	0.817740	0.000753	-0.016264	0.006855
103.30	3.75	undefined	undefined	0.811792	0.000610	-0.015307	0.006661
103.30	3.75	undefined	undefined	0.811792	0.000610	-0.015307	0.006661
99.53	848.88	1818.35	89.48	0.808011	0.017061	0.048596	0.003705
107.07	3.88	undefined	undefined	0.818821	0.000740	-0.017842	0.006682
100.10	4.36	3029.40	1101.59	0.812455	0.000936	0.031321	0.009940
102.87	4.19	3484.64	849.87	0.815492	0.000602	0.023351	0.005632
100.43	3.71	undefined	undefined	0.815526	0.000936	-0.007925	0.006535
105.70	4.88	3170.04	1221.08	0.813387	0.001351	0.028605	0.011181
101.43	3.82	undefined	undefined	0.815713	0.000627	-0.012380	0.005930
101.43	3.82	undefined	undefined	0.815713	0.000627	0.001380	0.005930
101.20	4.50	2534.64	393.69	0.811398	0.000501	0.042229	0.004440
102.45	4.12	2577.68	595.45	0.811650	0.000684	0.041156	0.006016
98.00	4.49	3118.35	719.97	0.812514	0.000675	0.029652	0.006119
100.61	5.47	2592.21	587.61	0.811976	0.000733	0.040383	0.006219
100.61	5.47	2592.21	587.61	0.811976	0.000733	0.040383	0.006219
97.30	3.84	2456.30	238.24	0.810624	0.000297	0.045999	0.002884
100.42	3.86	4391.85	741.57	0.814360	0.000485	0.012898	0.003986
107.25	54.32	1810.07	119.00	0.809693	0.002479	0.063598	0.002083
101.22	4.03	undefined	undefined	0.816093	0.000558	-0.002542	0.005669
99.72	3.64	4560.00	628.70	0.814937	0.000526	0.006330	0.004885
100.45	4.06	3792.94	302.90	0.812119	0.000527	0.027612	0.004835
101.40	4.06	3792.94	302.90	0.812119	0.000527	0.027612	0.004835
101.52	2.86	2948.32	546.68	0.812732	0.000501	0.033897	0.004001
108.05	4.54	undefined	undefined	0.820959	0.000921	-0.029215	0.008994
101.76	4.30	3720.83	950.80	0.813799	0.000637	0.020133	0.006531
97.45	4.12	3697.64	348.18	0.813166	0.000549	0.020495	0.005179
119.50	4.13	3130.64	626.61	0.813014	0.000560	0.029395	0.005379
102.78	4.16	2534.64	393.69	0.811398	0.000501	0.042229	0.004440
99.19	4.65	2754.80	762.71	0.812474	0.000532	0.019738	0.004441
100.77	4.47	4560.00	456.76	0.814745	0.000448	0.009694	0.002696
102.72	3.44	3817.74	547.47	0.814047	0.000524	0.018920	0.004368
100.50	5.27	4560.00	1775.30	0.814956	0.001316	0.007285	0.011354
101.13	3.96	4560.00	585.78	0.815248	0.000526	0.005437	0.005101
101.13	3.96	4560.00	585.78	0.815248	0.000526	0.005437	0.005101
100.59	3.71	undefined	undefined	0.819941	0.000599	-0.039414	0.005637
99.79	5.49	1724.29	319.05	0.808700	0.000526	0.066753	0.005068
13.50	30.84	414.95	115.75	0.800143	0.000047	0.131030	0.004253
104.30	4.96	undefined	undefined	0.818455	0.001025	-0.019364	0.008821
413.26	14.02	4560.00	657.77	0.816140	0.001560	0.010288	0.002992
101.43	3.82	undefined	undefined	0.815713	0.000627	-0.012380	0.005930
101.32	3.44	undefined	undefined	0.815584	0.000531	0.002460	0.001641
104.61	4.21	undefined	undefined	0.817864	0.001061	-0.013855	0.009156
98.21	4.65	2799.79	1301.88	0.811619	0.036259	0.041319	0.013819
98.82	3.41	4185.66	496.31	0.813932	0.000305	0.014838	0.002968
101.32	3.44	undefined	undefined	0.815584	0.000531	0.002460	0.001641
101.32	3.54	undefined	undefined	0.816398	0.000681	-0.007567	0.004406
98.32	3.57	4560.00	806.04	0.814699	0.000632	0.006598	0.005953
99.40	3.87	4560.00	621.92	0.815047	0.000611	0.004604	0.005741

Dakota Fm

Analysis Details

%Minimum Discordance	%Maximum Discordance	Preferred Discordia Scan #	Discordia Scans	2σ Conc. Scans	Relative u-Energy	Analysis_#
-999.99%	87.06%	157	161	112	0.0755	131001_1
-152.26%	85.10%	204	161	65	0.1385	131001_2
-999.43%	84.13%	113	161	51	0.1339	131001_3
-999.99%	72.46%	136	161	130	0.2539	131001_4
-13.83%	20.07%	182	161	130	2.2539	131001_5
-999.99%	68.23%	157	161	77	0.4263	131001_6
-999.99%	72.46%	136	161	92	0.1388	131001_7
-999.99%	82.66%	144	161	79	0.1841	131001_8
-673.54%	82.56%	103	161	63	0.1430	131001_9
-197.79%	81.26%	157	161	79	0.3345	131001_10
-999.99%	84.67%	230	161	112	0.1447	131001_11
-974.23%	91.06%	229	161	112	0.1447	131001_12
-999.99%	84.67%	230	161	74	0.1083	131001_13
-999.99%	68.25%	91	161	88	0.2813	131001_14
-999.99%	88.43%	110	161	37	0.1877	131001_15
-999.

See accompanying file **UPNCP.doc** for discription of each parameter and methods to calculate the values presented.

Preferred Age (Ma)	Absolute Error (±2σ)	Age Type	Analysis #	Product of Qualities
96.10	2.62	Pb/U wmean conc.	113101_18	0.8547
96.66	2.59	Pb/U wmean conc.	113101_45	0.6949
97.17	2.60	Pb/U wmean conc.	113101_24	0.9697
97.44	2.65	Pb/U wmean conc.	113101_49	0.9803
97.98	2.65	Pb/U wmean conc.	113101_27	0.9346
98.13	2.64	Pb/U wmean conc.	113101_28	0.8782
98.46	2.63	Pb/U wmean conc.	113101_46	0.9523
99.11	2.67	Pb/U wmean conc.	113101_48	0.9113
99.19	2.67	Pb/U wmean conc.	113101_7	0.4544
99.46	2.71	Pb/U wmean conc.	113101_50	0.9868
99.80	2.71	Pb/U wmean conc.	113101_37	0.5888
99.80	2.77	Pb/U wmean conc.	113101_35	0.4963
99.93	2.69	Pb/U wmean conc.	113101_22	0.6950
99.99	2.79	Pb/U wmean conc.	113101_47	0.5000
100.51	2.69	Pb/U wmean conc.	113101_42	0.8871
100.60	2.81	Pb/U wmean conc.	113101_15	0.3815
100.62	2.70	Pb/U wmean conc.	113101_31	0.7278
100.64	2.71	Pb/U wmean conc.	113101_3	0.9282
100.77	4.47	lower int.	113101_32	0.9998
100.88	2.73	Pb/U wmean conc.	113101_14	0.8615
100.97	2.78	Pb/U wmean conc.	113101_43	0.9747
101.19	2.74	Pb/U wmean conc.	113101_36	0.6721
101.21	2.74	Pb/U wmean conc.	113101_2	0.8877
101.49	2.95	Pb/U wmean conc.	113101_40	0.9999
101.53	2.86	Pb/U wmean conc.	113101_38	0.9661
101.92	2.80	Pb/U wmean conc.	113101_16	0.7510
102.05	2.74	Pb/U wmean conc.	113101_11	0.6557
102.07	2.76	Pb/U wmean conc.	113101_9	0.7654
102.29	2.74	Pb/U wmean conc.	113101_12	0.8829
102.73	2.93	Pb/U wmean conc.	113101_19	0.9413
102.75	2.77	Pb/U wmean conc.	113101_21	0.8516
102.91	2.75	Pb/U wmean conc.	113101_4	0.3045
103.33	2.80	Pb/U wmean conc.	113101_17	0.8199
103.57	2.79	Pb/U wmean conc.	113101_8	0.6927
103.81	2.84	Pb/U wmean conc.	113101_29	0.8988
104.34	2.93	Pb/U wmean conc.	113101_25	0.9316
104.38	2.86	Pb/U wmean conc.	113101_23	0.8603
104.59	2.93	Pb/U wmean conc.	113101_34	0.9983
105.01	3.13	Pb/U wmean conc.	113101_13	0.9915
105.23	2.87	Pb/U wmean conc.	113101_44	0.4507
106.50	2.92	Pb/U wmean conc.	113101_6	0.9881
106.81	2.97	Pb/U wmean conc.	113101_26	0.4600
106.99	2.89	Pb/U wmean conc.	113101_1	0.4642
109.52	3.04	Pb/U wmean conc.	113101_10	0.7806
113.27	3.24	Pb/U wmean conc.	113101_39	1.0000
411.56	12.40	Pb/U wmean conc.	113101_41	0.9150
1181.19	34.60	Pb/U wmean conc.	113101_30	0.8985
1760.66	40.05	Pb/Pb wmean conc.	113101_5	0.9994
1810.07	119.00	upper int.	113101_20	0.9783
rej 102	_ _ _ _ _ M	Pb/U wmean conc.	113101_33	0.9991
0.00	0.00	0.00	0.00	0.0000

378
379.5
381
382.5
384
385.5
387
388.5
390
391.5
393
394.5
396
397.5
399
400.5
402
403.5
405
406.5
408
409.5
411
412.5
414
415.5
417
418.5
420
421.5
423
424.5
426
427.5
429
430.5
432
433.5
435
436.5
438
439.5
441
442.5
444
445.5
447
448.5
450
451.5
453
454.5
456
457.5
459
460.5
462
463.5
465
466.5
468
469.5
471
472.5
474
475.5
477
478.5
480
481.5
483
484.5
486
487.5
489
490.5
492
493.5
495
496.5
498
499.5
501
502.5
504
505.5
507
508.5
510
511.5
513
514.5
516
517.5
519
520.5
522
523.5
525
526.5
528
529.5
531
532.5
534
535.5
537
538.5
540
541.5
543
544.5
546
547.5
549
550.5
552
553.5
555
556.5
558
559.5
561
562.5
564
565.5
567
568.5
570
571.5
573
574.5
576
577.5
579
580.5
582
583.5
585
586.5
588
589.5
591
592.5
594
595.5
597
598.5
600
601.5
603
604.5
606
607.5
609
610.5
612
613.5
615
616.5
618
619.5
621
622.5
624
625.5
627
628.5
630
631.5
633
634.5
636
637.5
639
640.5
642
643.5
645
646.5
648
649.5
651
652.5
654
655.5
657
658.5
660
661.5
663
664.5
666
667.5
669
670.5
672
673.5
675
676.5
678
679.5
681
682.5
684
685.5
687
688.5
690
691.5
693
694.5
696
697.5
699
700.5
702
703.5
705
706.5
708
709.5
711
712.5
714
715.5
717
718.5
720
721.5
723
724.5
726
727.5
729
730.5
732
733.5
735
736.5
738
739.5
741
742.5
744
745.5
747
748.5
750
751.5
753
754.5
756

1516.5
1518
1519.5
1521
1522.5
1524
1525.5
1527
1528.5
1530
1531.5
1533
1534.5
1536
1537.5
1539
1540.5
1542
1543.5
1545
1546.5
1548
1549.5
1551
1552.5
1554
1555.5
1557
1558.5
1560
1561.5
1563
1564.5
1566
1567.5
1569
1570.5
1572
1573.5
1575
1576.5
1578
1579.5
1581
1582.5
1584
1585.5
1587
1588.5
1590
1591.5
1593
1594.5
1596
1597.5
1599
1600.5
1602
1603.5
1605
1606.5
1608
1609.5
1611
1612.5
1614
1615.5
1617
1618.5
1620
1621.5
1623
1624.5
1626
1627.5
1629
1630.5
1632
1633.5
1635
1636.5
1638
1639.5
1641
1642.5
1644
1645.5
1647
1648.5
1650
1651.5
1653
1654.5
1656
1657.5
1659
1660.5
1662
1663.5
1665
1666.5
1668
1669.5
1671
1672.5
1674
1675.5
1677
1678.5
1680
1681.5
1683
1684.5
1686
1687.5
1689
1690.5
1692
1693.5
1695
1696.5
1698
1699.5
1701
1702.5
1704
1705.5
1707
1708.5
1710
1711.5
1713
1714.5
1716
1717.5
1719
1720.5
1722
1723.5
1725
1726.5
1728
1729.5
1731
1732.5
1734
1735.5
1737
1738.5
1740
1741.5
1743
1744.5
1746
1747.5
1749
1750.5
1752
1753.5
1755
1756.5
1758
1759.5
1761
1762.5
1764
1765.5
1767
1768.5
1770
1771.5
1773
1774.5
1776
1777.5
1779
1780.5
1782
1783.5
1785
1786.5
1788
1789.5
1791
1792.5
1794
1795.5
1797
1798.5
1800
1801.5
1803
1804.5
1806
1807.5
1809
1810.5
1812
1813.5
1815
1816.5
1818
1819.5
1821
1822.5
1824
1825.5
1827
1828.5
1830
1831.5
1833
1834.5
1836
1837.5
1839
1840.5
1842
1843.5
1845
1846.5
1848
1849.5
1851
1852.5
1854
1855.5
1857
1858.5
1860
1861.5
1863
1864.5
1866
1867.5
1869
1870.5
1872
1873.5
1875
1876.5
1878
1879.5
1881
1882.5
1884
1885.5
1887
1888.5
1890
1891.5
1893
1894.5

1896
1897.5
1902
1906.5
1908
1909.5
1910
1910.5
1911
1913.5
1914
1915.5
1916
1917
1918.5
1919
1920.5
1922
1923.5
1926
1927.5
1929
1931.5
1934
1935.5
1936
1937.5
1938
1939.5
1940
1941.5
1942
1943.5
1944
1945.5
1946
1947.5
1948
1949.5
1950
1951.5
1954
1955.5
1956
1957.5
1958
1959.5
1960
1961.5
1962
1963.5
1964
1965.5
1966
1967.5
1968
1969.5
1970
1971.5
1972
1973.5
1974
1975.5
1976
1977.5
1978
1979.5
1980
1981.5
1982
1983.5
1984
1985.5
1986
1987.5
1988
1989.5
1990
1991.5
1992
1993.5
1994
1995.5
1996
1997.5
1998
1999.5
2000
2001.5
2002
2003.5
2004
2005.5
2006
2007.5
2008
2009.5
2010
2011.5
2012
2013.5
2014
2015.5
2016
2017.5
2018
2019.5
2020
2021.5
2022
2023.5
2024
2025.5
2026
2027.5
2028
2029.5
2030
2031.5
2032
2033.5
2034
2035.5
2036
2037.5
2038
2039.5
2040
2041.5
2042
2043.5
2044
2045.5
2046
2047.5
2048
2049.5
2050
2051.5
2052
2053.5
2054
2055.5
2056
2057.5
2058
2059.5
2060
2061.5
2062
2063.5
2064
2065.5
2066
2067.5
2068
2069.5
2070
2071.5
2072
2073.5
2074
2075.5
2076
2077.5
2078
2079.5
2080
2081.5
2082
2083.5
2084
2085.5
2086
2087.5
2088
2089.5
2090
2091.5
2092
2093.5
2094
2095.5
2096
2097.5
2098
2099.5
2100
2101.5
2102
2103.5
2104
2105.5
2106
2107.5
2108
2109.5
2110
2111.5
2112
2113.5
2114
2115.5
2116
2117.5
2118
2119.5
2120
2121.5
2122
2123.5
2124
2125.5
2126
2127.5
2128
2129.5
2130
2131.5
2132
2133.5
2134
2135.5
2136
2137.5
2138
2139.5
2140
2141.5
2142
2143.5
2144
2145.5
2146
2147.5
2148
2149.5
2150
2151.5
2152
2153.5
2154
2155.5
2156
2157.5
2158
2159.5
2160
2161.5
2162
2163.5
2164
2165.5
2166
2167.5
2168
2169.5
2170
2171.5
2172
2173.5
2174
2175.5
2176
2177.5
2178
2179.5
2180
2181.5
2182
2183.5
2184
2185.5
2186
2187.5
2188
2189.5
2190
2191.5
2192
2193.5
2194
2195.5
2196
2197.5
2198
2199.5
2200
2201.5
2202
2203.5
2204
2205.5
2206
2207.5
2208
2209.5
2210
2211.5
2212
2213.5
2214
2215.5
2216
2217.5
2218
2219.5
2220
2221.5
2222
2223.5
2224
2225.5
2226
2227.5
2228
2229.5
2230
2231.5
2232
2233.5
2234
2235.5
2236
2237.5
2238
2239.5
2240
2241.5
2242
2243.5
2244
2245.5
2246
2247.5
2248
2249.5
2250
2251.5
2252
2253.5
2254
2255.5
2256
2257.5
2258
2259.5
2260
2261.5
2262
2263.5
2264
2265.5
2266
2267.5
2268
2269.5
2270
2271.5
2272
2273.5
2274

2275.5
2277
2278.5
2279.5
2280
2281.5
2282
2283.5
2284.5
2286
2287.5
2289
2290.5
2292
2293.5
2295
2296.5
2298
2299.5
2301
2302.5
2304
2305.5
2307
2308.5
2310
2311.5
2313
2314.5
2316
2317.5
2319
2320.5
2322
2323.5
2325
2326.5
2328
2329.5
2331
2332.5
2334
2335.5
2337
2338.5
2340
2341.5
2343
2344.5
2346
2347.5
2349
2350.5
2352
2353.5
2355
2356.5
2358
2359.5
2361
2362.5
2364
2365.5
2367
2368.5
2370
2371.5
2373
2374.5
2376
2377.5
2379
2380.5
2382
2383.5
2385
2386.5
2388
2389.5
2391
2392.5
2394
2395.5
2397
2398.5
2400
2401.5
2403
2404.5
2406
2407.5
2409
2410.5
2412
2413.5
2415
2416.5
2418
2419.5
2421
2422.5
2424
2425.5
2427
2428.5
2430
2431.5
2433
2434.5
2436
2437.5
2439
2440.5
2442
2443.5
2445
2446.5
2448
2449.5
2451
2452.5
2454
2455.5
2457
2458.5
2460
2461.5
2463
2464.5
2466
2467.5
2469
2470.5
2472
2473.5
2475
2476.5
2478
2479.5
2481
2482.5
2484
2485.5
2487
2488.5
2490
2491.5
2493
2494.5
2496
2497.5
2499
2500.5
2502
2503.5
2505
2506.5
2508
2509.5
2511
2512.5
2514
2515.5
2517
2518.5
2520
2521.5
2523
2524.5
2526
2527.5
2529
2530.5
2532
2533.5
2535
2536.5
2538
2539.5
2541
2542.5
2544
2545.5
2547
2548.5
2550
2551.5
2553
2554.5
2556
2557.5
2559
2560.5
2562
2563.5
2565
2566.5
2568
2569.5
2571
2572.5
2574
2575.5
2577
2578.5
2580
2581.5
2583
2584.5
2586
2587.5
2589
2590.5
2592
2593.5
2595
2596.5
2598
2599.5
2601
2602.5
2604
2605.5
2607
2608.5
2610
2611.5
2613
2614.5
2616
2617.5
2619
2620.5
2622
2623.5
2625
2626.5
2628
2629.5
2631
2632.5
2634
2635.5
2637
2638.5
2640
2641.5
2643
2644.5
2646
2647.5
2649
2650.5
2652
2653.5

2655
2659.5
2662
2661
2664
2665.5
2666.5
2668.5
2671
2673
2675.5
2676
2677.5
2679
2680.5
2682
2683.5
2685
2686.5
2688.5
2690.5
2692.5
2694
2695.5
2697
2699
2701.5
2703
2705.5
2706
2707.5
2709
2710.5
2713.5
2715
2716.5
2718
2719.5
2721
2722.5
2723.5
2724
2725.5
2727
2728.5
2729.5
2731.5
2733
2734.5
2736
2737.5
2739
2740.5
2742.5
2744.5
2745.5
2747.5
2748
2749.5
2751
2752.5
2754
2755.5
2757
2758.5
2760
2761.5
2763
2764.5
2766
2767.5
2769
2770.5
2772
2773.5
2775
2776.5
2778
2779.5
2781
2782.5
2784
2785.5
2787
2788.5
2790
2791.5
2793
2794.5
2796
2797.5
2799
2800.5
2802
2803.5
2805
2806.5
2808
2809.5
2811
2812.5
2814
2815.5
2817
2818.5
2820
2821.5
2823
2824.5
2826
2827.5
2829
2830.5
2832
2833.5
2835
2836.5
2838
2839.5
2841
2842.5
2844
2845.5
2847
2848.5
2850
2851.5
2853
2854.5
2856
2857.5
2859
2860.5
2862
2863.5
2865
2866.5
2868
2869.5
2871
2872.5
2874
2875.5
2877
2878.5
2880
2881.5
2883
2884.5
2886
2887.5
2889
2890.5
2892
2893.5
2895
2896.5
2898
2899.5
2901
2902.5
2904
2905.5
2907
2908.5
2910
2911.5
2913
2914.5
2916
2917.5
2919
2920.5
2922
2923.5
2925
2926.5
2928
2929.5
2931
2932.5
2934
2935.5
2937
2938.5
2940
2941.5
2943
2944.5
2946
2947.5
2949
2950.5
2952
2953.5
2955
2956.5
2958
2959.5
2961
2962.5
2964
2965.5
2967
2968.5
2970
2971.5
2973
2974.5
2976
2977.5
2979
2980.5
2982
2983.5
2985
2986.5
2988
2989.5
2991
2992.5
2994
2995.5
2997
2998.5
3000
3001.5
3003
3004.5
3006
3007.5
3009
3010.5
3012
3013.5
3015
3016.5
3018
3019.5
3021
3022.5
3024
3025.5
3027
3028.5
3030
3031.5
3033

3034.5
3036
3037.5
3039
3040.5
3042
3043.5
3045
3046.5
3048
3049.5
3051
3052.5
3054
3055.5
3057
3058.5
3060
3061.5
3063
3064.5
3066
3067.5
3069
3070.5
3072
3073.5
3075
3076.5
3078
3079.5
3081
3082.5
3084
3085.5
3087
3088.5
3090
3091.5
3093
3094.5
3096
3097.5
3099
3100.5
3102
3103.5
3105
3106.5
3108
3109.5
3111
3112.5
3114
3115.5
3117
3118.5
3120
3121.5
3123
3124.5
3126
3127.5
3129
3130.5
3132
3133.5
3135
3136.5
3138
3139.5
3141
3142.5
3144
3145.5
3147
3148.5
3150
3151.5
3153
3154.5
3156
3157.5
3159
3160.5
3162
3163.5
3165
3166.5
3168
3169.5
3171
3172.5
3174
3175.5
3177
3178.5
3180
3181.5
3183
3184.5
3186
3187.5
3189
3190.5
3192
3193.5
3195
3196.5
3198
3199.5
3201
3202.5
3204
3205.5
3207
3208.5
3210
3211.5
3213
3214.5
3216
3217.5
3219
3220.5
3222
3223.5
3225
3226.5
3228
3229.5
3231
3232.5
3234
3235.5
3237
3238.5
3240
3241.5
3243
3244.5
3246
3247.5
3249
3250.5
3252
3253.5
3255
3256.5
3258
3259.5
3261
3262.5
3264
3265.5
3267
3268.5
3270
3271.5
3273
3274.5
3276
3277.5
3279
3280.5
3282
3283.5
3285
3286.5
3288
3289.5
3291
3292.5
3294
3295.5
3297
3298.5
3300
3301.5
3303
3304.5
3306
3307.5
3309
3310.5
3312
3313.5
3315
3316.5
3318
3319.5
3321
3322.5
3324
3325.5
3327
3328.5
3330
3331.5
3333
3334.5
3336
3337.5
3339
3340.5
3342
3343.5
3345
3346.5
3348
3349.5
3351
3352.5
3354
3355.5
3357
3358.5
3360
3361.5
3363
3364.5
3366
3367.5
3369
3370.5
3372
3373.5
3375
3376.5
3378
3379.5
3381
3382.5
3384
3385.5
3387
3388.5
3390
3391.5
3393
3394.5
3396
3397.5
3399
3400.5
3402
3403.5
3405
3406.5
3408
3409.5
3411
3412.5

3414
3415
3416
3417
3418
3419
3420
3421
3422
3423
3424
3425
3426
3427
3428
3429
3430
3431
3432
3433
3434
3435
3436
3437
3438
3439
3440
3441
3442
3443
3444
3445
3446
3447
3448
3449
3450
3451
3452
3453
3454
3455
3456
3457
3458
3459
3460
3461
3462
3463
3464
3465
3466
3467
3468
3469
3470
3471
3472
3473
3474
3475
3476
3477
3478
3479
3480
3481
3482
3483
3484
3485
3486
3487
3488
3489
3490
3491
3492
3493
3494
3495
3496
3497
3498
3499
3500
3501
3502
3503
3504
3505
3506
3507
3508
3509
3510
3511
3512
3513
3514
3515
3516
3517
3518
3519
3520
3521
3522
3523
3524
3525
3526
3527
3528
3529
3530
3531
3532
3533
3534
3535
3536
3537
3538
3539
3540
3541
3542
3543
3544
3545
3546
3547
3548
3549
3550
3551
3552
3553
3554
3555
3556
3557
3558
3559
3560
3561
3562
3563
3564
3565
3566
3567
3568
3569
3570
3571
3572
3573
3574
3575
3576
3577
3578
3579
3580
3581
3582
3583
3584
3585
3586
3587
3588
3589
3590
3591
3592
3593
3594
3595
3596
3597
3598
3599
3600
3601
3602
3603
3604
3605
3606
3607
3608
3609
3610
3611
3612
3613
3614
3615
3616
3617
3618
3619
3620
3621
3622
3623
3624
3625
3626
3627
3628
3629
3630
3631
3632
3633
3634
3635
3636
3637
3638
3639
3640
3641
3642
3643
3644
3645
3646
3647
3648
3649
3650
3651
3652
3653
3654
3655
3656
3657
3658
3659
3660
3661
3662
3663
3664
3665
3666
3667
3668
3669
3670
3671
3672
3673
3674
3675
3676
3677
3678
3679
3680
3681
3682
3683
3684
3685
3686
3687
3688
3689
3690
3691
3692
3693
3694
3695
3696
3697
3698
3699
3700
3701
3702
3703
3704
3705
3706
3707
3708
3709
3710
3711
3712
3713
3714
3715
3716
3717
3718
3719
3720
3721
3722
3723
3724
3725
3726
3727
3728
3729
3730
3731
3732
3733
3734
3735
3736
3737
3738
3739
3740
3741
3742
3743
3744
3745
3746
3747
3748
3749
3750
3751
3752
3753
3754
3755
3756
3757
3758
3759
3760
3761
3762
3763
3764
3765

1264.5
1265
1265.5
1266
1266.5
1267
1267.5
1268
1268.5
1269
1269.5
1270
1270.5
1271
1271.5
1272
1272.5
1273
1273.5
1274
1274.5
1275
1275.5
1276
1276.5
1277
1277.5
1278
1278.5
1279
1279.5
1280
1280.5
1281
1281.5
1282
1282.5
1283
1283.5
1284
1284.5
1285
1285.5
1286
1286.5
1287
1287.5
1288
1288.5
1289
1289.5
1290
1290.5
1291
1291.5
1292
1292.5
1293
1293.5
1294
1294.5
1295
1295.5
1296
1296.5
1297
1297.5
1298
1298.5
1299
1299.5
1300
1300.5
1301
1301.5
1302
1302.5
1303
1303.5
1304
1304.5
1305
1305.5
1306
1306.5
1307
1307.5
1308
1308.5
1309
1309.5
1310
1310.5
1311
1311.5
1312
1312.5
1313
1313.5
1314
1314.5
1315
1315.5
1316
1316.5
1317
1317.5
1318
1318.5
1319
1319.5
1320
1320.5
1321
1321.5
1322
1322.5
1323
1323.5
1324
1324.5
1325
1325.5
1326
1326.5
1327
1327.5
1328
1328.5
1329
1329.5
1330
1330.5
1331
1331.5
1332
1332.5
1333
1333.5
1334
1334.5
1335
1335.5
1336
1336.5
1337
1337.5
1338
1338.5
1339
1339.5
1340
1340.5
1341
1341.5
1342
1342.5
1343
1343.5
1344
1344.5
1345
1345.5
1346
1346.5
1347
1347.5
1348
1348.5
1349
1349.5
1350
1350.5
1351
1351.5
1352
1352.5
1353
1353.5
1354
1354.5
1355
1355.5
1356
1356.5
1357
1357.5
1358
1358.5
1359
1359.5
1360
1360.5
1361
1361.5
1362
1362.5
1363
1363.5
1364
1364.5
1365
1365.5
1366
1366.5
1367
1367.5
1368
1368.5
1369
1369.5
1370
1370.5
1371
1371.5
1372
1372.5
1373
1373.5
1374
1374.5
1375
1375.5
1376
1376.5
1377
1377.5
1378
1378.5
1379
1379.5
1380
1380.5
1381
1381.5
1382
1382.5
1383
1383.5
1384
1384.5
1385
1385.5
1386
1386.5
1387
1387.5
1388
1388.5
1389
1389.5
1390
1390.5

1391
1391.5
1392
1392.5
1393
1393.5
1394
1394.5
1395
1395.5
1396
1396.5
1397
1397.5
1398
1398.5
1399
1399.5
1400
1400.5
1401
1401.5
1402
1402.5
1403
1403.5
1404
1404.5
1405
1405.5
1406
1406.5
1407
1407.5
1408
1408.5
1409
1409.5
1410
1410.5
1411
1411.5
1412
1412.5
1413
1413.5
1414
1414.5
1415
1415.5
1416
1416.5
1417
1417.5
1418
1418.5
1419
1419.5
1420
1420.5
1421
1421.5
1422
1422.5
1423
1423.5
1424
1424.5
1425
1425.5
1426
1426.5
1427
1427.5
1428
1428.5
1429
1429.5
1430
1430.5
1431
1431.5
1432
1432.5
1433
1433.5
1434
1434.5
1435
1435.5
1436
1436.5
1437
1437.5
1438
1438.5
1439
1439.5
1440
1440.5
1441
1441.5
1442
1442.5
1443
1443.5
1444
1444.5
1445
1445.5
1446
1446.5
1447
1447.5
1448
1448.5
1449
1449.5
1450
1450.5
1451
1451.5
1452
1452.5
1453
1453.5
1454
1454.5
1455
1455.5
1456
1456.5
1457
1457.5
1458
1458.5
1459
1459.5
1460
1460.5
1461
1461.5
1462
1462.5
1463
1463.5
1464
1464.5
1465
1465.5
1466
1466.5
1467
1467.5
1468
1468.5
1469
1469.5
1470
1470.5
1471
1471.5
1472
1472.5
1473
1473.5
1474
1474.5
1475
1475.5
1476
1476.5
1477
1477.5
1478
1478.5
1479
1479.5
1480
1480.5
1481
1481.5
1482
1482.5
1483
1483.5
1484
1484.5
1485
1485.5
1486
1486.5
1487
1487.5
1488
1488.5
1489
1489.5
1490
1490.5
1491
1491.5
1492
1492.5
1493
1493.5
1494
1494.5
1495
1495.5
1496
1496.5
1497
1497.5
1498
1498.5
1499
1499.5
1500
1500.5
1501
1501.5
1502
1502.5
1503
1503.5
1504
1504.5
1505
1505.5
1506
1506.5
1507
1507.5
1508
1508.5
1509
1509.5
1510
1510.5
1511
1511.5
1512
1512.5
1513
1513.5
1514
1514.5
1515
1515.5
1516
1516.5
1517

1517.5
1518
1518.5
1519
1519.5
1520
1520.5
1521
1521.5
1522
1522.5
1523
1523.5
1524
1524.5
1525
1525.5
1526
1526.5
1527
1527.5
1528
1528.5
1529
1529.5
1530
1530.5
1531
1531.5
1532
1532.5
1533
1533.5
1534
1534.5
1535
1535.5
1536
1536.5
1537
1537.5
1538
1538.5
1539
1539.5
1540
1540.5
1541
1541.5
1542
1542.5
1543
1543.5
1544
1544.5
1545
1545.5
1546
1546.5
1547
1547.5
1548
1548.5
1549
1549.5
1550
1550.5
1551
1551.5
1552
1552.5
1553
1553.5
1554
1554.5
1555
1555.5
1556
1556.5
1557
1557.5
1558
1558.5
1559
1559.5
1560
1560.5
1561
1561.5
1562
1562.5
1563
1563.5
1564
1564.5
1565
1565.5
1566
1566.5
1567
1567.5
1568
1568.5
1569
1569.5
1570
1570.5
1571
1571.5
1572
1572.5
1573
1573.5
1574
1574.5
1575
1575.5
1576
1576.5
1577
1577.5
1578
1578.5
1579
1579.5
1580
1580.5
1581
1581.5
1582
1582.5
1583
1583.5
1584
1584.5
1585
1585.5
1586
1586.5
1587
1587.5
1588
1588.5
1589
1589.5
1590
1590.5
1591
1591.5
1592
1592.5
1593
1593.5
1594
1594.5
1595
1595.5
1596
1596.5
1597
1597.5
1598
1598.5
1599
1599.5
1600
1600.5
1601
1601.5
1602
1602.5
1603
1603.5
1604
1604.5
1605
1605.5
1606
1606.5
1607
1607.5
1608
1608.5
1609
1609.5
1610
1610.5
1611
1611.5
1612
1612.5
1613
1613.5
1614
1614.5
1615
1615.5
1616
1616.5
1617
1617.5
1618
1618.5
1619
1619.5
1620
1620.5
1621
1621.5
1622
1622.5
1623
1623.5
1624
1624.5
1625
1625.5
1626
1626.5
1627
1627.5
1628
1628.5
1629
1629.5
1630
1630.5
1631
1631.5
1632
1632.5
1633
1633.5
1634
1634.5
1635
1635.5
1636
1636.5
1637
1637.5
1638
1638.5
1639
1639.5
1640
1640.5
1641
1641.5
1642
1642.5
1643
1643.5

1644
1644.5
1645
1645.5
1646
1646.5
1647
1647.5
1648
1648.5
1649
1649.5
1650
1650.5
1651
1651.5
1652
1652.5
1653
1653.5
1654
1654.5
1655
1655.5
1656
1656.5
1657
1657.5
1658
1658.5
1659
1659.5
1660
1660.5
1661
1661.5
1662
1662.5
1663
1663.5
1664
1664.5
1665
1665.5
1666
1666.5
1667
1667.5
1668
1668.5
1669
1669.5
1670
1670.5
1671
1671.5
1672
1672.5
1673
1673.5
1674
1674.5
1675
1675.5
1676
1676.5
1677
1677.5
1678
1678.5
1679
1679.5
1680
1680.5
1681
1681.5
1682
1682.5
1683
1683.5
1684
1684.5
1685
1685.5
1686
1686.5
1687
1687.5
1688
1688.5
1689
1689.5
1690
1690.5
1691
1691.5
1692
1692.5
1693
1693.5
1694
1694.5
1695
1695.5
1696
1696.5
1697
1697.5
1698
1698.5
1699
1699.5
1700
1700.5
1701
1701.5
1702
1702.5
1703
1703.5
1704
1704.5
1705
1705.5
1706
1706.5
1707
1707.5
1708
1708.5
1709
1709.5
1710
1710.5
1711
1711.5
1712
1712.5
1713
1713.5
1714
1714.5
1715
1715.5
1716
1716.5
1717
1717.5
1718
1718.5
1719
1719.5
1720
1720.5
1721
1721.5
1722
1722.5
1723
1723.5
1724
1724.5
1725
1725.5
1726
1726.5
1727
1727.5
1728
1728.5
1729
1729.5
1730
1730.5
1731
1731.5
1732
1732.5
1733
1733.5
1734
1734.5
1735
1735.5
1736
1736.5
1737
1737.5
1738
1738.5
1739
1739.5
1740
1740.5
1741
1741.5
1742
1742.5
1743
1743.5
1744
1744.5
1745
1745.5
1746
1746.5
1747
1747.5
1748
1748.5
1749
1749.5
1750
1750.5
1751
1751.5
1752
1752.5
1753
1753.5
1754
1754.5
1755
1755.5
1756
1756.5
1757
1757.5
1758
1758.5
1759
1759.5
1760
1760.5
1761
1761.5
1762
1762.5
1763
1763.5
1764
1764.5
1765
1765.5
1766
1766.5
1767
1767.5
1768
1768.5
1769
1769.5
1770
1770.5

1770.5
1771
1771.5
1772
1772.5
1773
1773.5
1774
1774.5
1775
1775.5
1776
1776.5
1777
1777.5
1778
1778.5
1779
1779.5
1780
1780.5
1781
1781.5
1782
1782.5
1783
1783.5
1784
1784.5
1785
1785.5
1786
1786.5
1787
1787.5
1788
1788.5
1789
1789.5
1790
1790.5
1791
1791.5
1792
1792.5
1793
1793.5
1794
1794.5
1795
1795.5
1796
1796.5
1797
1797.5
1798
1798.5
1799
1799.5
1800
1800.5
1801
1801.5
1802
1802.5
1803
1803.5
1804
1804.5
1805
1805.5
1806
1806.5
1807
1807.5
1808
1808.5
1809
1809.5
1810
1810.5
1811
1811.5
1812
1812.5
1813
1813.5
1814
1814.5
1815
1815.5
1816
1816.5
1817
1817.5
1818
1818.5
1819
1819.5
1820
1820.5
1821
1821.5
1822
1822.5
1823
1823.5
1824
1824.5
1825
1825.5
1826
1826.5
1827
1827.5
1828
1828.5
1829
1829.5
1830
1830.5
1831
1831.5
1832
1832.5
1833
1833.5
1834
1834.5
1835
1835.5
1836
1836.5
1837
1837.5
1838
1838.5
1839
1839.5
1840
1840.5
1841
1841.5
1842
1842.5
1843
1843.5
1844
1844.5
1845
1845.5
1846
1846.5
1847
1847.5
1848
1848.5
1849
1849.5
1850
1850.5
1851
1851.5
1852
1852.5
1853
1853.5
1854
1854.5
1855
1855.5
1856
1856.5
1857
1857.5
1858
1858.5
1859
1859.5
1860
1860.5
1861
1861.5
1862
1862.5
1863
1863.5
1864
1864.5
1865
1865.5
1866
1866.5
1867
1867.5
1868
1868.5
1869
1869.5
1870
1870.5
1871
1871.5
1872
1872.5
1873
1873.5
1874
1874.5
1875
1875.5
1876
1876.5
1877
1877.5
1878
1878.5
1879
1879.5
1880
1880.5
1881
1881.5
1882
1882.5
1883
1883.5
1884
1884.5
1885
1885.5
1886
1886.5
1887
1887.5
1888
1888.5
1889
1889.5
1890
1890.5
1891
1891.5
1892
1892.5
1893
1893.5
1894
1894.5
1895
1895.5
1896
1896.5


Dakota Fm
histogram
pdf
Relative Age Frequency
U-Pb Age (Ma)


Dakota Fm
pdf
U/Th
Relative Age Frequency (red curve)
U/Th (blue points)
U-Pb Age (Ma)


Dakota Fm
gauss
pdf
Relative Age Frequency
U-Pb Age (Ma)


Dakota Fm
U/Th
U/Th
U-Pb Age (Ma)


Dakota Fm
peak
Frequency
U-Pb Age (Ma)


Dakota Fm
U ppm
U ppm
U-Pb Age (Ma)


Dakota Fm
Lower-Age Intercept (Ma)
Upper-Age Intercept (Ma)


Dakota Fm
Th ppm
Th ppm
U-Pb Age (Ma)

Primary Standard

UPbICP Standard Index	6	Standard	F
Accepted Age	1099		
Isotopic Ratio	**Mean Age**	**FF Relative Error**	
207Pb/235U	1094.49	0.0087	
207Pb/235Uc	1098.24	0.0087	
206Pb/238U	1097.90	0.0131	
207Pb/206Pb	1099.00	0.0109	
208Pb/232Th	1096.60	0.0182	

Secondary Standard

UPbICP Standard Index	2	Standard	T
Accepted Age	61.23		
Uncorrected Age	59.94		

28Si 91Zr All Standards

Mineral	Zrn		
Statistics	**91Zr/28Si**	**28Si**	**91Zr**
median	0.759	2415364	1834113
standard deviation	0.114	238144	206892
minimum	not applicable	1972684	1558450
maximum	not applicable	2879354	2343284

Session	Grain	Preferred (Measured) Age (Ma)	Absolute Error (±2σ)	Age Type	Accepted Age (Ma)	Absolute Error (±2σ)	(Accepted - Preferred) / Accepted	Average (perfect=0)	Accepted / Measured	Average (perfect=1)
z100110	TR_1	60.90	1.62	Pb/U wmean conc.	61.23	0.11	0.0054	-0.0037	1.0054	0.9972
z100110	TR_2	61.92	1.66	Pb/U wmean conc.	61.23	0.11	-0.0112		0.9889	
z100110	TR_3	61.38	1.65	Pb/U wmean conc.	61.23	0.11	-0.0024		0.9976	
z100110	TR_5	58.69	1.58	Pb/U wmean conc.	61.23	0.11	0.0415		1.0433	
z100110	TR_6	60.84	1.64	Pb/U wmean conc.	61.23	0.11	0.0064		1.0065	
z100110	TR_7	60.45	1.64	Pb/U wmean conc.	61.23	0.11	0.0128		1.0130	
z100110	TR_8	62.47	1.71	Pb/U wmean conc.	61.23	0.11	-0.0203		0.9801	
z100110	TR_9	62.96	1.70	Pb/U wmean conc.	61.23	0.11	-0.0282		0.9726	
z100110	TR_11	64.25	1.83	Pb/U wmean conc.	61.23	0.11	-0.0493		0.9531	
z100110	TR_12	61.35	1.67	Pb/U wmean conc.	61.23	0.11	-0.0020		0.9980	
z100110	TR_13	61.62	1.68	Pb/U wmean conc.	61.23	0.11	-0.0064		0.9937	
z100110	TR_14	60.45	1.63	Pb/U wmean conc.	61.23	0.11	0.0127		1.0128	
z100110	FC5_1	1096.93	31.49	Pb/U wmean conc.	1099.00	0.60	0.0019		1.0019	
z100110	FC5_10	1095.36	31.62	Pb/U wmean conc.	1099.00	0.60	0.0033		1.0033	
z100110	FC5_2	1112.47	31.94	Pb/U wmean conc.	1099.00	0.60	-0.0123		0.9879	
z100110	FC5_3	1095.72	31.46	Pb/U wmean conc.	1099.00	0.60	0.0030		1.0030	
z100110	FC5_4	1100.93	31.65	Pb/U wmean conc.	1099.00	0.60	-0.0018		0.9982	
z100110	FC5_5	1120.66	32.23	Pb/U wmean conc.	1099.00	0.60	-0.0197		0.9807	
z100110	FC5_6	1112.76	31.98	Pb/U wmean conc.	1099.00	0.60	-0.0125		0.9876	
z100110	FC5_7	1098.72	31.56	Pb/U wmean conc.	1099.00	0.60	0.0003		1.0003	
z100110	FC5_8	1110.61	31.91	Pb/U wmean conc.	1099.00	0.60	-0.0106		0.9896	
z100110	FC5_9	1102.62	31.66	Pb/U wmean conc.	1099.00	0.60	-0.0033		0.9967	
z100110	IFCan_1	28.62	0.77	Pb/U wmean conc.	28.20	0.01	-0.0149		0.9854	
z100110	IFCan_10	28.49	0.76	Pb/U wmean conc.	28.20	0.01	-0.0104		0.9897	
z100110	IFCan_2	26.52	0.76	Pb/U wmean conc.	28.20	0.01	0.0595		1.0633	
z100110	IFCan_3	31.84	0.86	Pb/U wmean conc.	28.20	0.01	-0.1289		0.8858	
z100110	IFCan_4	29.34	0.79	Pb/U wmean conc.	28.20	0.01	-0.0402		0.9613	
z100110	IFCan_5	29.97	0.82	Pb/U wmean conc.	28.20	0.01	-0.0627		0.9410	
z100110	IFCan_6	26.65	0.86	Pb/U wmean conc.	28.20	0.01	0.0549		1.0581	
z100110	IFCan_7	27.64	0.73	Pb/U wmean conc.	28.20	0.01	0.0199		1.0203	
z100110	IFCan_8	27.26	0.73	Pb/U wmean conc.	28.20	0.01	0.0334		1.0346	
z100110	IFCan_9	29.37	0.96	Pb/U wmean conc.	28.20	0.01	-0.0416		0.9601	
z100110	MD_10	98.20	2.68	Pb/U wmean conc.	99.12	0.14	0.0092		1.0093	
z100110	MD_2	99.40	2.66	Pb/U wmean conc.	99.12	0.14	-0.0028		0.9972	
z100110	MD_3	96.35	2.72	Pb/U wmean conc.	99.12	0.14	0.0279		1.0287	
z100110	MD_4	97.24	2.59	Pb/U wmean conc.	99.12	0.14	0.0190		1.0193	
z100110	MD_5	98.32	2.63	Pb/U wmean conc.	99.12	0.14	0.0080		1.0081	
z100110	MD_6	99.50	2.65	Pb/U wmean conc.	99.12	0.14	-0.0039		0.9961	
z100110	MD_8	103.16	2.78	Pb/U wmean conc.	99.12	0.14	-0.0408		0.9608	
z100110	MD_9	99.56	2.67	Pb/U wmean conc.	99.12	0.14	-0.0044		0.9956	
z100110	TEM2_1	410.23	11.40	Pb/U wmean conc.	416.78	0.33	0.0157		1.0160	
z100110	TEM2_10	422.09	12.28	Pb/U wmean conc.	416.78	0.33	-0.0127		0.9874	
z100110	TEM2_2	418.88	11.75	Pb/U wmean conc.	416.78	0.33	-0.0050		0.9950	
z100110	TEM2_3	413.82	11.53	Pb/U wmean conc.	416.78	0.33	0.0071		1.0071	
z100110	TEM2_4	417.75	11.86	Pb/U wmean conc.	416.78	0.33	-0.0023		0.9977	
z100110	TEM2_5	427.23	11.73	Pb/U wmean conc.	416.78	0.33	-0.0251		0.9755	
z100110	TEM2_6	399.15	10.99	Pb/U wmean conc.	416.78	0.33	0.0423		1.0442	
z100110	TEM2_8	410.70	11.45	Pb/U wmean conc.	416.78	0.33	0.0146		1.0148	
z100110	TEM2_9	419.14	11.51	Pb/U wmean conc.	416.78	0.33	-0.0057		0.9944	

Results		
# ages total	49	
# ages in population	*45*	
weighted mean age ± 2σ (Ma; incl. fractionation error)	*101.60*	*± 2.25*
MSWDw	*5.21*	
P(MSWDw)	*0.000000*	

Settings	
Min Fit Quality	0.0500
Min Clean Quality	0.3333
Min 204Pb Quality	0.3333
Min Uppm Quality	9
Mineral	Zrn
# of 1σ Absolute Errors	*3.00*

APPENDIX E



APPENDIX E

1. *Pseudoceratium retusum* Brideaux, 1977; SV082200-2, Dakota Fm.
2. *Oligosphaeridium totum* Brideaux, 1971; SV082200-2, Dakota Fm.
3. *Cribroperidinium edwardsi* (Cookson and Eisenack, 1958) Davey, 1969; SV082200-2, Dakota Fm.
4. *Spiniferites ramosus* (Ehrenberg, 1838) Mantell, 1854; SV082200-2, Dakota Fm.
5. *Oligosphaeridium complex* (White, 1842) Davey and Williams, 1966; SV082200-2, Dakota Fm.
6. *Ovoidinium scabrosum* (Cookson and Hughes, 1964) Davey, 1970; SV082200-2, Dakota Fm.
7. *Ovoidinium verrucosum* (Cookson and Hughes, 1964) Davey, 1970; SV082200-2, Dakota Fm.
8. *Canninginopsis colliveri* (Cookson and Eisenack, 1960) Backhouse, 1988; SV082200-2, Dakota Fm.
9. *Oligosphaeridium complex* (White, 1842) Davey and Williams, 1966; SV082200-2, Dakota Fm.
10. *Cribroperidinium orthoceras* (Eisenack, 1958) Davey, 1969; SV082200-2, Dakota Fm.
11. *Tenua* sp. (*Baltisphaeridium* B of Singh, 1971); SV082200-2, Dakota Fm.
12. *Pseudoceratium regium* Singh, 1971; SV082200-2, Dakota Fm.
13. *Chlamydophorella nyei* Cookson and Eisenack, 1958; SV082200-2, Dakota Fm.
14. *Florentinia cooksoniae* (Singh, 1971) Duxbury, 1980; SV082200-2, Dakota Fm.
15. *Pterospermopsis* sp.; SV082200-2, Dakota Fm.
16. *Oligosphaeridium pulcherrimum* (Deflandre and Cookson, 1955) Davey and Williams, 1966; SV082200-2, Dakota Fm.
17. *Tasmanites* sp.; SV082200-2, Dakota Fm.
18. *Aptea polymorpha* Eisenack, 1958; SV082200-2, Dakota Fm.

REFERENCES

Backhouse, J., 1988, Late Jurassic and early Cretaceous palynology of the Perth Basin, western Australia: Geological Survey of Western Australia Bulletin 135, 233 p, 51 plates.

Brideaux, W.W., 1971, Palynology of the lower Colorado Group, central Alberta, Canada, I. Introductory remarks, Geology and microplankton studies: Palaeontographica, Abteilung B, Band 135, Lieferung 3-6, p. 53-114, plates 21-30.

Brideaux, W.W., 1977, Taxonomy of Upper Jurassic-Lower Cretaceous microplankton from the Richardson Mountains, District of Mackenzie, Canada: Geological Survey of Canada Bulletin 281, plates 1-7 p.

Cookson, I.C., and Eisenack, A., 1958, Microplankton from Australian and New Guinea upper Mesozoic sediments: Proceedings of the Royal Society of Victoria, v. 70, no. 1, p. 19-79, plates 1-12.

Cookson, I.C., and Eisenack, A., 1960, Upper Mesozoic microplankton from Australia and New Guinea: Palaeontology, v. 2, no. 2, p. 243-261, plates 37-39.

Cookson, I.C., and Hughes, N.F., 1964, Microplankton from the “Cambridge Greensand” (mid-Cretaceous): Palaeontology, v. 7, no. 1, p. 37-59, plates 5-11.

Davey, R.J., 1969, Non-calcareous microplankton from the Cenomanian of England, northern France and North America, part I: Museum (Natural History) Geology Bulletin, v. 17, p. 103-180, plates 1-11.

Davey, R.J., 1970, Non-calcareous microplankton from the Cenomanian of England, northern France and North America, part II: British Museum (Natural History) Geology Bulletin, v. 18, no. 8, p. 333-397, plates 1-10.

Davey, R.J., and Williams, G.L., 1966, V. The genus Hystrichospaeridium and its allies, *in* Davey, R.J., Downie, C., Sarjeant, W.A.S., and Williams, G.L., editors, Studies on Mesozoic and Cainozoic dinoflagellate cysts: British Museum (Natural History) Geology Bulletin, Supplement 3, p. 53-106.

Deflandre, G., and Cookson, I.C., 1955, Fossil microplankton from Australian Late Mesozoic and Tertiary sediments: Australian Journal of Marine and Freshwater Research, v. 6, no. 2, p. 242-313, plates 1-9.

Duxbury, S., 1980, Barremian phytoplankton from Speeton, east Yorkshire: Palaeontographica, Abteilung B, Band 186, Lieferung 1-3, p. 18-80, plates 1-10.

Ehrenberg, C.G., 1838, Űber das Massenverhaltniss der jetzt lebenden Kiesel-Infusorien und űber ein neues Infusorien-Conglomerat als Polierschiefer von Jastraba in Ungarn: Kőniglich Akademie der Wissenschaften zu Berlin Abhandlungen, 1836, v. 7, p. 109-135, plates 1-2.

Eisenack, A., 1958, Mikroplankton aus dem nordeutschen Apt, nebst einigen Bemerkungen uberfossile Dinoflagellaten: Neues Jahrbuch fűr Geologie und Paläontologie, Abhandlungen, v. 106, no. 3, p. 383-422, plates 21-27.

Mantell, G.A., 1854, The Medals of Creation: or first lessons in Geology and the Study of Organic Remains (second edition): London, UK, Henry G. Bohn, 930 p.

Singh, C., 1971, Lower Cretaceous microfloras of the Peace River area, northwestern Alberta: Research Council of Alberta Bulletin 28, p. 301-542, plates 39-80.

White, H.H., 1842, On fossil *Xanthida*: Microscopical Journal, v. 11, p. 35-40, plate 4.

APPENDIX F


1
2
3
4
5a
5b
6
7
8
9
10
11
12
13
14
15
16
17
18a
18b
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
0 20 40 60 80 100 µ

APPENDIX F

1. *Deltoidospora diaphana* Wilson and Webster, 1946; SV082200-2, Dakota Fm.
2. *Camarozonosporites insignis* Norris, 1967; SV082200-2, Dakota Fm.
3. *Camarozonosporites heskemensis* (Pflanzl in Murriger and Pflanzl, 1955) Krutzsch, 1959; SV082200-2, Dakota Fm.
4. *Cicatricosisporites hallei* Delcourt and Sprumont, 1955; SV082200-2, Dakota Fm.
5. *Cicatricosisporites annulatus* Archangelsky and Gamerro, 1966; a: proximal view; b: distal view; SV082200-2, Dakota Fm.
6. *Undulatisporites undulapolus* Brenner, 1963; SV082200-2, Dakota Fm.
7. *Klukisporites pseudoreticulatus* Couper, 1958; SV082200-2, Dakota Fm.
8. *Appendicisporites tricornitatus* Weyland and Greifeld, 1953; SV082200-2, Dakota Fm.
9. *Appendicisporites potomacensis* Brenner, 1963; PA060806-1, Dakota Fm.
10. *Concavissimisporites variverrucatus* (Couper, 1958) Brenner, 1963; PA060806-1, Dakota Fm.
11. *Taurocusporites segmentatus* Stover, 1962; PA060806-1, Dakota Fm.
12. *Concavissimisporites punctatus* (Delcourt and Sprumont, 1955) Brenner, 1963; PA060806-1, Dakota Fm.
13. *Costatoperforosporites foveolatus* Deák, 1962; SV082200-2, Dakota Fm.
14. *Neoraistrickia robusta* Brenner, 1963; PA060806-1, Dakota Fm.
15. *Gleicheniidites senonicus* Ross, 1949; SV082200-2, Dakota Fm.
16. *Gleicheniidites senonicus* Ross, 1949; SV082200-2, Dakota Fm.
17. *Trilobosporites minor* Pocock, 1962; SV082200-2, Dakota Fm.
18. *Asteropollis asteroides* Hedlund and Norris, 1968; SKM05-20-05-04, Dakota Fm., a: surface view; b: optical section view.
19. *Densoisporites microrugulatus* Brenner, 1963; PA060806-1, Dakota Fm.
20. Bisaccate sp.; SV082200-2, Dakota Fm.
21. *Taxodiaceaepollenites hiatus* (Potonie, 1961) Kremp, 1949; SV082200-2, Dakota Fm.
22. *Classopollis classoides* Pflug, 1953; SV082200-2, Dakota Fm.
23. *Rouseisporites reticulatus* Pocock, 1962; PA060806-1, Dakota Fm.
24. *Psilatriletes circumundulatus* Brenner, 1963; PA060806-1, Dakota Fm.
25. *Rugubivesiculites rugosus* Pierce, 1961; SV082200-2, Dakota Fm.
26. *Cerebropollenites mesozoicus* (Couper, 1958) Nilsson, 1958; SV082200-2, Dakota Fm.
27. *Reticulatisporites* sp.; PA060806-1, Dakota Fm.
28. *Pilosisporites trichopapillosus* (Thiergart, 1949) Delcourt and Sprumont, 1955; PA060806-1, Dakota Fm.
29. Bisaccate sp.; SV082200-2, Dakota Fm.
30. *Araucariacites australis* Cookson and Hughes, 1964; SV082200-2, Dakota Fm.
31. *Callialasporites dampieri (Balme, 1957)* Sukh, 1961; PA060806-1, Dakota Fm.
32. *Trilobosporites crassus* Brenner, 1963; PA060806-1, Dakota Fm.

33. Trilobosporites apiverrucatus Couper, 1958; PA060806-1, Dakota Fm.
34. Trilobosporites apiverrucatus Couper, 1958; PA060806-1, Dakota Fm.
35. *Pilosisporites trichopapillosus* (Thiergart, 1949) Delcourt and Sprumont, 1955; PA060806-1, Dakota Fm.
36. *Callialasporites dampieri* (Balme, 1957) Sukh, 1961; PA060806-1, Dakota Fm.

REFERENCES

Archangelsky, S., and Gamerro, J.C., 1966, Estudio palinológico de la Formación Baqueró (Cretácico), provincia de Santa Cruz II: Ameghiniana, v. 4, no. 6, p. 201-209.

Balme, B.E., 1957, Spores and pollen grains from the Mesozoic of Western Australia: The Commonwealth Scientific and Industrial Research Organisation (C.S.I.R.O.), Australia Coal Research Section, Reference T.C. 25, p. 1-48, plates 1-11.

Brenner, G.J., 1963, Spores and pollen of the Potomac Group of Maryland: Maryland Department of Geology, Mines, and Water Resources Bulletin 27, 215 p, 43 plates.

Cookson, I.C., and Hughes, N.F., 1964, Microplankton from the “Cambridge Greensand” (mid-Cretaceous): Palaeontology, v. 7, no. 1, p. 37-59, plates 5-11.

Couper, R.A., 1958, British Mesozoic microspores and pollen grains, a systematic and stratigraphic study: Palaeontographica, Abteilung B, Band 103, Lieferung 4-6, p. 75-179.

Deák, M.H., 1962, Deux nouveaux genres de spore de la série d’argiles et de marmes aptoemmes: Földtani Kőzlőny, v. 92, no. 2, p. 230-235.

Delcourt, A.F., and Sprumont, F., 1955, Les spores et grains de pollen du Wealdien du Hainaut: Mémoires de la Société Belge de Géologie, de Paléontologie et d'Hydrologie, v. 5, p. 1-73.

Hedlund, R.W., and Norris, G., 1968, Spores and pollen grains from Fredericksburgian (Albian) strata, Marshall County, Oklahoma: Pollen et Spores, v. 10, no. 1, p. 129-159.

Kremp, G., 1949, Pollenanalytische Untersuchung des Miozänen Braunkohlenlagers von Konin an der Warthe: Palaeontographica, Abteilung B, Band 90, Lieferung 1-3, p. 53-93, plates 3-9.

Krutzsch, W., 1959, Mikropälaontologische (sporenpälaontologische) Untersucungen in der Braunkohle des Geiseltales: Geologie, Jahr 8, 425 p.

Murriger, F., and Pflanzl, G., 1955, Pollenanalytische Datierungen einiger hessischer Braunkohlen: Notizbl. Hess. L.- Amt. Bodenforsch, v. 83, no. 3, p. 71-89.

Nilsson, T., 1958, Űber das Vorkommen eines mesozoischen Sapropelogesteins in Schonen, Lunds Universitets Årsskrift, N. F., Avd. 2, Band 54, no. 10, p. 112.

Norris, G., 1967, Spores and pollen from the lower Colorado Group (Albian-?Cenomanian) of central Alberta: Palaeontographica, Abteilung B, Band 120, Lieferung 1-4, p. 2-115.

Pflug, H.D., 1953, Zur Entstehung und Entwicklung des Angiospoeriden Pollens in der Erdgeschichte: Palaeontogaphica, Abteilung B, Band 96, Lieferung 4-6, p. 60-171, plates 15-25.

Pierce, R.L., 1961, Lower Upper Cretaceous plant microfossils from Minnesota: Minnesota Geological Survey Bulletin 42, 86 p.

Pocock, S.A.J., 1962, Microfloral analysis and age determination of strata at the Jurassic-Cretaceous boundary in the Western Canada Plains: Palaeontographica, Abteilung B,

Band 111, Lieferung 1-3, p. 1-95.
Potonie, R., 1961, Pollenformen aus tertiaeren Braunkohlen (3. Mitt.): Jb. Preuss. Geol. L.A., v. 52, plates 1-7.
Ross, N.E., 1949, On a Cretaceous pollen and spore-bearing clay deposit of Scania: Bulletin of the Geological Institution of the University of Upsala, v. 34, p. 25-43.
Stover, L.E., 1962, *Taurocusporites*, a new trilete spore genus from the Lower Cretaceous of Maryland: Micropaleontology, v. 8, no. 1, p. 55-59.
Sukh, D., 1961, The fossil flora of the Jabalpur series - 3: spores and pollen grains: Palaeobotanist, v. 8, no. 1-2, p. 43-56, plates 1-8.
Thiergart, F., 1949, Der stratigraphische Wert mesozoischer Pollen und Sporen: Palaeontographica, Abteilung B, Band 94, Lieferung 1-4, p. 1-138.
Weyland, H., and Greifeld, G., 1953, Űber strukturbietende Blätter und pflanzliche Mikrofossilien aus den untersenonen Tonen der Gegend von Quedlinberg: Palaeontographica, Abteilung B, Band 95, Lieferung 1-3, p. 30-52.
Wilson, L.R., and Webster, R.M., 1946, Plant microfossils from a Fort Union Coal of Montana: American Journal of Botany, v. 33, no. 4, p. 271-278.