

GILSONITE VEINS OF THE UINTA BASIN, UTAH

by Taylor Boden and Bryce T. Tripp



SPECIAL STUDY 141
UTAH GEOLOGICAL SURVEY
a division of
UTAH DEPARTMENT OF NATURAL RESOURCES
2012

GILSONITE VEINS OF THE UINTA BASIN, UTAH

by Taylor Boden¹ and Bryce T. Tripp²

¹Utah Geological Survey

²Utah Geological Survey, retired

Cover photo: Red Wash gilsonite vein cropping out in member C of the Uinta Formation in NW¼NE¼ section 19, T. 8 S., R. 23 E., Salt Lake Base Line and Meridian, Uintah County.

ISBN 978-1-55791-856-7



SPECIAL STUDY 141
UTAH GEOLOGICAL SURVEY
a division of
UTAH DEPARTMENT OF NATURAL RESOURCES
2012

STATE OF UTAH

Gary R. Herbert, Governor

DEPARTMENT OF NATURAL RESOURCES

Michael Styler, Executive Director

UTAH GEOLOGICAL SURVEY

Richard G. Allis, Director

PUBLICATIONS

contact

Natural Resources Map & Bookstore

1594 W. North Temple

Salt Lake City, UT 84116

telephone: 801-537-3320

toll-free: 1-888-UTAH MAP

website: mapstore.utah.gov

email: geostore@utah.gov

UTAH GEOLOGICAL SURVEY

contact

1594 W. North Temple, Suite 3110

Salt Lake City, UT 84116

telephone: 801-537-3300

website: geology.utah.gov

Although this product represents the work of professional scientists, the Utah Department of Natural Resources, Utah Geological Survey, makes no warranty, expressed or implied, regarding its suitability for a particular use. The Utah Department of Natural Resources, Utah Geological Survey, shall not be liable under any circumstances for any direct, indirect, special, incidental, or consequential damages with respect to claims by users of this product.

CONTENTS

| | |
|---|----|
| ABSTRACT..... | 1 |
| INTRODUCTION..... | 1 |
| Definitions and Gilsonite History..... | 2 |
| Study Area..... | 4 |
| Project Background..... | 4 |
| Scope..... | 4 |
| Methods..... | 5 |
| Previous Work..... | 5 |
| GILSONITE USES..... | 5 |
| STRUCTURAL SETTING..... | 6 |
| STRATIGRAPHIC SETTING..... | 7 |
| ORIGINS OF GILSONITE..... | 9 |
| EXPLORATION TECHNIQUES..... | 10 |
| DEVELOPMENT..... | 11 |
| Mining History..... | 11 |
| Mining Methods and Transportation..... | 13 |
| HEALTH ISSUES..... | 14 |
| GILSONITE VEINS..... | 15 |
| Bonanza Area Veins..... | 15 |
| Red Wash and Little Red Wash..... | 15 |
| Cowboy (Eureka)..... | 15 |
| Independent (Bonanza), Tabor, and Little Bonanza..... | 15 |
| Little Chepetta, Chepetta, and Caldwell..... | 17 |
| Wagon Hound..... | 18 |
| Little Emma (Uinta)..... | 18 |
| Augustine..... | 18 |
| Weaver..... | 18 |
| Rainbow Area Veins..... | 19 |
| Little Butte..... | 19 |
| Rustler..... | 19 |
| Alabama..... | 19 |
| Little Alabama..... | 19 |
| Black Dragon..... | 19 |
| Rainbow..... | 19 |
| Pride of the West..... | 20 |
| Harrison..... | 21 |
| Little Asphalt Wash..... | 21 |
| Asphalt Wash..... | 21 |
| Snow..... | 22 |
| Saddletree..... | 22 |
| South Harrison..... | 22 |
| Little Chief..... | 22 |
| Fox..... | 22 |
| Kings Well..... | 23 |
| Neal..... | 23 |
| Bitter Water..... | 23 |
| Long Draw..... | 23 |
| Sage Brush..... | 24 |
| Antler and Little Antler..... | 24 |
| Little Bitter Creek #1..... | 25 |
| Ouray Area Veins..... | 25 |
| Jumbo..... | 25 |
| Pride of Utah..... | 25 |
| Antelope..... | 26 |
| Unreported Occurrences..... | 26 |
| Cottonwood..... | 26 |
| Black Cat..... | 26 |
| Willow Creek/Wall..... | 27 |
| Ouray..... | 27 |

| | |
|---|----|
| Little Seam..... | 27 |
| Gem..... | 27 |
| Little Boy..... | 27 |
| Cliff Dweller..... | 27 |
| Willow Group..... | 28 |
| Turtle Shell..... | 29 |
| Petroglyph..... | 29 |
| Black Diamond..... | 29 |
| Turkey Trail Hill..... | 30 |
| Willow Creek #1 and #2..... | 30 |
| O.K., Original Owner, Black Dog, and Florence..... | 30 |
| Pariette Area Veins..... | 31 |
| Castle Peak, Baxter, and Pariette..... | 31 |
| Gilsonite Draw..... | 31 |
| Fort Duchesne Area Veins..... | 31 |
| U.S. Bureau of Land Management Gilsonite Mapping..... | 31 |
| Unconfirmed Deposits..... | 32 |
| CONCLUSION..... | 33 |
| ACKNOWLEDGMENTS..... | 33 |
| REFERENCES..... | 34 |
| SUPPLEMENTAL BIBLIOGRAPHY..... | 36 |
| APPENDIX..... | 45 |

FIGURES

| | |
|---|----|
| Figure 1. Uinta Basin location map..... | 1 |
| Figure 2. Gilsonite textures showing columnar structure and conchoidal fracturing..... | 2 |
| Figure 3. Hunt's (1979) classification of natural bitumens and coals..... | 4 |
| Figure 4. Tertiary basin and uplift structures, and physiographic features in relation to study area..... | 6 |
| Figure 5. Stratigraphic column of the Eocene section hosting gilsonite veins in the Uinta Basin..... | 7 |
| Figure 6. Cross section of a typical gilsonite vein..... | 8 |
| Figure 7. Gilsonite sill on the Rainbow vein in the lower Uinta Formation..... | 9 |
| Figure 8. Gilsonite float eroding from the northwest segment of the Harrison vein..... | 10 |
| Figure 9. Gilsonite annual production and value..... | 12 |
| Figure 10. Gilsonite underground mining methods..... | 13 |
| Figure 11. Red Wash and Little Red Wash veins cropping out in member C of the Uinta Formation..... | 17 |
| Figure 12. Black Dragon vein cropping out in the Douglas Creek Member of the Green River Formation..... | 20 |
| Figure 13. Mined open cut on the Rainbow vein..... | 20 |
| Figure 14. Harrison vein cropping out in the Uinta Formation..... | 21 |
| Figure 15. Antler vein outcrop showing the main vein and adjacent secondary vein..... | 24 |
| Figure 16. Prospect on the Jumbo vein..... | 25 |
| Figure 17. Prospect shaft on the Cliff Dweller vein..... | 28 |
| Figure 18. Black Diamond mine..... | 29 |

TABLES

| | |
|--|----|
| Table 1. Physical and chemical characteristics of gilsonite..... | 3 |
| Table 2. Summary of gilsonite vein attribute data..... | 16 |

PLATE

| | |
|--|-------|
| Plate 1. Gilsonite veins of the Uinta Basin..... | on CD |
|--|-------|

GILSONITE VEINS OF THE UINTA BASIN, UTAH

by Taylor Boden and Bryce T. Tripp

ABSTRACT

Gilsonite is a solid hydrocarbon that forms a swarm of subparallel, northwest-trending, near-vertical, laterally and vertically extensive veins in the Uinta Basin of Utah and Colorado. The Uinta Basin hosts the world's largest deposits of gilsonite and is the only place where gilsonite is economically produced in large quantities. Gilsonite is sourced from the Mahogany oil shale zone of the Green River Formation and is hosted in the Tertiary Wasatch, Green River, Uinta, and Duchesne River Formations. The veins formed in two stages associated with thermal maturation of the Mahogany oil shale. Overpressuring deep in the Uinta Basin expelled large quantities of thermal water from the reservoir rocks and hydrofractured the overlying strata. Subsequently, thick liquid gilsonite was expelled from the reservoir rocks, forcing open the existing fractures in the overlying strata. The gilsonite later solidified in these fractures, probably primarily through cooling and polymerization.

This study included examination and mapping of 59 veins, vein systems, and isolated vein outcrops totaling more than 120 miles in length. In addition, we collected 1474 Global Positioning System data points with associated attribute data, obtained field data from previous geologic mapping by the U.S. Geological Survey and U.S. Bureau of Land Management, and examined recent National Agriculture Imagery Program high-resolution color aerial photography. A total of 71 significant veins, vein systems, and vein extensions were documented in our study, having a total combined vein length of more than 170 miles.

Gilsonite is a valuable resource and has a wide variety of uses, including asphalt paving mixes and coatings; chemical components in metallurgical, adhesive, coating, binder, ink, and paint products; and uses in metal foundry and oil well drilling and well completions. Even though significant amounts of the approximately 45-million-short-ton original gilsonite resource have been mined, millions of tons of the resource still remain. This resource tends to be in the deeper parts of the veins and in thinner, more remote veins that will likely be more expensive to mine than veins mined in the past. At the recent industry production rate of 60,000 to 80,000 tons per year, gilsonite could continue to be mined in the Uinta Basin for decades.

INTRODUCTION

The Uinta Basin of northeastern Utah and northwestern Colorado (figure 1) contains vast hydrocarbon deposits. Oil, natural gas, tar sand, oil shale, coal, and solid hydrocarbons (asphaltites), including gilsonite, wurtzilite, tabbyite, and ozokerite, have a long history of exploration and/or production in the basin. The Uinta Basin is the only place where gilsonite is economically produced in large quantities, and contains the world's largest deposits. Gilsonite is remarkable for its unusual geologic origin, chemical and physical properties, and industrial uses. Also notable are the ingenuity and persistence of the gilsonite pioneers who created a new industry, and over the past 100 years have solved mining, processing, transportation, marketing, and other challenges to continue to supply this unique material to world markets.

The gilsonite veins in Utah are the largest in the world and have a long, colorful history of profitable mining. Consequently, gilsonite has been studied and described




Figure 1. Uinta Basin location map.

in a large body of work dating back to the 1880s. However, some ambiguity has always existed in the descriptions of the veins, especially in their locations and extent, so accurate summaries have been lacking. The goal of the Utah Geological Survey (UGS) study was to use Global Positioning System (GPS) and geographic information system technology to generate accurate maps of the gilsonite veins, collect vein attribute data, and combine these new data with existing data into an accurate, up-to-date compilation of Utah's gilsonite resource. Historical production data and an extensive bibliography were also compiled. All of the data were compiled into a map (except for isolated vein outcrops) and accompanying text, which includes detailed descriptions of the character of the gilsonite deposit for each vein, vein system, vein extension, isolated vein outcrop, and evidence of mining and prospecting.

Definitions and Gilsonite History

Gilsonite is naturally occurring hydrocarbon bitumen that occurs in dikes (veins), sills, fracture fillings, and disseminated blebs, commonly in association with oil shale and tar sand. Gilsonite has a dull, black, coal-like appearance on weathered surfaces and a shiny, black, obsidian-like appearance on fresh surfaces. Fracture surfaces vary from conchoidal to columnar (pencillated) to flaky or scaly (figure 2). Occasionally gilsonite in deep parts of some veins is in a semi-solid state. Industry once defined three major subdivisions of gilsonite based on appearance and fusing temperature: selects, seconds, and jet. Select material is very shiny, fuses from 300 to 334°F, and tends to occur in centers of veins. Seconds are somewhat duller than selects, fuse from 306 to 361°F, and tend to occur along vein margins, sometimes having columnar jointing (pencillated texture). Pencillated tex-





Figure 2. Gilsonite textures: **A.** gilsonite showing columnar "pencillated" structure, and **B.** "select" gilsonite from underground workings on the Independent vein showing conchoidal fracturing.

ture forms at right angles to vein walls and penetrates about 6 inches into the ore (Verbeek and Grout, 1993). Gilsonite in some localities also has flaky or scaly texture. A third, unusual variety, called jet gilsonite, has a brilliantly shiny surface, a bluish-black color, and fuses from 390 to 446°F (Abraham, 1960). To date, it has been found only in the Cowboy vein. Gilsonite is now classified into five ranges, according to fusing temperature, which are used in different applications and sell for different prices. Gilsonite from different veins or different parts of veins is sometimes mixed to achieve a product with a specific fusing temperature range. Physical and chemical characteristics of gilsonite (table 1) are important for differentiating it from other asphaltites and in determining possible industrial applications.

Gilsonite, named after Samuel H. Gilson, was discovered in the 1860s. Gilson was not one of the original discoverers of gilsonite, but his enthusiastic development and promotional efforts linked the material to him, and people in the region began referring to this material as gilsonite rather than using its scientific name, “uintahite” (Kretchman, 1957; Covington, 1964). The name gilsonite further solidified in common usage when an early mining company adopted and trademarked the name.

People in the late 1800s were uncertain about the exact nature of gilsonite, whether it was asphaltum, coal, or mineral wax. Samples were shipped to several scientists for examination. Wurtz (1869) described a sample in the Columbia College School of Mine’s collection and noted the similarity of this material to the grahamite that is present in dikes in Ritchie County, West Virginia. Blake (1890) examined this same sample and determined it to be “uintahite,” an asphaltite that he had described in an earlier scientific paper (Blake, 1885).

Gilsonite is classified as a member of the asphaltite group of hydrocarbon bitumens. These are a group of naturally occurring solid hydrocarbons that are somewhat similar in appearance, occurrence, and properties. Scientists describing these hydrocarbons have developed various classification schemes for them. Abraham (1960) developed a systematic classification scheme based on physical and chemical characteristics such as solubility, physical state (solid or liquid), fusibility, and oxygen content. Hunt (1963) slightly modified Abraham’s classification system. King and others (1963) devised a classification scheme starting with the geologic origin of the hydrocarbon. Hunt (1979) combined aspects of Abraham’s and King and others’ classifications while deleting petroleum from the scheme and subdividing some of the hydrocarbons based on their hydrogen-to-carbon ratios (figure 3). Other proposed hydrocarbon classification systems have been published by Rogers and others (1974), Jacob (1983), Curiale (1986), and Cornelius (1987), but Hunt’s (1979) classification scheme is commonly accepted and is easy to apply using basic laboratory data.

Table 1. Physical and chemical characteristics of gilsonite.

| Characteristic | Value | Data Source |
|-------------------------------------|--|---------------------------|
| Color | black | - |
| Fracture | usually conchoidal, occasionally columnar to platy | - |
| Luster | bright | - |
| Streak | brown | Abraham (1960) |
| Hardness (Mohs scale) | 2 | Abraham (1960) |
| Specific gravity (g/cc) | 1.03–1.10 | Abraham (1960) |
| Bulk density (kg/m ³) | 641 | Ziegler (2008) |
| Solubility in CS ₂ (%) | 98 | Abraham (1960) |
| Solubility in petroleum naphtha (%) | 10–60 | Abraham (1960) |
| Carbon (wt %) | 85–86 | Abraham (1960) |
| Hydrogen (wt %) | 8.5–10 | Abraham (1960) |
| Sulfur (wt %) | 0.22–0.53 | Bell and Hunt (1963) |
| Nitrogen (wt %) | 2.25–3.29 | Bell and Hunt (1963) |
| Oxygen (wt %) | 1.5 | Wen and others (1978) |
| Refractive index | 1.59–1.64 | Bell and Hunt (1963) |
| H/C ratio | 1.42–1.47 | Bell and Hunt (1963) |
| Fusing (softening) point (°C) | 161–230 | Bell and Hunt (1963) |
| (ring and ball method) | | |
| Moisture content (wt %) | 0.5 | Ziegler (2008) |
| Heating value (J/kg) | 4.2 x 10 ⁷ | Ziegler (2008) |
| Specific heat (168°C) | 0.52 | Ziegler (2008) |
| Penetration (28°C) | 0–3 | Ziegler (2008) |
| Porphyrim content (wt %) | 0.004–0.03 | McGee (1956) |
| (nickel complex) | | |
| Molecular weight | 8,130–12,300 | Dickie and Yen (1967) |
| Saturated hydrocarbons (wt %) | 2.4–2.8 | American Gilsonite (2008) |
| Asphaltenes (wt %) | 56.7–76.2 | American Gilsonite (2008) |
| Resins (maltenes, wt %) | 21.0–26.7 | American Gilsonite (2008) |
| Volatile matter | 75.2–86.5 | American Gilsonite (2008) |
| (wt %, dry, ash-free basis) | | |
| Fixed carbon (wt %) | 13.8–18.6 | Hunt (1963) |
| Ash (wt %) | 0.3–0.7 | American Gilsonite (2008) |
| Resistivity (ohm-m) | 1.9 x 10 ¹⁰ | Neel (1980) |
| Reflectance (% RM) | 0.11–0.13 | Jacob (1983) |




Figure 3. Hunt's (1979) classification of natural bitumens and coals.

Study Area

The study area (plate 1) is principally in Uintah County and to a lesser extent in Duchesne County. Gilsonite veins are exposed in the vicinity of the Green and White Rivers, south of the towns of Roosevelt and Vernal. Gilsonite forms a subparallel set of relatively straight, linearly continuous, vertical veins in a northwest-southeast-trending zone approximately 64 miles long by 30 miles wide. The landscape is moderately dissected, and typified by small mesas, low buttes, dry washes, and sparse vegetation.

Project Background

This project began as an evaluation of the gilsonite resources of the Uinta Basin for the Utah School and Institutional Trust Lands Administration (SITLA). This evaluation included compilation of existing data, as well as examination and GPS plots of most known veins. Summary reports of this work for SITLA were published by Tripp (2004), Tripp and White (2006), and Boden and Tripp (2008).

Scope

Although many good, early summary reports have been published, they lack detailed and/or accurate vein information. Reliable, detailed information about gilsonite deposits in large areas of the Uinta Basin is contained only in proprietary company files and has not been published. The lack of detailed, accurate published data is partially due to the remoteness of the gilsonite veins, poor-quality base maps at the time of some of the earlier studies, and the greater difficulty of accurately plotting more than 100 miles of veins before GPS technology was available. While some of the largest and most economically significant veins are well described in published reports, other veins containing significant resources are not well described. Therefore, for this investigation we sought to locate, accurately map, and collect data on all the gilsonite veins, especially the "lesser-known" deposits. This study fills many of the data gaps; however, we were unable to obtain access to gilsonite deposits on the Uintah and Ouray Indian Reservation, so veins located there remain inadequately described. Also, we were

unable to examine deposits in the area of Fort Duchesne where the deposits have been obscured by development.

Methods

Mapping of the gilsonite veins in the study area was accomplished using recreational grade Garmin GPS equipment with horizontal accuracy of approximately 15 feet, and U.S. Geological Survey (USGS) 7.5-minute topographic quadrangle maps. National Agriculture Imagery Program (NAIP) 1 meter, high-resolution, color aerial photography of 2006 from the Utah Automated Geographic Reference Center was also used to locate and map veins. We used the gilsonite deposit maps produced by Pruitt (1961) and Cashion (1967) as initial base maps to approximately locate reported veins. The Cowboy, Independent/Tabor, Little Bonanza, Little Chepetta, Chepetta, Caldwell, Augustine, and Colorado segment of the Weaver vein locations were digitized from previous USGS mapping (Cashion, 1974, 1977, 1978, and 1986). The locations of the digitized vein traces were improved through examination of NAIP aerial photography from 2006. We used early work by Cashion (1967) for general geological information as it relates to the gilsonite deposits. To compile the geology shown on the gilsonite veins map (plate 1), we adopted the geologic map units developed during recent mapping in the Uinta Basin by Sprinkel (2007, 2009).

Previous Work

Gilsonite is a geologically interesting and economically significant resource, so it has been the subject of numerous studies and publications from the late 1800s to the present. Some of the more important publications include those of Eldridge (1901), who gave an early description of the gilsonite deposits and geologic setting. Abraham (1960), Bell and Hunt (1963), Hatcher and others (1992), Hunt (1963, 1979), Hunt and others (1954), and Monson and Parnell (1992a, 1992b) investigated origins and compositions of gilsonite deposits and various other hydrocarbons in the Uinta Basin. Verbeek and Grout (1992, 1993) studied the geometry and structural evolution of the gilsonite veins. Henderson (1957) described the refining of gilsonite by the American Gilsonite Company. Covington (1964), Kretchman (1957), Remington (1959), Tripp (2004), and Tripp and White (2006) described the history of the gilsonite industry and mining. Cashion (1967) described the geologic setting and estimated gilsonite reserves. Boden and Tripp (2008), Crawford and Pruitt (1963), Davis (1957), and Pruitt (1961) described the location, occurrence, and geologic setting of gilsonite deposits.

In the early 1980s, U.S. Bureau of Land Management (BLM) employees mapped selected gilsonite veins on the Ouray, Ouray SE, Big Pack Mtn. NW, Big Pack Mtn. NE,

Archy Bench, Asphalt Wash, and Rainbow 7.5-minute topographic quadrangle maps (Peter Sokolosky, BLM, written communication, April 2, 2004). The survey appears to have focused on veins in old BLM prospecting permit application areas. Many of the veins mapped by the BLM are on the Uintah and Ouray Indian Reservation and were not visited by UGS geologists. However, UGS geologists observed some of these veins in road cuts and outcrops along the public roads that pass through the reservation. In the areas covered by both field investigations, mapping mostly agrees except for small discrepancies. Plate 1 shows the Pride of Utah, Natural Buttes, Black Gnat, Workman, Willow #2 southeast extension, Black Bridge, and southeast extensions of the Willow Creek #1 and #2 veins as mapped by the BLM, most of which are on the Uintah and Ouray Indian Reservation.

GILSONITE USES

Gilsonite has an extremely wide range of uses that have changed over time with new technology and industrial needs. Kemmerer (1934), Carey and Roberts (1949), Davis (1951a, 1951b), Kretchman (1957), and Remington (1959) give good summaries of past gilsonite uses, which include use as a component of the durable black paint used by Ford Motor Company on early Model T automobiles. Gilsonite has hundreds of uses, but they can be grouped into the following five major categories (American Gilsonite Company, 2008):

- 1) Asphalt paving mixes and coatings – gilsonite added to asphalt paving mixes (black top) improves pavement performance. Gilsonite can also be incorporated into solvent-based and emulsion-type pavement sealers.
- 2) Chemical products – gilsonite is used in metallurgy, in adhesives and coatings, as a binder in refractories, and as a binder in brake pads.
- 3) Metal foundry uses – gilsonite is used as an additive in molding sand mixes in iron and steel casting.
- 4) Inks and paints – gilsonite, added to ink and paint formulations, can save money and improve product properties.
- 5) Oil well drilling and well completions – gilsonite has long been used to improve characteristics of water-based, oil-based, and synthetic-based drilling mud systems.

Gilsonite is also used in low-density cementing slurry for lost circulation control. More details on these applications are provided by American Gilsonite Company (2008) and are summarized by Tripp and White (2006).

STRUCTURAL SETTING

Gilsonite veins are hosted in gently dipping Tertiary sedimentary rocks on the southern flank of the Uinta Basin. The basin is an asymmetric, intermontane basin along the northern edge of the Colorado Plateau. The Uinta Basin is bordered by the Uinta Mountains to the north, Douglas Creek arch to the east, Uncompahgre uplift to the southeast, San Rafael Swell to the southwest, and Wasatch Range to the west (figure 4). The closely related Piceance Creek Basin to the east is separated from the Uinta Basin by the north-south trending Douglas Creek arch. Structural features in the Uinta Basin and Uinta Mountains region have a development history that is long and complex, with some structures like the Uinta rift basin forming during Proterozoic time, and other structural activity possibly occurring during the Pennsylvanian-Permian ancestral Rocky Mountains uplift (Stone, 1993).

The Uinta Basin, Uinta Mountains, and associated folds were formed by west-southwest to east-northeast compression during the Cretaceous-early Tertiary Laramide orogeny (Erslev, 1993; Stone, 1993). Some fault systems that may have been formed during or were reactivated by the Laramide orogeny, were subsequently reactivated during the late Tertiary by extensional deformation producing normal-slip displacement.

Structural features within the study area include folds, faults, joints, and dikes, and are influenced by the regional structural trends exhibited throughout the Uinta Basin and Uinta Mountains. Faulting in the study area is minor in both abundance and displacement. The eastern part of the east-west-striking, normal-slip, Duchesne fault system (plate 1) occurs in the northwestern part of the study area. Other significant fault systems in the study area exist in the southwestern part trending northwest




Figure 4. Tertiary basin and uplift structures, and physiographic features in relation to study area (gray box)(modified from Cashion, 1967).

and in the southeastern part, near the Utah-Colorado border, trending northeast. Two gilsonite veins in the Pariette Draw area, the O.K. and Original Owner, apparently postdate movement on the Duchesne fault. The O.K. vein is along the fault and trends east-west. The Original Owner vein is approximately 0.3 mile south of the O.K. and also trends east-west. Also, the Florence vein has an unusual "kink" that may be related to faulting in the area.

In addition to the fractures that host the gilsonite veins, two other significant regional fracture/joint sets are common in the Uinta Basin; they trend northwest and northeast (Verbeek and Grout, 1993). According to Verbeek and Grout (1993), the older, northwest-trending set of joints dominates the fracture network in many areas and developed during the strongest period of post-Laramide extensional deformation. These joints strike almost parallel to the gilsonite veins; however, they are not directly related to (and appear to postdate) the fractures containing gilsonite, as shown by abutting relations between vein walls and joint surfaces, and differences in orientation and physical characteristics. The younger, northeast-trending joints are most numerous where the older northwest-trending joints are least developed. The gradual regional shift of the relative abundance of one joint set over the other reflects the gradually changing stress gradients over time in the basin (Verbeek and Grout, 1993).

STRATIGRAPHIC SETTING

Gilsonite veins are hosted in the Eocene strata of the upper Wasatch, Green River, Uinta, and lower Duchesne River Formations (figure 5). The rocks comprising these formations were deposited in lacustrine and fluvial depositional environments and range in composition from

chemically precipitated carbonate rocks to clastic rocks, and include minor tuffaceous rocks. Contacts between these formations are gradational, having complex inter-tonguing relationships and abrupt facies changes that reflect fluctuating paleo-lake (Lake Uinta) levels.

The lowest stratigraphic unit containing gilsonite veins is the Eocene Renegade Tongue of the upper Wasatch Formation. The predominantly fluvial Renegade Tongue consists of massive, irregularly bedded, brown and gray sandstone, and red and gray shale and siltstone that were deposited around the edges of Lake Uinta (Cashion, 1967). The Renegade Tongue interfingers with the primarily lacustrine Douglas Creek Member of the lower Green River Formation. Gilsonite veins in the Renegade Tongue crop out on the eastern side of the Uinta Basin near the Utah-Colorado border. The Weaver vein (in Colorado it is known as the Colorado vein) is present in the Renegade Tongue inside Colorado and is reported to pinch out farther to the southeast in Colorado and downward into shale beds in the Renegade Tongue (Cashion, 1967). The Black Dragon vein is located approximately 12 miles south of the Weaver vein and approximately 1 mile west of the Utah-Colorado border where it pinches out downward into shale beds at the contact zone between the Renegade Tongue and Douglas Creek Member.

Eocene Green River Formation lacustrine deposits host gilsonite veins in the basal Douglas Creek and upper Parachute Creek Members; however, no gilsonite is known to cross-cut the Mahogany oil shale zone contained in the Parachute Creek Member. The Douglas Creek Member consists primarily of sandstone, siltstone, and shale; algal, oolitic, and ostracodal limestone; and a few oil shale beds (Cashion, 1967). Veins below the Mahogany oil shale zone are thickest in the sandstone beds of the Douglas Creek Member and interfingering Renegade Tongue of the Wasatch Formation.




Figure 5. Stratigraphic column of the Eocene section hosting gilsonite veins in the Uinta Basin (modified from Sprinkel, 2007, 2009).

The Parachute Creek Member overlies and interfingers with the Douglas Creek Member and consists primarily of marlstone, oil shale, siltstone, sandstone, and tuff (Cashion, 1967). Bedding within the Parachute Creek Member is thin to laminated, even, and laterally continuous. The Mahogany oil shale zone is a sequence of rich oil shale beds within the Parachute Creek Member. The Mahogany bed is a particularly rich and laterally extensive oil shale bed in the Mahogany zone; its lateral continuity makes it a good regional marker bed (plate 1). Petrographically the oil shale is dolomitic marlstone having a high content of organic matter in the form of kerogen. All but two (Weaver and Black Dragon) of the known significant gilsonite veins are located above the Mahogany oil shale zone. Gilsonite veins that are above or beneath the Parachute Creek Member branch into veinlets and pinch out within the thin-to-laminated marlstone and oil shale beds contained in this member (Cashion, 1967). The veins can thicken in the upper beds of the Parachute Creek Member near the contact with the Uinta Formation.

Gilsonite veins in the eastern part of the basin are known to achieve their greatest thickness in the lower Uinta Formation (Pruitt, 1961; Cashion, 1967). The Eocene Uinta Formation interfingers with the underlying Parachute Creek Member and consists of marginal lacustrine deposits in the lower part, mixed fluvial and marginal lacustrine deposits in the middle part, and entirely fluvial deposits in the upper part. Marginal lacustrine deposits in the lower part consist primarily of thick, laterally continuous, medium-bedded to massive sandstone containing interbedded siltstone and thin intervals of marlstone and tuff. Fluvial beds in the middle and upper parts are composed of channel-form sandstone, variegated mudstone, and minor conglomerate that tend to be laterally discontinuous and thinner than marginal lacustrine deposits down section. Gilsonite veins commonly split, become discontinuous, and/or pinch out in the mudstone-rich upper Uinta Formation.

Rocks of the Eocene lower Duchesne River Formation are the youngest strata to host gilsonite veins. Conformability between the Uinta Formation and overlying Duchesne River Formation varies depending on location. The Duchesne River Formation consists primarily of fluvial interbedded mudstone and weakly cemented channel-form sandstone (Cashion, 1967). In the area of Fort Duchesne, Utah, the Carbon and Raven veins are located in the Brennan Basin Member of the lower Duchesne River Formation (Sprinkel, 2007) and were the first veins to be mined in the Uinta Basin (Pruitt, 1961).

Gilsonite veins can maintain their widths over extensive vertical and horizontal distances; vein continuity is related to the stratigraphy and lithology of the host formations (Pruitt, 1961; Cashion, 1967). Gilsonite veins

are commonly widest in the thick, competent, laterally continuous sandstone beds of the lower Uinta Formation and also in similar beds at the top of the Green River Formation (figure 6). Continuing up section, veins tend to split and be less continuous in thinner bedded marlstone, thicker bedded mudstone and siltstone, and discontinuous channel sandstones in the middle and upper Uinta Formation. In the eastern part of the study area, gilsonite veins above the source beds are in many cases eroded down near their roots, and the remaining veins generally can be expected to split and thin with depth. Gilsonite veins exposed beneath the source beds attain their greatest thickness in the Douglas Creek Member and the interfingering Renegade Tongue and can be expected to thin both downward and upward from these units. In the central and northwestern parts of the study area, bedrock hosting gilsonite veins is generally less eroded, and the veins rooted in the Green River Formation oil shale have greater vertical extent and generally can be expected to widen with depth from exposures in the middle and upper Uinta Formation.




Figure 6. Cross section of a typical gilsonite vein (from Eldridge, 1901).

ORIGINS OF GILSONITE

Gilsonite deposits primarily form as long, essentially vertical dikes (veins) that predominantly trend north-west and can range in width from less than an inch to more than 20 feet. Gilsonite sills (figure 7) are occasionally associated with the gilsonite dikes. The continuity of the veins is impressive; they stretch in relatively long, straight ribbons across the hills of the Uinta Basin, with the longest vein system (Pride of the West–Rainbow–Black Dragon) extending more than 22 miles. The gilsonite veins are also vertically continuous, ranging from hundreds to more than 3000 feet, commonly having only small variations in width.

The source and mechanisms that may have formed the gilsonite vein deposits in the Uinta Basin have long been debated, and the theories are well summarized by Verbeek and Grout (1993). Laboratory and field evidence strongly suggests that the gilsonite was sourced from kerogen-rich oil shale beds contained in and around the Mahogany oil shale zone in the Parachute Creek Member of the Green River Formation. Mechanisms for propagating the fractures that contain gilsonite appear to involve elevated pore pressure within the hydrocarbon source beds of the Green River Formation and regional stress fields.

The processes that formed the gilsonite vein deposits in the Uinta Basin began during Eocene time when a large lake, Lake Uinta, developed within a basin formed during the Late Cretaceous–early Tertiary Laramide orogeny. Lake Uinta covered an area now containing the Uinta Basin in Utah and Piceance Creek Basin in Colorado, but varied in position, size, and water chemistry over its millions of years of existence. During middle Eocene time, the lake reached its maximum areal extent and greatest depth, extending open-lacustrine conditions over a wide region. The lake also appears to have been chemically stratified during this time, having a lower, stagnant saline layer enriched in hydrogen sulfide and devoid of free oxygen, as indicated by the preservation of organic matter and the common occurrence of pyrite in the lake-bottom sediments (Hunt and others, 1954). Large amounts of organic material (primarily algae) accumulated on the bottom of the central area of the lake, along with bedded to laminated sediments composed primarily of calcium, magnesium, and sodium carbonates precipitated out of the lake water. Later, the heat and pressure of burial changed the organic-rich sediment into the thick, kerogen-bearing oil shales of the middle and upper Green River Formation.

Hunt and others (1954) and Hunt (1963) summarized compelling evidence that supports the Mahogany oil shale



Figure 7. Gilsonite sill (right of rock hammer) on the Rainbow vein in the lower Uinta Formation in NW¼NE¼ section 25, T. 11 S., R. 24 E., SLBLM, Uintah County.

zone as the source of the gilsonite. Their work involved comparing physical and chemical characteristics of vein and tar sand hydrocarbons to that of soluble organic matter contained in associated rocks to identify similarities that would connect solid hydrocarbon deposits to their source rocks. With respect to gilsonite, numerous sites were sampled starting approximately 1000 feet above the base of the Green River Formation, and the organic matter was analyzed for refractive index, infrared spectra, liquid chromatography results, and elemental composition. Near Bonanza, Utah, a strong similarity was demonstrated between the oil shale in and around the Mahogany zone and the nearby Cowboy and Bonanza (Independent) gilsonite veins. Hatcher and others (1992) used gas chromatography and mass spectrometry to compare hopane, sterane, and carotenoid biomarker compositions of gilsonite, tar sand, oil, and Green River Formation oil shale in the Uinta Basin. Results of their work demonstrated a direct relationship between Green River Formation oil shale and gilsonite, tar seeps, Asphalt Ridge tar sand, and asphaltic crude oils.

Strong field evidence also supports the Mahogany oil shale zone as the source of the gilsonite. No gilsonite veins are known to cross-cut the Mahogany zone. The Black Dragon vein is located beneath the Mahogany zone and in Threemile Canyon can be traced as it thins directly below, and subsequently terminates in, the Mahogany oil shale. The Bonanza (Independent) vein is located above the Mahogany zone and is reported to split at its base into numerous veinlets as it encounters the rich oil shale of the Mahogany (Hunt and others, 1954).

Verbeek and Grout (1993) concluded that the fractures filled by gilsonite veins are large-scale hydraulic extension features that resulted from the overpressurization of pore-space fluid in the hydrocarbon-rich source beds in the Green River Formation. Early-stage, post-Laramide, regional tectonic extension is also believed to be a factor in fracture formation (Verbeek and Grout, 1993). Fractures were forcefully propagated first by formational water, which in some instances bleached wall rock, deposited limonite and calcite on vein walls, and deposited chlorite in adjacent sandstones. Subsequently, fractures were widened as viscous, liquid asphaltite was injected under high pressure and later solidified into gilsonite, probably through cooling and polymerization. The gilsonite contains authigenic quartz and barite and occasionally 1- to 3-millimeter vesicles (Monsen and Parnell, 1992b) that originally contained water or gas.

EXPLORATION TECHNIQUES

All of the veins mined today were discovered by early prospectors who located surface exposures. Many of

the veins have prominent surface expressions, appearing as black, straight bands irrespective of topography, and contrasting strongly with light-colored host rocks in a sparsely vegetated terrain. Sections of the veins are, however, frequently covered by alluvium and colluvium and are generally well exposed only on ridges and in bedrock drainages, disappearing under alluvial valleys and reappearing on adjacent ridges. Where covered, veins can usually be mapped by the presence of eroded gilsonite flakes or evidence of past mining such as headframes and shafts sunk through the colluvium. The veins are generally straight and continuous, and the simple regional geology makes it easy to extrapolate between good exposures. Gilsonite is lightweight and can wash down colluvium-covered slopes, so streaks of gilsonite float (figure 8) can be offset from actual vein locations by 20 feet or more (Tripp and White, 2006).




Figure 8. Gilsonite float eroding from the northwest segment of the Harrison vein in $S\frac{1}{2}SW\frac{1}{4}$ section 29, T. 10 S., R. 23 E., SLBLM, Uintah County. Notebook is 7.5 inches long.

In areas where alluvium or colluvium is relatively thick, the traditional exploration technique was to project the trend of a vein and then trench or sink shallow shafts to verify location and determine the extent and quality of the gilsonite. Current exploration practices include angle and vertical drilling of suspected vein extensions using a truck-mounted drill rig; such drilling is important because some past mines were abandoned because the vein pinched out at shallow depths. Before a mine is placed in operation, the shaft location typically is core drilled to various depths; additional angle drilling is generally completed at 400-foot intervals along the strike of the vein on both sides of the shaft. Typical drill intercepts are at depths of 100, 300, 600, and 800 feet. Data collected during exploration include vein width, wall-rock characteristics, and ore quality. Quality characteristics of gilsonite veins, particularly the melt point, viscosity, and ash content, can change rapidly along strike and/or with depth. This information is of paramount importance for mine planning and determining potential markets for the gilsonite (Tripp and White, 2006).

Geochemical and geophysical exploration methods for gilsonite have yielded some success, but none have become standard exploration procedures. Botbol (1961) collected soil samples along closely spaced transects across covered extensions of known veins and then concentrated and measured the gilsonite fraction in the soil. The resulting data showed that the gilsonite contents in closely spaced samples were high over the expected trends of the veins. Electrical resistivity surveys across known gilsonite veins produced sharp anomalies with peak values over the center of the veins and smaller auxiliary peaks symmetrically spaced on both sides of the veins (Hays and others, 1967). Boleneus (2007) used electrical resistivity imaging to determine the presence and extent of gilsonite at abandoned mine sites on the Cowboy vein and a proposed shaft site on the Cottonwood vein. Near the northwest end of the Cowboy vein, electrical resistivity imaging was used to estimate the reclamation needs at abandoned gilsonite mines by determining the extent of underground workings and the location of unmined gilsonite. On the southeast segment of the Cottonwood vein, electrical resistivity imaging was used to locate gilsonite at a proposed shaft site for underground mining by Lexco, which operated two shafts farther to the northwest. The Cottonwood vein survey located an approximately 30-inch-wide vein using an electrode spacing of 16.4 feet in the proposed shaft area where there was no sign of gilsonite at the surface, neither in outcrop nor evidenced from soil or rock outcrops overlying the vein (Boleneus, 2007). The relatively wide electrode spacing shows that this type of geophysical survey can be used for rapid reconnaissance of a wide area to explore for concealed gilsonite veins of minable width.

DEVELOPMENT

Mining History

Gilsonite has a long, colorful mining history, which includes early exploration and development in rugged, remote country by prospectors; development of uses and markets; competition for the resource; conflicts among Native Americans, developers, and the U.S. Government; passage of laws; consolidation of development companies; development of processing and transportation infrastructure; and competition for markets. While the history before 1938 is very interesting, we will focus on the history since then. Crawford (1957), Kretchman (1957), Remington (1959), Covington (1964), and Bender (1970) discussed the pre-1938 period of gilsonite discovery and development in detail, and their accounts have also been summarized by Tripp (2004), and Tripp and White (2006).

In 1938, The Gilson Asphaltum Company, which had acquired the dominant gilsonite position in the Uinta Basin, changed its name to Barber Asphalt Company (the predecessor of Barber Oil Company). Barber Oil Company sold part ownership in the gilsonite operation to Standard Oil of California (now ChevronTexaco Corp.) in 1948 (Lewis, 1994). The new, jointly owned company was named the American Gilsonite Company (AGC). The new company renewed efforts to make gasoline and high-purity electrode carbon from gilsonite (Kretchman, 1957). Refinery design problems were solved by 1954, but a lower cost method of transporting the gilsonite was needed to make the project economic. A 72-mile-long slurry pipeline was constructed on an abandoned railroad right-of-way to a new (in August 1957) petroleum refinery at Gilsonite, Colorado, built specifically to refine gilsonite (Kretchman, 1957). The intended market for gasoline from the refinery was western Colorado, and most of the electrode carbon was for sale to the growing aluminum smelting industry of the Pacific Northwest (Henderson, 1957). Use of gilsonite for refinery feedstock greatly expanded production. In 1957, AGC started mining gilsonite with water jet cutters to speed production and prevent gilsonite dust explosions; Kilborn (1964) described the hydraulic mining technique in detail. AGC discontinued refining gilsonite and sold the refinery in 1973. Mine production declined from about 360,000 tons per year to about 54,000 tons per year as a result. AGC re-emphasized non-fuel uses of gilsonite and within 10 years AGC's gilsonite production expanded to 91,000 tons per year (Jackson, 1985). Around 1981, AGC replaced their 30-year-old, 4500-tons-per-month mill with a new, \$6 million processing plant. The new mill was designed to process 9000 tons per month of ore, be more efficient, improve product quality, and meet environmental and safety regulations (Jackson, 1985). In January 1981,

Chevron bought out Barber Oil's share of the business (Jackson, 1985; Hawes, 1990). AGC has simultaneously mined ore from several mines on different veins to provide the various grades of gilsonite specified by customers. In 1983, AGC mined ore from 11 gilsonite mines near Bonanza (Shushan, 1983). In 1991, Chevron sold their gilsonite operation to Stratford Enterprises Company of Tulsa, Oklahoma (Lewis, 1994). AGC then became an independent, publicly held company until March 2008, when they were acquired by American Gilsonite Holding Company, a privately held portfolio company of Palladium Equity Partners III, L.P. (Business Wire, 2008). In 2007, AGC employed 91 people at its mines and Bonanza mill and office (U.S. Mine Safety and Health Administration, 2008).

Two smaller companies have also mined gilsonite in the recent past: Ziegler Chemical & Mineral Corporation (Ziegler) and Lexco, Inc. Ziegler was incorporated in 1944 as a New Jersey-based producer of pitches, asphalts, and resins. The company, then known as G.S. Ziegler and Company, purchased gilsonite from small independent gilsonite producers. Ziegler acquired its own gilsonite properties in 1952 from the Utah Gilsonite Company, which had been in operation since 1920. In

1953, Ziegler also purchased some of the property of the American Asphaltum Association (a company formed in 1902 by former stakeholders in The Gilson Asphaltum Company). Ziegler additionally acquired the assets of the Standard Gilsonite Company in 1962 and renamed the combined company the Ziegler Chemical & Mineral Corporation (Lewis, 1994; Ziegler, 2008). The consolidated operation's office and plant were at Little Bonanza, Utah. Ziegler typically has produced less ore than AGC, but in 1994 produced more than 50% of the gilsonite mined in the Uinta Basin (Ziegler, 2008). In 2007, an average of 15 Ziegler employees worked 26,619 hours at the mines and the Little Bonanza mill (U.S. Mine Safety and Health Administration, 2008).

Lexco, Inc. was relatively new to the gilsonite industry when it began operating in 1988. Lexco operated a processing plant southeast of Fort Duchesne, in central Uintah County, and produced gilsonite from the ITM and Cottonwood mines. As recently as 2008, Lexco operated the Cottonwood mine where the vein width averages 36 to 38 inches. At the ITM mine, the vein width averages 18 inches and was mined in sections as narrow as 14 inches. In 2007, an average of 19 Lexco employees worked 32,187 hours at the mine and mill (U.S. Mine Safety and Health




Figure 9. Gilsonite annual production (red dots) and value (yellow triangles). Data gaps occur where production data were not preserved or where data were withheld by U.S. Bureau of Mines. Data from U.S. Bureau of Mines; Utah Division of Oil, Gas and Mining; Utah Geological Survey; Aurand (1920); Baker (1950); Carey and Roberts (1949); Cashion (1969); Covington (1964); Crawford (1957); Dennis (1930); Eldridge (1901); Garvin (1966); Kretchman (1957); Ladoo and Myers (1951); Romney (1963); Utah Mining Association (1955, 1959, 1967).

Administration, 2008). Lexco was recently acquired by AGC and its mines and processing plant were idle as of 2010. In general, gilsonite production and value has increased gradually from 1885 to 2008 (figure 9) except for the production spike from 1957 to 1973 that was associated with operation of AGC's refinery.

Mining Methods and Transportation

Early gilsonite mining was predominantly by open-cut mining with picks, shovels, and horse-powered hoists. All the gilsonite produced today from the Uinta Basin is by underground mining methods (figure 10). Mining consists of two major phases: shafts are sunk at regular intervals along the veins, and drifts and slopes are then extended laterally from the shafts. The top 30 feet of the gilsonite is not mined for safety and reclamation reasons. The following mining method description was largely taken from Tripp and White (2006).

Mining begins with main shaft development either by hand sinking directly into the ore or by drilling the shaft with a Teton drill rig that has a large-diameter (7-foot) bit. The method selected depends on the width of the vein at the surface and at depth. Hand sinking is preferred if the vein is at least 2 feet wide at the surface because it

yields salable gilsonite. Shaft drilling contaminates the gilsonite with wall rock; however, it is sometimes necessary on veins that, on outcrop, are too narrow for hand mining, but that widen at depth. On the Cowboy vein, where outcrops are generally not very wide but vein width increases with depth, the large-diameter drill is used to develop a shaft down to the depth where the vein is at least 4 feet wide. Once this width has been verified, the shaft sinking continues by hand. Drilled sections of shafts are usually about 220 feet deep. Typically, shafts developed in the ore body extend 26 feet along the length of the vein. All of the shaft-support equipment, including landings and guide rails, are installed in conjunction with hand sinking. The shaft contains three compartments; the two outside compartments contain pipes for the air lift and compressed air, and a central compartment contains a skip for miners.

Once the main shaft has been developed to a depth where the ore is wide enough for mining, or is at least 30 feet below the surface, a drift is started in one or both directions from the shaft. These drifts extend 500 to 700 feet from the shaft and serve as the escapeway drifts for the initial phase of mining. Once the initial drifts are complete, a small escapeway shaft is raised to the surface or is drilled if the distance from the surface down to the




Figure 10. Gilsonite underground mining methods (from Tripp and White, 2006, courtesy of American Gilsonite Company).

escapeway drift is more than 50 feet. In cases where the ore width is insufficient to conduct shaft raising or sinking in the vein, an escapeway shaft is drilled to intersect the escapeway drift.

Recently, AGC changed its mining methods by sinking the main shaft to the bottom of the ore deposit and then developing the initial drifts at this lower level and extracting ore from the bottom of the mine up to the top of the mine. The major benefit of this method is reduced shaft and escapeway maintenance. Once the ore has been extracted from the lower levels, those portions of the drifts and shaft do not need to be maintained. If mining is conducted from the top of the mine to the bottom, all of the drifts and shafts must be maintained for the life of the mine.

Gilsonite slope mining is generally initiated by developing two drifts, 200 vertical feet apart, from the main shaft to the escapeway shafts. Development can take place along strike in both directions from the shaft. Once these drifts are completed, mining begins on the floor of the upper drift downward and a 45° slope is created. This slope is extended along the upper drift until the bottom of the slope reaches the lower drift. This creates a slope in the ore body, which is approximately 140 feet long. Mining progresses from the bottom of the slope, which is the lower drift, to the top of the slope, which is the upper drift. A miner using a chipping hammer cuts a 4-foot-high bench on the bottom of the slope. The miner then cuts this bench up the 45° incline to the top of the slope. As the ore is cut with the chipping hammer, it slides down the slope and is directed into the air lift that conveys the ore to the surface. In a 5-foot-wide vein, each bench mined up the slope yields 79 tons of gilsonite. These benches are mined until the slope reaches the escapeway shaft. The remaining triangle of ore is mined from the top to the bottom of the slope so that a straight face of ore is left.

The gilsonite ore usually is mined using pneumatic chipping hammers that weigh approximately 12 pounds and are equipped with 12-inch-long, hardened steel moils. Compressed air for the chipping hammers comes from a 150-horsepower air compressor on the surface via a 3-inch-diameter steel pipe down the shaft to the working face (Jackson, 1985). On occasion, permissible explosives will be used if the ore is particularly hard to mine.

Mine timbers placed on 5-foot centers provide ground support in gilsonite mines. The timbers used in the shaft and on working floors are "hitched" into holes approximately 2 inches deep and slightly larger than the diameter of the timber and are locked into these hitches with wooden wedges. With the support resting on the 2-inch-wide rock ledge of the hitch, the timber will remain in place regardless of shrinkage. Timbers that are only set to control the wall rock are not hitched, but are placed

on 2 x 10 x 12-inch cap boards on each end, then wedged tightly into place.

AGC conveys the mined ore from the shaft, drift, or slope by means of an air lift system. Each system is generally equipped with three centrifugal fans, each powered by a 101-horsepower electric motor. The Sprout Waldron centrifugal fans are arranged in series and together develop a vacuum of 140 inches of water gage. The fans are connected to a 12-inch-diameter, steel, air lift pipe which is connected to a bag house and bin arrangement. The air lift pipe extends from the bag house into the mine. Depth of workings and distance between shafts is partially controlled by the capacity of the air lift system. The air lift system can efficiently transport gilsonite in 1100-foot-deep shafts having 700-foot-long drifts, but performance drops off after these distances are exceeded. The air lift system performs three additional functions: (1) it ventilates the mine, (2) it breaks the ore into smaller fragments (gilsonite fractures easily during mining and transport), and (3) it removes the small amount of water that percolates into the mine. Occasionally, large inflows of water require pumping by 50-horsepower staged pumps. While the air lift system is expensive to operate, it is a versatile and proven technology.

Additional mechanization of gilsonite mining has been limited by (1) lack of equipment designed to operate in narrow veins, (2) the highly explosive nature of gilsonite dust, which complicates use of explosives and diesel-, gas-, and electric-powered equipment, and (3) the need to prevent contamination by wall rock fragments, which favors careful hand mining of the ore. Despite these constraints, AGC has experimented with several mechanical mining systems and has recently evaluated a hydraulically powered continuous mining machine designed for narrow veins.

HEALTH ISSUES

Gilsonite is friable and mining generates large quantities of very fine, dark-brown dust, which can be explosive, but which does not represent any special respiratory risk to workers. In the early days of gilsonite mining, ventilation was often inadequate, candles were used for illumination, and explosives were sometimes used, all of which resulted in occasional, spectacular mine explosions and fires. Modern mining techniques and regulations have eliminated most of the explosions and fires. The U.S. Occupational Safety and Health Administration's Material Safety Data Sheets classify gilsonite as non-toxic, noncarcinogenic, and nonmutagenic, so no extreme safety measures are necessary. However, a respirator is recommended in heavy dust concentrations. Gilsonite is also approved for use as a coating on surfaces that contact food (Ziegler, 2008).

GILSONITE VEINS

In our field study, 59 veins, vein systems, and isolated vein outcrops were mapped and described in detail. The total combined vein length mapped was more than 120 miles. The UGS work, combined with previous mapping, identified 71 significant veins, vein systems and vein extensions having a total combined vein length of more than 170 miles. Cashion (1967) estimated the original in-place gilsonite resource of all the veins to be approximately 45 million short tons. In the following sections each vein, vein system, vein extension, isolated vein outcrop, or field investigation is summarized from field data and/or previous work by other investigators. Gilsonite veins in a particular area are generally listed north to south. Significant gilsonite veins that were located during the field study, or confirmed in the literature, are shown on plate 1 (except the Carbon and Raven veins) and summarized in table 2. All township, range, and section locations in the following text are relative to the Salt Lake Base Line and Meridian (SLBLM) unless otherwise noted.

Bonanza Area Veins

Red Wash and Little Red Wash

The Red Wash and Little Red Wash veins (figure 11; plate 1) are closely related and are located in sections 18 and 19, T. 8 S., R. 23 E., and section 13, T. 8 S., R. 22 E. We mapped the vein system for approximately 1.9 miles, from a 7-inch-wide outcrop located approximately 1 mile northwest of Red Wash at the southeast end to a 7-inch-wide outcrop located a short distance southeast of the paved road on Glen Bench at the northwest end. The vein system generally strikes N. 65° W., having a maximum width of 1.1 feet on the south vein and a maximum width of 1 foot on the north vein. Both maximum widths are located a short distance northwest of the vein system's midpoint. The veins are hosted at the surface by the mudstone and minor sandstone belonging to Sprinkel's (2007) member C of the upper Uinta Formation (plate 1). The vein system consists of two main veins less than 200 feet apart that run subparallel, and also has thinner, closely spaced, subparallel, secondary veins cropping out in places. The southernmost vein is 1.9 miles long and generally wider than the north vein, which is 1.2 miles long. The north vein begins a short distance southeast of the vein system's midpoint and continues with the south vein to the northwest termination. Both veins are locally sinuous, and pinch and swell throughout their lengths. Bitumen-impregnated sandstone outcrops (figure 10) are present along the veins at various stratigraphic levels, and one relatively thick sandstone bed appears to contain the most significant bitumen deposits. Bitumen-impregnated sandstone outcrops are black adjacent to the veins, and quickly lighten away from the veins. The

black, bituminous sandstone is not widespread and is found only adjacent to the veins. The thicker sandstone unit that hosts the most significant bitumen-impregnated sandstone outcrops is, however, fairly gray and widespread but also bleached in places. The vein system is unmined and only one or two superficial prospects exist.

Cowboy (Eureka)

The Cowboy (Eureka) vein system (plate 1) contains the widest gilsonite vein in the Uinta Basin, as well as the largest single gilsonite deposit, and has supported mining operations since the late 1800s (Pruitt, 1961). Vein locations on plate 1 were derived from USGS mapping (Cashion, 1978, 1986) and our NAIP aerial-photography mapping. The vein system is located in sections 7, 8, 16, 17, 22, and 23, T. 9 S., R. 25 E., sections 1, 2, 3, 4, and 12, T. 9 S., R. 24 E., and sections 32 and 33, T. 8 S., R. 24 E. The vein system is reported to be more than 10 miles long, generally strikes from N. 60 to 70° W., has a maximum width of 22 feet, and has an estimated maximum vertical extent of 1000 feet (Pruitt, 1961; Cashion, 1967). At the southeast end, the vein system begins near the north wall of the White River canyon in the SW¼ section 23, approximately 1.3 miles west of the Utah-Colorado state line. In places, the southeast end of the vein appears to be eroded away where it intersects drainages that cut through the upper Green River Formation. Historical open-trench mining at the southeast end appears to have been shallow, which suggests the vein is splitting and pinching out as it cuts deeper into the upper Green River Formation. Nahcolite (NaHCO₃) dissolution cavities in the upper Green River Formation have also been filled with gilsonite at the southeast end of the vein. The Cowboy vein splits in places, having offshoots of significant width on the northwest segment. Pruitt (1961) reported that a 5- to 6-foot-wide vein called the "E.B." (extra black) vein is located parallel and approximately 100 feet to the north of the northwestern part of the Cowboy vein. Upper Green River Formation (Parachute Creek Member) and Uinta Formation sandstone and mudstone host the vein system at the surface (plate 1). The Cowboy vein system attains its greatest width in the massive sandstone in the lower part of the Uinta Formation. The base of the vein is about 150 feet above the Mahogany bed, and the upper termination is in thin mudstone and sandstone beds in the Uinta Formation (Cashion, 1967).

Independent (Bonanza), Tabor, and Little Bonanza

The Independent (Bonanza), Tabor, and Little Bonanza vein system (plate 1) contains the second-widest gilsonite vein in the Uinta Basin and has supported mining operations since the late 1800s (Pruitt, 1961). Vein locations on plate 1 were derived from USGS mapping

Table 2. Summary of gilsonite vein attribute data.

| Vein Name | Mapped Length (mi) | Max. Surface Width (ft) | General Strike | Surface Hosting Formation | Mining | Data Source |
|-----------------------------|--------------------|-------------------------|----------------|---------------------------|--------|-------------|
| BONANZA AREA | | | | | | |
| Red Wash | 1.9 | 1.1 | N65°W | Uinta | No | UGS |
| Little Red Wash | 1.2 | 1 | N65°W | Uinta | No | UGS |
| Cowboy | 10 | 22 | N60-70°W | Green River - Uinta | Yes | USGS |
| Independent-Tabor | 7.5 | 14 | N55-62°W | Green River - Uinta | Yes | USGS |
| Little Bonanza | 5.5 | 13 | N68°W | Uinta | Yes | USGS |
| Little Chepetta | 2.3 | 2.5 | N62°W | Uinta | ? | USGS |
| Chepetta | 3 | 1.5 | N65°W | Uinta | ? | USGS |
| Caldwell | 0.7 | ? | N60°W | Uinta | ? | USGS |
| Wagon Hound | 5 | 3.3 | N63°W | Uinta | Yes | UGS |
| Little Emma | 5 | 2.8 | N65°W | Uinta | Yes | UGS |
| Augustine | 0.9 | 2.2 | N50°W | Green River | ? | USGS |
| Weaver | 2.3 | 2 | N58°W | Lower Green River | Yes | UGS |
| RAINBOW AREA | | | | | | |
| Little Butte | 0.5 | 1.1 | N60°W | Uinta | No | UGS |
| Rustler | 0.8 | 1.2 | N55°W | Uinta | No | UGS |
| Alabama | 3.9 | 2 | N50-65°W | Uinta | No | UGS |
| Little Alabama | 0.2 | 0.2 | N55°W | Uinta | No | UGS |
| Black Dragon | 3.9 | 10 | N47°W | Lower Green River | Yes | UGS |
| Rainbow | 4.3 | 10 | N45-55°W | Green River - Uinta | Yes | UGS |
| Pride of the West | 13.8 | 7 | N50-65°W | Green River - Uinta | Yes | UGS |
| Harrison | 10.6 | 4 | N50-65°W | Green River - Uinta | Yes | UGS |
| Little Asphalt Wash | 0.8 | 2 | N50-65°W | Green River - Uinta | No | UGS |
| Asphalt Wash | 2.5 | 3.5 | N50-60°W | Green River - Uinta | No | UGS |
| Snow | 1.1 | 0.25 | N67°W | Uinta | No | UGS |
| Saddletree | 1.4 | 2.2 | N58°W | Uinta | No | UGS |
| South Harrison | 6.1 | 3.5 | N58°W | Uinta | No | UGS |
| Little Chief | 0.9 | 0.7 | N57°W | Uinta | No | UGS |
| Fox | 2.9 | 2.5 | N55-65°W | Uinta | No | UGS |
| Kings Well | 5.6 | 6 | N55-65°W | Green River - Uinta | Yes | UGS |
| Neal | 1.8 | 3 | N60°W | Uinta | Yes | UGS |
| Bitter Water | 7 | 3 | N58-70°W | Uinta | No | UGS |
| Long Draw | 0.6 | 2 | N59°W | Green River - Uinta | No | UGS |
| Sage Brush | 1.8 | 1 | N65°W | Green River - Uinta | No | UGS |
| Little Antler | 0.5 | 1 | N62°W | Green River - Uinta | No | UGS |
| Antler | 0.6 | 2 | N58°W | Green River - Uinta | No | UGS |
| Little Bitter Creek #1 | 0.3 | 0.5 | N56°W | Uinta | No | UGS |
| OURAY AREA | | | | | | |
| Ouray | 0.3 | 1.5 | N62°W | Uinta | Yes | UGS |
| Little Seam | 0.2 | 0.25 | N56°W | Uinta | No | UGS |
| Gem | 0.2 | ? | N46°W | Uinta | No | UGS |
| Little Boy | 0.5 | 2 | N45°W | Uinta | No | UGS |
| Cliff Dweller | 0.15 | 0.5 | N55°W | Uinta | No | UGS |
| Jumbo | 1.8 | 1.1 | N48-65°W | Uinta | No | UGS |
| Pride of Utah | 6 | 2 | N58°W | Uinta | Yes | BLM |
| Natural Buttes | 2.2 | 0.7 | N55°W | Uinta | No | BLM |
| Black Gnat | 2.4 | 2.5 | N54°W | Uinta | Yes | BLM |
| Workman | 0.9 | 1.2 | N53°W | Uinta | Yes | BLM |
| Willow #1 | 1.8 | 2.2 | N50-70°W | Uinta | No | UGS |
| Willow #2 | 1.4 | 1.5 | N63°W | Uinta | No | UGS |
| Willow #2 extension | 1.3 | 1.2 | N54°W | Uinta | No | BLM |
| Willow #3 | 0.5 | 1.8 | N68°W | Uinta | No | UGS |
| Willow #4 | 0.3 | 0.6 | N60°W | Uinta | No | UGS |
| Turtle Shell | 0.8 | 1 | N66°W | Uinta | No | UGS |
| Petroglyph | 1.2 | 1 | N56°W | Uinta | No | UGS |
| Antelope | 0.4 | 0.8 | N47°W | Uinta | No | UGS |
| Black Diamond | 0.4 | 1.4 | N56°W | Uinta | Yes | UGS |
| Turkey Trail Hill | 1 | 2 | N60°W | Uinta | ? | UGS |
| Willow Creek #1 | 3.3 | 1 | N50-58°W | Uinta | No | UGS |
| Willow Creek #2 | 1.2 | 1.8 | N54°W | Uinta | No | UGS |
| Willow Creek extension | 0.87 | 0.9 | N57°W | Uinta | No | BLM |
| Black Bridge | 2.2 | 2.5 | N57-80°W | Uinta | ? | BLM |
| Cottonwood | 7 | 4.5 | N60-70°W | Uinta | Yes | UGS |
| Black Cat | 0.5 | 1 | N60°W | Uinta | No | UGS |
| Willow Creek-Wall | 3 | 3.5 | N60°W | Uinta | Yes | UGS |
| Black Dog | 0.4 | 0.6 | N50°W | Uinta | No | UGS |
| O.K. | 0.7 | 0.4 | N80-90°W | Uinta | No | UGS |
| Florence | 0.8 | 1.8 | N47-80°W | Uinta | Yes | UGS |
| Original Owner | 0.25 | ? | N80-90°W | Uinta | No | UGS |
| PARIETTE AREA | | | | | | |
| Castle Peak-Baxter-Pariette | 5 | 2 | N30-40°W | Uinta | Yes | UGS |
| Gilsonite Draw | 0.25 | 2(?) | N35°W | Uinta | ? | UGS |
| FORT DUCHESNE AREA | | | | | | |
| Raven | 3 | 2 | N37°W | Duchesne River | Yes | UGS |
| Carbon | 3 | 4 | N40°W | Duchesne River | Yes | UGS |

Total vein length = 172.4

(Cashion, 1974, 1977, 1986) and our NAIP aerial-photography mapping. The vein system is located in sections 29, 30, 32, and 33, T. 9 S., R. 25 E., and sections 8, 15, 16, 17, 22, 23, 24, and 25, T. 9 S., R. 24 E. Upper Green River Formation (Parachute Creek Member) and Uinta Formation sandstone and mudstone host the Independent, Tabor, and Little Bonanza vein system at the surface (plate 1).

The Independent and Tabor veins are a single vein but are called by different names northwest and southeast, respectively, of the junction with the Little Bonanza vein in NW $\frac{1}{4}$ section 30. The Independent/Tabor vein is reported to be more than 7.5 miles long, generally strikes from N. 55 to 62° W., has a maximum width of 14 feet, and has an estimated maximum vertical extent of 1100 feet (Cashion, 1967). The southeast end of the Tabor vein segment begins north of the White River canyon in NW $\frac{1}{4}$ section 33, and contains a short split-off vein approximately 0.5 mile from the southeast end. Pruitt (1961) reported that the Independent vein segment is 6 feet wide near the junction with the Little Bonanza vein, reaches a width of 13.5 feet approximately 2 miles farther to the northwest, and then decreases to 7 or 8 feet wide 3.5 miles from the vein junction. The Little Bonanza vein is reported to be more than 5.5 miles long, generally strikes N. 68° W., has a maximum width of 13 feet and has an estimated maximum vertical extent of 1200

feet (Cashion, 1967). Pruitt (1961) reported that the Little Bonanza vein width averages 9 feet or more along its length and at the northwest end decreases to 4 feet and begins to split. All of the veins in the system are widest in the massive sandstone contained in the lower part of the Uinta Formation. The veins all pinch out down section in marlstone and oil shale above the Mahogany bed in the Parachute Creek Member, and the up-section pinch out is in thinly bedded sandstone and mudstone of the Uinta Formation (Cashion, 1967).

Little Chepetta, Chepetta, and Caldwell

The Little Chepetta, Chepetta, and Caldwell veins and/or systems (plate 1) are all within a mile southwest of the Independent, Tabor, and Little Bonanza vein system. Vein locations on plate 1 were derived from USGS mapping (Cashion, 1974, 1977, 1986) and our NAIP aerial-photography mapping. The veins are located in sections 29, 30, 31, and 32, T. 9 S., R. 25 E., and sections 25 and 26, T. 9 S., R. 24 E. Uinta Formation sandstone and mudstone primarily host the Little Chepetta, Chepetta, and Caldwell veins at the surface (plate 1).

The Little Chepetta vein system is located about 1000 feet southwest of the Tabor vein, extends approximately



Figure 11. Red Wash vein and bitumen-impregnated sandstone (foreground) and Little Red Wash vein (background) cropping out in member C of the Uinta Formation in SE $\frac{1}{4}$ SE $\frac{1}{4}$ section 13, T. 8 S., R. 22 E., SLBLM, Uintah County.

2.3 miles along a strike of N. 62° W. (Cashion, 1974, 1986), and ranges in width from 1 to 2.5 feet over a length of 1.5 miles (Pruitt, 1961). A short subparallel vein is present a small distance north of the southeast segment of the Little Chepetta vein and may be related to it. To the northwest, the Little Chepetta vein trace disappears for a short distance and then reappears for approximately a third of a mile. The Chepetta vein system is located south of the Little Chepetta vein system and is approximately 3 miles long, generally strikes N. 65° W. (Cashion, 1974, 1986), and ranges in width from 1 to 1.5 feet over its length (Pruitt, 1961). A short subparallel vein is present a small distance north of the southeast segment of the Chepetta vein and may be related to it. The Caldwell vein is located south of the Chepetta vein system and is approximately 0.7 mile long, generally striking N. 60° W. (Cashion, 1986). Little else is known about the Caldwell vein.

Wagon Hound

The Wagon Hound vein (plate 1) is located in section 6, T. 10 S., R. 25 E., section 1, T. 10 S., R. 24 E., and sections 27, 28, 34, 35, and 36, T. 9 S., R. 24 E. We mapped the vein for approximately 5 miles, from a 1.5-foot-wide outcrop half a mile west of the White River canyon at the southeast end to a 1.4-foot-wide outcrop at the northwest end. The vein generally strikes N. 63° W. and has a maximum width of 3.3 feet, as exposed in a prospect trench 0.7 mile southeast of State Route 45. Cashion (1967) gave an estimated vertical extent of the vein of 1300 feet. Uinta Formation sandstone and mudstone host the vein at the surface; however, at the southeast end the vein terminates in beds at the top of the Parachute Creek Member of the Green River Formation (plate 1). The Wagon Hound vein consists of one main vein ranging in width from 2 to 3 feet along most of its length. The vein walls are commonly stained by iron oxide and, in places, are bitumen-impregnated. Sandstone beds hosting the vein locally contain hematite concretions, which range in size and occur within the bitumen-impregnated and iron-oxide-stained vein walls. The vein contains prospects along its entire length and appears to have been mined, via shafts and trenches, starting west of State Route 45. In 2004, the vein hosted one active mine shaft near its northwest end in NE¼SE¼ section 28, as well as several other modern, but inactive, mine shafts.

Little Emma (Uinta)

The Little Emma (Uinta) vein (plate 1) is located in sections 2 and 3, T. 10 S., R. 24 E., sections 29, 30, 32, 33, and 34, T. 9 S., R. 24 E., and section 25, T. 9 S., R. 23 E. We mapped the vein for approximately 5 miles, from an occurrence of gilsonite float chips a short distance west of the White River bridge on State Route 45 at the southeast end to an escape shaft at the northwest end.

The vein generally strikes N. 65° W. and has a maximum width of 2.8 feet, as exposed in a shaft at the middle of the northwest segment of the vein. Cashion (1967) gave an estimated vertical extent of the vein of 1000 feet. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The Little Emma vein consists of one main vein having an average width of 2 feet along most of its length. The vein walls are commonly stained by iron oxide and, in places, sandstone beds contain concretions. The vein contains prospects along its entire length and appears to have been mined via shafts, both modern and historical, especially the northwest segment where numerous shafts are present.

Augustine

A gilsonite vein is reported to be located approximately 1 mile north of the Weaver vein, in sections 2, 3, and 11, T. 10 S., R. 25 E. (Cashion, 1977). The area is located on the crest of a high ridge accessed by a rough, four-wheel-drive road. A brief field reconnaissance of the area resulted in the discovery of only a few small gilsonite chips as float in SE¼SE¼ section 3. A gilsonite vein probably exists in the area; however, without trenching we could not determine its exact location. A geological map of the Weaver Ridge quadrangle (Cashion, 1977) includes a vein named the Augustine (plate 1), which is mapped in the area where the float was discovered. Cashion (1977) reported the vein to be approximately 0.9 mile long, striking N. 50° W., more than 2 feet wide at its northwest end, and exposed at an elevation a few hundred feet above the Mahogany oil shale zone in the area.

Weaver

The Weaver vein (plate 1) straddles the Utah and Colorado state line (in Colorado it is referred to as the Colorado vein), and in Utah the vein is located in sections 10, 13, and 14, T. 10 S., R. 25 E. We mapped the generally N. 58° W.-striking vein for approximately 2.3 miles, from an outcrop 1.2 feet wide a short distance inside Colorado, to a short 2-foot-wide prospect trench (the vein's maximum known width in Utah) on the south side of Weaver Canyon at the northwest end. Cashion (1967) gave an estimated vertical extent of the vein of 1400 feet. The lower part of the Parachute Creek Member and the complete Douglas Creek Member of the Green River Formation host the vein at the surface in Utah and into Colorado (plate 1). In Colorado, the vein continues southeast for about another 4 miles before eventually pinching out in Wasatch Formation shale beds (Cashion, 1967). To the northwest in Weaver Canyon (SE¼ section 10), we observed the vein to abruptly terminate under thick beds of marlstone and shale in the lower Parachute Creek Member, approximately 300 feet beneath the Mahogany oil shale zone. No signs of significant mining were observed on the Weaver vein in Utah, but there are sporadic prospects along its

length. However, in Colorado it was one of the first veins to be mined in the basin around 1890 (Pruitt, 1961).

Rainbow Area Veins

Little Butte

The Little Butte vein (plate 1) is located in NW $\frac{1}{4}$ section 26 and NE $\frac{1}{4}$ section 27, T. 10 S., R. 24 E. We mapped the vein for approximately 0.5 mile, from a 1-foot-wide outcrop at the southeast end to another 1-foot-wide outcrop at the northwest end. The vein generally strikes N. 60° W. and averages 1 foot wide along its length. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein appears to be present only in the upper part of a dissected butte. An attempt to follow the vein a mile to the northwest at a lower elevation was unsuccessful; the vein also does not crop out at lower elevations to the southeast. The White River oil shale mine is at the northwest end of the vein and is private land, so the northwest terminus could not be located. We observed the vein continuing across the top of the butte into the mine area. The vein shows no signs of mining or prospecting.

Rustler

The Rustler vein (plate 1) is located in SW $\frac{1}{4}$ SW $\frac{1}{4}$ section 2, SE $\frac{1}{4}$ section 3, and NW $\frac{1}{4}$ section 11, T. 11 S., R. 24 E. We mapped the vein for 0.8 mile, from an occurrence of gilsonite float chips at the southeast end to an 8-inch-wide outcrop at the northwest end. The vein generally strikes N. 55° W. and has a maximum width of 1.2 feet at an outcrop near the vein's midpoint. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein may continue to the southeast, concealed by colluvium, but a fairly extensive search over a distance of approximately 1 mile to the southeast located only a few small gilsonite chips as float in dry washes. The Rustler vein is in a remote location having no easy access; one rough, four-wheel-drive road crosses the vein. The vein shows no signs of mining or prospecting.

Alabama

The Alabama vein (plate 1) is located in sections 35 and 36, T. 10 S., R. 23 E., and sections 4, 5, 6, and 9, T. 11 S., R. 24 E. We mapped the vein for approximately 3.9 miles, from a fracture zone containing gilsonite veinlets at the southeast end to an outcrop 2.5 inches wide at the northwest end. The vein generally strikes from N. 50 to 65° W. and has a maximum width of 2 feet, as exposed in a prospect trench near the vein's midpoint. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein consists primarily of one continuous vein between 0.5 and 2 feet wide having very few splits

or secondary veins. The vein appears to step to the north near the midpoint, where the vein is the widest. The vein averages 1.8 feet wide over an approximately 2-mile segment, which begins at the first observed outcrop south-east of Asphalt Wash and continues southeast until a jeep trail crosses the vein in NW $\frac{1}{4}$ NE $\frac{1}{4}$ section 9. The vein eventually thins to the southeast and northwest and subsequently disappears. No signs of significant mining were observed on the Alabama vein. However, prospect trenches are located on the widest section of the vein, beginning a short distance northwest of the vein's midpoint, and continuing for approximately 1.7 miles to the southeast.

Little Alabama

The Little Alabama vein (plate 1) is located in NW $\frac{1}{4}$ SW $\frac{1}{4}$ section 13, T. 11 S., R. 24 E., and is about 2.5 miles south-east of, but approximately on trend with, the main Alabama vein. The vein strikes N. 55° W., was mapped for 0.2 mile, and has a maximum width of 2 inches. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). This vein is difficult to trace to the southeast or the northwest because it is mostly covered by alluvium and colluvium, and is extremely thin.

Black Dragon

The Black Dragon vein system (plate 1) is located in sections 4, 5, 9, 10, 11, and 14, T. 12 S., R. 25 E. We mapped the vein system for approximately 3.9 miles, from an occurrence of gilsonite float chips on the top of a high ridge east of Dragon Canyon at the southeast end, to a fenced, caved-in shaft or subsidence in a drainage 0.7 mile northwest of Threemile Canyon. The vein system generally strikes N. 47° W. and has a maximum width from 8 to 10 feet, as estimated at a significant mine site in Dragon Canyon (figure 12). Cashion (1967) gave an estimated maximum vertical extent of the vein of 1100 feet, extending down from the lower part of the Parachute Creek Member through the Douglas Creek Member of the Green River Formation (plate 1). We observed the vein's upper extent to abruptly thin and terminate beneath the Mahogany oil shale marker bed in Threemile Canyon. Between Dragon and Threemile Canyons, the main vein splits in many places into two closely spaced, parallel veins of significant width. The vein system pinches out to the northwest as it cuts up section into Parachute Creek Member oil shale beds, and disappears to the southeast in the Douglas Creek Member and interfingering Renegade Tongue of the Wasatch Formation. Signs of significant historical mining and prospecting activity are present along most of the Black Dragon vein system.

Rainbow

The Rainbow vein system (plate 1) is located in sections

30, 31, and 32, T. 11 S., R. 25 E., and sections 14, 15, 23, 24, and 25, T. 11 S., R. 24 E. We mapped the vein system for approximately 4.3 miles, from an isolated group of veins that crop out for approximately 0.5 mile in upper Green River Formation marlstone at the southeast end to a deep mine trench in the East Fork of Asphalt Wash, where the division between the Rainbow and the Pride of the West extensions is located in section 15. The vein system generally strikes from N. 45 to 55° W. and has a maximum width on a single vein from 8 to 10 feet. Cashion (1967) gave an estimated maximum vertical extent of the vein of 600 feet. Upper Green River Formation (Parachute Creek Member) and lower Uinta Formation sandstone and mudstone host the veins at the surface (plate 1). The Rainbow vein splits in many places and consists of a group of closely spaced, parallel veins of significant width, especially in the area of the Green River and Uinta For-

mation contact. Northwest of the isolated group of veins at the southeast end, approximately 0.7 mile of the vein appears to have been eroded away, exposing Mahogany oil shale beds. The southeast end of the Rainbow vein system is less than 1 mile from the northwest end of the Black Dragon vein system, and lies several hundred feet to the southwest of the projected Black Dragon vein system. Signs of significant historical mining (figure 13) and prospecting activity are present along most of the Rainbow vein system.

Pride of the West

The Pride of the West vein system (plate 1) is located in sections 6, 7, 8, 9, 15, and 16, T. 11 S., R. 24 E., sections 1 and 2, T. 11 S., R. 23 E., sections 19, 28, 29, 30, 33, 34, and 35, T. 10 S., R. 23 E., and sections 14, 15, 23, and 24, T. 10



Figure 12. Black Dragon vein cropping out in the Douglas Creek Member of the Green River Formation in Dragon Canyon in SW $\frac{1}{4}$ NE $\frac{1}{4}$ section 14, T. 12 S., R. 25 E., SLBLM, Uintah County.



Figure 13. Mined open cut on the Rainbow vein in NE $\frac{1}{4}$ SE $\frac{1}{4}$ section 23, T. 11 S., R. 24 E., SLBLM, Uintah County.

S., R. 22 E. We mapped the vein system for approximately 13.8 miles, from the division between the Rainbow and Pride of the West extensions at the southeast end to a 4-inch-wide outcrop at the northwest end. The vein system generally strikes from N. 50 to 65° W. and has a maximum width on a single vein of 7 feet near the southeast end of the vein. Upper Green River Formation (Parachute Creek Member) and Uinta Formation sandstone and mudstone host the veins at the surface (plate 1). The Pride of the West vein splits in many places, and at the southeast end consists of a group of closely spaced, parallel veins of significant width. From Asphalt Wash toward the northwest end, one main vein is present along with occasional thinner, secondary veins. The vein eventually thins to the northwest and disappears approximately 1 mile northwest of Bitter Creek. Signs of significant historical mining and prospecting activity are present along most of the southeast segment of the vein system, and prospects also sporadically continue for the rest of the system's length.

Harrison

The Harrison vein (plate 1) is located in sections 7, 8, 16, 17, 21, 22, 23, and 26, T. 11 S., R. 24 E., sections 1, 2, 3, and 12, T. 11 S., R. 23 E., and sections 29, 30, 32, and 33, T. 10 S., R. 23 E. We mapped the vein for approximately 10.6 miles, from a 1-foot-wide outcrop approximately 1 mile southwest of the historic Rainbow town site at the southeast end to a thin outcrop approximately 1.2 miles northwest of Saddletree Draw at the northwest end. The vein generally strikes from N. 50 to 65° W. and has a maximum width estimated at 4 feet in a deep mine trench approximately 0.8 mile southeast of a prospect in the West Fork of Asphalt Wash. Upper Green River Formation (Parachute Creek Member) and Uinta Formation sandstone and mudstone (figure 14) host the vein at the surface (plate 1). The Harrison vein consists primarily of one main vein that varies from 1 to 4 feet wide along most of its length. However, in places the main vein splits into thinner, closely spaced, parallel veins, especially toward the ends of the southeast and northwest segments. Approximately 0.7 mile northwest of Atchees Wash, the vein disappears. It reappears about 0.4 mile to the northwest on a ridge top, having stepped approximately 600 feet north of the previous trend. The vein eventually thins to the northwest and disappears approximately 1.2 miles northwest of Saddletree Draw. The vein contains prospects along most of its length. Significant historical mining and prospecting activity have occurred on the widest segments of the vein in SW $\frac{1}{4}$ SW $\frac{1}{4}$ section 8, SE $\frac{1}{4}$ section 16, and section 22.

Little Asphalt Wash

The Little Asphalt Wash vein (plate 1) is located in sections 22 and 27, T. 11 S., R. 24 E., and is approximately 900 feet northeast of, and runs subparallel to, the

Asphalt Wash vein. We mapped the vein for 0.8 mile, from a sparse occurrence of gilsonite float chips at the southeast end to an occurrence of small float chips at the northwest end. The vein generally strikes from N. 50 to 65° W. and has a maximum width estimated at 2 feet on an outcrop approximately 700 feet northwest of an improved dirt road in NE $\frac{1}{4}$ section 27. Upper Green River Formation (Parachute Creek Member) and lower Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein is mainly a single one but in places splays into veinlets cropping out in fracture zones. The Little Asphalt Wash vein also appears to be rather short and quickly disappears in both directions. The vein shows no signs of mining or prospecting.

Asphalt Wash

The Asphalt Wash vein system (plate 1) is located in sections 22, 26, 27, 35, and 36, T. 11 S., R. 24 E. This vein




Figure 14. Harrison vein cropping out in the Uinta Formation in the West Fork of Asphalt Wash in NE $\frac{1}{4}$ SW $\frac{1}{4}$ section 7, T. 11 S., R. 24 E., SLBLM, Uintah County.

was previously known as the South Harrison and was renamed Asphalt Wash to avoid confusion with another more extensive South Harrison vein, which is located west of and not on trend with this vein. We mapped the vein system for approximately 2.5 miles, from a gilsonite veinlet outcrop approximately 700 feet northwest of an improved dirt road in NW¼ section 36 at the southeast end to a 4-inch-wide outcrop east of the Center Fork of Asphalt Wash at the northwest end. The vein system generally strikes from N. 50 to 60° W. and has a maximum width of 3.5 feet at the middle of the southeast segment of the southern vein. Upper Green River Formation (Parachute Creek Member) and lower Uinta Formation sandstone and mudstone host the veins at the surface (plate 1). The vein system consists of two, closely spaced, parallel veins of significant width. The southern vein is longer and generally wider than the northern vein, which crops out in the middle section of the vein system. The primary southern vein maintains a fairly consistent width of 2 to 3 feet throughout most of its length, and also contains a short split-off vein at the southeast end. Keighin (1977) reported that the vein continues to the northwest across the Center Fork of Asphalt Wash and beyond; however, we were unable to locate that extension. The vein appears to terminate to the southeast in the top of the Parachute Creek Member, where sub-inch gilsonite sills were observed in the shale beds. The vein system has been prospected sporadically along its length, but we observed no signs of significant mining.

Snow

The Snow vein (plate 1) is located in sections 2, 3, and 11, T. 11 S., R. 23 E. We mapped the vein for approximately 1.1 miles, from a 1-inch-wide outcrop at the southeast end to an occurrence of large gilsonite float chips at the northwest end in Atchees Wash. The snow vein generally strikes N. 67° W. and has a maximum width of 3 inches measured at several locations along the vein. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein is poorly exposed and is mostly traceable by sporadic gilsonite float and vein occurrences, but appears to be thin, having at least one other thin vein occurring a short distance to the north. The vein shows no signs of mining or prospecting.

Saddletree

The Saddletree vein (plate 1) is located in sections 4 and 5, T. 11 S., R. 23 E., and SE¼ section 31, T. 10 S., R. 23 E. We mapped the vein for approximately 1.4 miles, from a 1-foot-wide outcrop at the southeast end to an occurrence of small gilsonite float chips at the northwest end. The vein generally strikes N. 58° W. and has a maximum width of 2.2 feet at the middle of the vein, a short distance north of the improved dirt road in the West Fork of Saddletree Draw. Uinta Formation sandstone and mudstone

host the vein at the surface (plate 1). Thin secondary veins and veinlets are also present. The southeast segment of the vein is fairly wide, varying between 1.4 and 2.2 feet. The vein appears to be short and quickly disappears to the northwest and southeast. No mines and only one possible prospect are present on the vein.

South Harrison

The South Harrison vein system (plate 1) is located in sections 18, 19, and 20, T. 11 S., R. 24 E., and sections 3, 4, 10, 11, 13, and 14, T. 11 S., R. 23 E. We mapped the vein system for approximately 6.1 miles, from a 1.8-foot-wide outcrop west of the Center Fork of Asphalt Wash at the southeast end to a 1.1-foot-wide outcrop a short distance east of the West Fork of Saddletree Draw at the northwest end. The vein system generally strikes N. 58° W. and has a maximum width of 3.5 feet, as exposed in a prospect between Atchees Wash and the West Fork of Saddletree Draw. Uinta Formation sandstone and mudstone primarily host the veins at the surface (plate 1). The South Harrison vein splits in many places and consists of a group of closely spaced, parallel veins, some of significant width, while others are thin secondary veins. The consistently widest section of vein is between 2 and 3.5 feet and is located in the middle of the northwest segment of the vein system. At the northwest end, the vein thins and subsequently disappears; however, at the southeast end the vein is still 1.8 feet thick where it disappears under colluvium in flatter terrain. The vein system has been prospected along most of its length, but we observed no signs of significant mining.

Little Chief

The Little Chief vein (plate 1) is located in sections 14 and 15, T. 11 S., R. 23 E. We mapped the vein for approximately 0.9 mile, from an 8-inch-wide outcrop at the southeast end to an occurrence of small gilsonite float chips a short distance east of Atchees Wash at the northwest end. The vein generally strikes N. 57° W. and has a maximum width of 8 inches exposed on the rim of a high mesa overlooking the West Fork of Asphalt Wash. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The southeast end of the vein is mostly covered by colluvium and difficult to trace. Gilsonite is fairly well exposed at the northwest end, where a thin secondary vein is also present. The vein appears to be quite short and quickly disappears in both directions. The vein shows no signs of mining or prospecting.

Fox

The Fox vein system (plate 1) is located in sections 14, 15, 16, 23, and 24, T. 11 S., R. 23 E. We mapped the vein system for approximately 2.9 miles, from a 1.3-foot-wide outcrop a short distance west of the West Fork of Asphalt

Wash at the southeast end to a 6-inch-wide outcrop a short distance east of the West Fork of Saddletree Draw at the northwest end. The vein system generally strikes from N. 55 to 65° W. and has a maximum width of 2.5 feet measured on an outcrop approximately 0.5 mile from the southeast end of the vein system. Uinta Formation sandstone and mudstone primarily host the veins at the surface (plate 1). The Fox vein splits in many places, especially east of Atchees Wash where it is composed of one main vein and several thinner, closely spaced, parallel, secondary veins. The main vein varies between 1 and 2 feet wide along most of its length until it thins and disappears to the northwest, but remains 1.3 feet wide until it disappears to the southeast. No signs of mining were observed, but sporadic prospects are found along the length of the main vein.

Kings Well

The Kings Well vein (plate 1) is located in section 4, T. 12 S., R. 24 E., sections 30, 32, and 33, T. 11 S., R. 24 E., and sections 22 and 23, T. 11 S., R. 23 E. We mapped the vein for approximately 5.6 miles, from a 1-foot-wide outcrop a short distance east of Kings Well at the southeast end to an occurrence of small gilsonite float chips a short distance east of Atchees Wash at the northwest end. The vein generally strikes from N. 55 to 65° W. and has a maximum width of 6 feet, as measured in a short open cut leading to an adit. The vein at this location (approximately 0.3 mile northwest of Kings Well) is anomalously thick compared to vein outcrops short distances in either direction that are barely half as wide. We also observed no evidence of significant mining activity. Upper Green River Formation (Parachute Creek Member) and Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The Kings Well vein locally splits into one main vein along with thinner, closely spaced, parallel, secondary veins. Approximately 1.8 miles northwest of Kings Well, the vein steps north approximately 500 feet. A few hundred feet farther to the northwest of this step, a new 2-foot-wide vein appears a short distance north of the main vein location, and crops out for about 300 feet. The main vein is traceable for 1 mile northwest of this location, and then it disappears for approximately 1.5 miles across the West Fork of Asphalt Wash, and finally reappears approaching a steep mesa slope. The vein eventually thins to the northwest and disappears just short of Atchees Wash. To the southeast it disappears in the Parachute Creek Member west of Long Draw. Prospects sporadically occur along the entire length of the vein, and two small historical mines are located in section 32, T. 11 S., R. 24 E. One is a 1.5-foot-wide mine trench that is several hundred feet long in SE $\frac{1}{4}$ NW $\frac{1}{4}$ section 32, and the other is a shaft in a short mine trench where a 3- to 4-foot-wide vein is exposed in NW $\frac{1}{4}$ NW $\frac{1}{4}$ section 32. The depth of mining at both sites appears to be shallow, only about 30 feet.

Neal

The Neal vein (plate 1) is located in sections 31 and 32, T. 11 S., R. 24 E. We mapped the vein for approximately 1.8 miles, from an occurrence of small gilsonite float chips about 0.7 mile west of Kings Well at the southeast end to a 1-foot-wide outcrop at the northwest end. The vein generally strikes N. 60° W. and has a maximum estimated width of 3 feet in a mine trench at the middle of the southeast segment of the vein. Lower Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The Neal vein consists of one main vein with some thinner, closely spaced, parallel, secondary veins cropping out in places. The vein steps north approximately 300 feet near the northwest end and continues for about 1200 more feet after this step before it thins and disappears. Prospects and mine trenches having shafts are present at the middle of the southeast segment of the vein in S $\frac{1}{2}$ section 32 where the vein is widest, averaging between 2.5 and 3 feet.

Bitter Water

The Bitter Water vein (plate 1) is located in sections 16, 17, 18, 21, 22, 26, and 27, T. 11 S., R. 23 E., and sections 11, 12, and 13, T. 11 S., R. 22 E. We mapped the vein for approximately 7 miles, from an occurrence of small gilsonite float chips a short distance west of the West Fork of Asphalt Wash at the southeast end to a 9-inch-wide outcrop a short distance east of Bitter Creek at the northwest end. The vein generally strikes from N. 58 to 70° W. and has a maximum width of 3 feet, as measured on an outcrop approximately 1.3 miles southeast of the northwest end of the vein. Uinta Formation sandstone and mudstone primarily host the vein at the surface (plate 1); however, both ends of the vein appear to be located in the uppermost beds of the Parachute Creek Member of the Green River Formation. The Bitter Water vein consists primarily of one main vein with thinner, closely spaced, parallel, secondary veins cropping out in places. At the southeast end, and especially at the northwest end of the vein, significant splitting produces vein groups that may represent the vein's termination. The vein is concealed under colluvium for 0.7 mile, from 1.3 miles northwest of the West Fork of Saddletree Draw to a steep drainage leading down to Bitter Creek, where the vein reappears. Few prospects and no mines are found along the vein; the prospects are present sporadically from 0.5 mile southeast of Atchees Wash to approximately 1 mile northwest of the West Fork of Saddletree Draw.

Long Draw

The Long Draw vein system (plate 1) is located in S $\frac{1}{2}$ section 10, T. 12 S., R. 24 E. We mapped the vein system for 0.6 mile, from an occurrence of gilsonite float chips at the southeast end to a 1.2-foot-wide outcrop at the northwest

end. The vein system generally strikes N. 59° W. and has a maximum width of 2 feet, as measured on the northernmost vein at the middle of the system. At the surface the veins are located in the contact zone between the Green River and Uinta Formations (plate 1). The Long Draw vein system consists of two significant veins and numerous veinlets; the northernmost vein appears to be the widest. Northwest-trending fractures in gray marlstone near the southeast end of the veins are locally filled with gilsonite veinlets. To the northwest, the veins are concealed by alluvium as they disappear down the middle of a drainage that leads to the Center Fork of Asphalt Wash. To the southeast, the veins disappear in thinly bedded marlstone of the upper Parachute Creek Member of the Green River Formation. The vein system shows no signs of mining or prospecting.

Sage Brush

The Sage Brush vein (plate 1) is located in NE¼ section 18, T. 12 S., R. 24 E., and section 12, T. 12 S., R. 23 E. We mapped the vein for approximately 1.8 miles, from a 3-inch-wide outcrop at the southeast end to a 1-foot-wide outcrop at the northwest end. The vein generally strikes N. 65° W. and has a maximum width of 1 foot on an outcrop at the northwest end. The vein is exposed in the contact zone between the Green River and Uinta Formations (plate 1). The vein consists of one main vein with thinner, closely spaced, parallel, secondary veins cropping out in places. The vein is generally poorly exposed and is especially difficult to trace in the southeast, where it is mostly concealed under colluvial and alluvial deposits. The northwest segment is better exposed, but disappears in alluvium a short distance to the northwest. The vein shows no signs of mining or prospecting.

Gilsonite float chip and veinlet occurrences roughly on trend to the southeast of this vein are present in SE¼NW¼ section 17, T. 12 S., R. 24 E., and may be related to the vein, extending it about another 0.7 mile. Another unreported gilsonite vein is located in SW¼NE¼ section 18, T. 12 S., R. 24 E., approximately 1300 feet south of the southeast end of the Sage Brush vein. The 4-inch-wide vein strikes N. 30° W. and is exposed for only a few feet in a sandstone outcrop. Approximately 0.8 mile to the southeast of this vein outcrop, the Little Antler vein is on trend with it and may be related.

Antler and Little Antler

The Antler and Little Antler veins (plate 1) are located in sections 17, 20, and 21, T. 12 S., R. 24 E. The veins are separated by approximately 500 feet and are subparallel. We mapped the Antler vein for 0.6 mile, from a 1.4-foot-wide outcrop at the southeast end to an occurrence of gilsonite float chips at the northwest end. The vein generally strikes N. 58° W. and has a maximum width esti-

mated at more than 2 feet on an outcrop a short distance southeast of an unimproved dirt road in NE¼NE¼ section 20. The Antler vein (figure 15) consists of one main vein with thinner, closely spaced, parallel, secondary veins cropping out in places. The vein maintains a width between 1 and 2 feet from a short distance northwest of where the unimproved dirt road intersects the vein to its southeast end.

We mapped the Little Antler vein for approximately 0.5 mile, from a 1-foot-wide outcrop at the southeast end (the vein's maximum known width) to an occurrence of gilsonite float chips at the northwest end. The vein generally strikes N. 62° W. and is generally poorly exposed,



Figure 15. Outcrop on the Antler vein showing the main 1.8-foot-wide vein and an adjacent thinner (1.2 feet), parallel, secondary vein cropping out on the right side, located west of the Center Fork of Asphalt Wash in NE¼NE¼ section 20, T. 12 S., R. 24 E., SLBLM, Uintah County. Notebook is 7.5 inches long.

traceable mainly by sporadic gilsonite float occurrences. The Antler and Little Antler veins are exposed in the contact zone between the Green River and Uinta Formations (plate 1). Both veins may continue beyond extents shown on plate 1 since they are difficult to trace due to colluvial cover. The mapped segments of the veins show no signs of mining or significant prospecting.

Little Bitter Creek #1

The Little Bitter Creek #1 vein (plate 1) is located in SW $\frac{1}{4}$ SW $\frac{1}{4}$ section 34, and possibly in SE $\frac{1}{4}$ section 33, T. 10 S., R. 22 E. We mapped one vein for a couple hundred feet up the side of a steep mesa on the west side of Bitter Creek. It has a width of 5 to 6 inches and a general strike of N. 56° W. on the steep slope. Aerial photographs suggest the vein continues a short distance farther to the northwest for a combined mapped length of approximately 0.3 mile. The vein did not appear to continue to the southeast on the east side of Bitter Creek. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). A 2- to 3-inch-wide vein is present approximately 200 feet to the north of the first one. Bitumen-impregnated sandstone occurs adjacent to both veins. The veins at this location appear to be thin, short, and show no signs of mining or prospecting. Other veins of small dimensions are likely present in northwest-trending fractures in the Bitter Creek drainage, in the area between the northwest ends of the Bitter Water and Pride of the West veins.

Ouray Area Veins

Jumbo

The Jumbo vein (plate 1) is located in sections 25, 26, and 36, T. 9 S., R. 21 E. We mapped the vein for 1.8 miles, from an occurrence of gilsonite float chips at the southeast end to an approximately 6-inch-wide outcrop a short distance south of the Uintah and Ouray Indian Reservation at the northwest end. The vein generally strikes N. 48° W. and has a maximum width of 1.1 feet, as measured at a prospect (figure 16) in the middle of the northwest segment of the vein. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein is poorly exposed and is mostly traceable by sporadic gilsonite float and vein occurrences, but appears to consist of one main vein having closely spaced, subparallel veinlets cropping out in places. Near the southeast end of the vein, the general strike abruptly changes to N. 65° W. and the vein appears to shift significantly eastward. The vein may continue to the northwest on the Uintah and Ouray Indian Reservation, but thins before disappearing to the southeast. The vein has been prospected sporadically along its length but shows no signs of mining.

Pruitt (1961) reported that south of Ouray, Utah, on the Uintah and Ouray Indian Reservation, the Jumbo vein lies

2 miles northeast of the Pride of Utah vein in the west half of T. 9 S., R. 21 E., and the east half of T. 9 S., R. 20 E. The report states that the vein is 5 miles long, strikes N. 67° W., and is 1 to 1.2 feet wide near the northwest end. According to Pruitt (1961) and unpublished BLM gilsonite maps from the 1980s, several short veins also exist between the Jumbo and Pride of Utah veins.

Pride of Utah

The UGS-mapped segment of the Pride of Utah vein system (plate 1) is located in N $\frac{1}{2}$ section 32, T. 9 S., R. 21 E. We mapped the vein for 0.6 mile, from an occurrence of gilsonite float chips a short distance west of Cottonwood Wash at the southeast end to another float occurrence at the northwest end. The vein segment generally strikes N. 47° W. and has a maximum width estimated at 2 feet in a group of closely spaced shafts that Pruitt (1961) referred to as the Anthill mine. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein



Figure 16. Prospect on the Jumbo vein exposing a 1.1-foot-wide section of vein in NE $\frac{1}{4}$ SE $\frac{1}{4}$ section 26, T. 9 S., R. 21 E., SLBLM, Uintah County.

is poorly exposed and traceable for only a short distance in both directions, away from the exposures in the mine shafts at the Anthill site. Sparse outcrops, middle Uinta Formation stratigraphy, and the generally low topographic relief in this area make locating this gilsonite vein difficult. The vein system is reported by the BLM to continue as a group of veins for more than 5.5 miles to the northwest into the Uintah and Ouray Indian Reservation.

Antelope

The Antelope vein (plate 1) is located in SE $\frac{1}{4}$ section 36, T. 9 S., R. 20 E. We mapped the vein for 0.4 mile, from an occurrence of gilsonite float chips at the southeast end to a vein fracture outcrop at the northwest end. The vein generally strikes N. 47° W. and has a maximum width of 10 inches, as measured in a prospect trench. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein is traceable for only a short distance away from the prospect trench; however, sporadic, small gilsonite float chips were observed in the alluvial cover for more than a mile to the southeast. A filled-in prospect shaft having a wooden collar is located on the vein a short distance south of an improved dirt road in NW $\frac{1}{4}$ SE $\frac{1}{4}$ section 36.

Unreported Occurrences

We located a number of unreported gilsonite occurrences in and around SW $\frac{1}{4}$ NW $\frac{1}{4}$ section 33, T. 9 S., R. 21 E. They are approximately 0.5 mile southeast of a group of shafts (Anthill mine) on the Pride of Utah vein, but do not appear to be directly related to the main vein. We mapped the N. 70° W.-striking occurrences for 0.3 mile east of Cottonwood Wash. The occurrences consist of gilsonite veinlets less than 1 inch wide in fracture zones and small patches of gilsonite float chips.

We also located two unreported gilsonite occurrences in the middle of section 2, T. 10 S., R. 21 E. They are approximately 3.1 miles southeast of a group of shafts (Anthill mine) on the Pride of Utah vein and a short distance east of a prominent butte in section 2. The occurrences consist of two isolated patches of small gilsonite float chips that are less than 1 inch in width. The two patches could not be linked to any other gilsonite deposits.

Cottonwood

The Cottonwood vein system (plate 1) is located in section 6, T. 11 S., R. 22 E., section 1, T. 11 S., R. 21 E., sections 28, 29, 30, 33, 34, and 35, T. 10 S., R. 21 E., and possibly section 24, T. 10 S., R. 20 E. We mapped the vein system for approximately 7 miles, from an occurrence of small gilsonite float chips approximately 0.7 mile east of Sand

Wash at the southeast end to another occurrence of small float chips at the northwest end. The vein system generally strikes from N. 60 to 70° W. and has a maximum width of 4.5 feet, as exposed in a shaft wall approximately 0.3 mile east of Cottonwood Wash in an area having several other shafts. Uinta Formation sandstone and mudstone host the veins at the surface (plate 1). The vein system consists of two main veins of significant width with thinner, closely spaced, parallel, secondary veins cropping out in places. In SW $\frac{1}{4}$ section 28, T. 10 S., R. 21 E., the vein splits into a northern and southern vein. Both veins continue southeast, sporadically cropping out toward modern mine shafts; the southernmost vein appears to host the modern mine shafts and be wider than the northern vein. Southeast of the shafts, the vein trace is concealed under alluvium and colluvium for approximately 0.6 mile, reappearing as a single vein in Sand Wash. The vein appears to thin to the northwest and eventually disappears. Near the southeast end, the vein splits into 2.2-foot and 4-inch-wide veins before disappearing a short distance farther to the southeast. The southeast end of the Cottonwood vein system and the northwest end of the Bitter Water vein are separated by approximately 4 miles but lie roughly along the same trend. We investigated the area between the southeast end of the Cottonwood vein system and the canyon rim above Bitter Creek, but were unable to locate any gilsonite on the surface. However, topographic relief in this area, northwest of the Bitter Creek drainage, is generally low and gilsonite veins in the area tend to be poorly exposed. The vein system is also reported by the BLM to continue into the Uintah and Ouray Indian Reservation, cropping out farther to the northwest as the Black Bridge vein group in the Willow Creek area. The veins have been prospected along most of their lengths and as recently as 2008 hosted two active mine shafts (operated by Lexco Inc.) on the southernmost vein in SW $\frac{1}{4}$ section 35, T. 10 S., R. 21 E. Signs of significant historical mining were also observed in section 29, T. 10 S., R. 21 E., beginning from the midpoint in section 29 and continuing southeast for approximately 0.6 mile.

Black Cat

The Black Cat vein (plate 1) is located in S $\frac{1}{2}$ SW $\frac{1}{4}$ section 30, and partially in NW $\frac{1}{4}$ NE $\frac{1}{4}$ section 31, T. 10 S., R. 21 E. We mapped the vein for approximately 0.5 mile, from an occurrence of gilsonite float chips at the southeast end to an 8-inch-wide outcrop at the northwest end. The vein generally strikes N. 60° W. and has a maximum width of 1 foot, as measured a short distance southeast of the vein's midpoint. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The Black Cat vein consists of one main vein averaging 9 inches wide that is traceable for only a short distance before it disappears in both directions. The vein shows no signs of mining or prospecting.

Willow Creek/Wall

The Willow Creek/Wall vein (plate 1) is located in section 32, T. 10 S., R. 21 E., and section 25, T. 10 S., R. 20 E. The northwest segment of the vein is called Willow Creek and the southeast segment is called Wall. We mapped the vein for approximately 3 miles, from a 2.5-foot-wide outcrop in a prospect trench a short distance west of Cottonwood Wash at the southeast end to an occurrence of gilsonite float chips at the northwest end. The vein generally strikes N. 60° W. and has a maximum width of 3.5 feet, as exposed in a prospect trench a short distance northwest of a reclaimed mine site near the northwest end of the vein. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein consists of one main vein having a width between 2 and 3.5 feet along most of its length. The vein trace disappears for approximately 1.5 miles in a flat to low hilly area in the middle section of the vein. The vein disappears to the northwest not far past a reclaimed mine site and disappears to the southeast in the Cottonwood Wash area. However, the BLM reports exposures of a vein group (Willow Creek extension) on the Uintah and Ouray Indian Reservation northwest of Willow Creek, and these veins are also reported to continue northwest off the reservation. The vein has been prospected along most of its exposed length, and the southeast Wall segment contains historical mine sites consisting of trenches and shafts. The reclaimed mine site (ITM mine) is located on the northwest Willow Creek segment in NE $\frac{1}{4}$ SW $\frac{1}{4}$ section 25, T. 10 S., R. 20 E.

Ouray

The Ouray vein (plate 1) is located in NW $\frac{1}{4}$ section 26, T. 4 S., R. 2 E., Uinta Base Line and Meridian (UBLM). We mapped the vein for approximately 0.3 mile, from a 6-inch-wide outcrop at the southeast end to an occurrence of gilsonite float chips at the northwest end. The vein generally strikes N. 62° W. and has a maximum width of 1.5 feet, as exposed in a prospect trench a short distance southeast of a mine shaft. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). We could only trace the vein a short distance southeast and northwest of the mine shaft located mid-vein. The historical mine site appears somewhat significant, having two cabins, a collapsed headframe, and a wooden-collared, open shaft. A small copper deposit, associated with fossilized wood, has been prospected within a mile of the vein to the southeast.

Little Seam

The Little Seam vein (plate 1) is located in NW $\frac{1}{4}$ NE $\frac{1}{4}$ section 36, T. 4 S., R. 2 E., UBLM. We mapped the vein for approximately 0.2 mile, from a 3-inch-wide outcrop (the only gilsonite outcrop encountered on the vein) exposed

in a prospect shaft wall at the southeast end to a prospect shaft at the northwest end. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein could only be traced a short distance and was locatable only via three prospect shafts aligned along a N. 56° W. trend. No other naturally occurring gilsonite outcrops or float occurrences were observed elsewhere along the trend of the shafts. The Little Seam vein may be related to the Ouray vein, which is located approximately on trend 1.3 miles to the northwest.

Gem

The Gem vein (plate 1) is located mostly in NW $\frac{1}{4}$ SW $\frac{1}{4}$ section 36, T. 4 S., R. 2 E., UBLM. We mapped the vein for 0.2 mile along a N. 46° W. trend between two prospect shafts. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein was locatable only via two prospect shafts containing gilsonite chips. No other naturally occurring gilsonite outcrops or float occurrences were observed elsewhere along the trend of the shafts.

Little Boy

The Little Boy vein (plate 1) is located in SW $\frac{1}{4}$ section 36, T. 4 S., R. 2 E., UBLM. We mapped the vein for approximately 0.5 mile, from a prospect shaft at the southeast end to a 6-inch-wide outcrop exposed in a prospect shaft wall at the northwest end. The vein generally strikes N. 45° W. and has a maximum width estimated to be 2 feet on an outcrop exposed in a prospect shaft a short distance northwest of the vein's midpoint. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein can be traced for only a short distance and was locatable mostly via prospect shafts; however, a few gilsonite outcrops and float chip occurrences are present along the vein trend.

Cliff Dweller

The Cliff Dweller vein (plate 1) is located mostly in SW $\frac{1}{4}$ SE $\frac{1}{4}$ section 2, T. 5 S., R. 2 E., UBLM. We mapped the vein for approximately 800 feet, from a 6-inch-wide outcrop at the southeast end to a 6-inch-wide outcrop exposed in a prospect shaft at the northwest end. The vein generally strikes N. 55° W. and has an average width of 6 inches along its length. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein was locatable mostly via three small, shallow, prospect shafts (figure 17). The Ouray 7.5-minute quadrangle map shows an adit near the west bank of the Green River, on the side of a steep butte, that is on trend with the prospect shafts on top of the butte.

A prospect shaft was observed in a gully northeast of the

Cliff Dweller vein that is presumably the location of the Green River vein reported by Pruitt (1961). We did not examine the Green River vein in this field investigation, but Pruitt (1961) reported that the Cliff Dweller, Green River, and Black Butte veins are within a mile of one another and similar in strike and dimension. The prospect shaft presumably on the Green River vein is located in SE $\frac{1}{4}$ section 2, T. 5 S., R. 2 E., UBLM.

Willow Group

At least four veins in the Willow Group (plate 1) are located in section 19, T. 9 S., R. 20 E, and sections 13 and 24, T. 9 S., R. 19 E. Three of the veins appear to originate from the same area in SE $\frac{1}{4}$ section 13. Rocks hosting the veins belong to the Uinta Formation (plate 1) and consist of interbedded, thin to thick beds of discontinuous chan-



Figure 17. Prospect shaft on the Cliff Dweller vein. Person is standing on the northwest-trending vein located in SW $\frac{1}{4}$ SE $\frac{1}{4}$ section 2, T. 5 S., R. 2 E., UBLM, Uintah County.

nel sandstone and thin- to thick-bedded mudstone. Gilsonite veins contained in these beds are discontinuous and tend to vary in width, meander, and split. The Willow Group veins are discussed from south to north.

Willow #1, the southernmost vein, may be two independent veins or one vein that has a slight bend. Colluvial cover in a flat area obscures one-third of a mile of the central area of the vein, which makes it difficult to determine whether or not it is a single vein; however, for simplicity it is discussed as a single vein. We mapped the Willow #1 vein for approximately 1.8 miles, from an occurrence of gilsonite float chips a short distance west of the Uintah and Ouray Indian Reservation boundary at the southeast end to a 2.2-foot-wide outcrop exposed in a prospect trench east of the Green River at the northwest end. The vein generally strikes from N. 50 to 70° W. and has a maximum width of 2.2 feet, as measured in a short prospect trench at the northwest end of the vein. The southeast segment of the vein meanders, contains splits, and is considerably narrower, never exceeding 8 inches. The northwest segment of the vein is straighter and wider and is explored by several short prospect trenches on a low hill, the only prospects located on the vein.

We mapped the Willow #2 vein for approximately 1.4 miles, from a 10-inch-wide outcrop a short distance west of the Uintah and Ouray Indian Reservation boundary at the southeast end to a 1.2-foot-wide outcrop at the northwest end, where it appears to intersect the Willow #1 vein to the south. The vein is sinuous, generally strikes N. 63° W., and has a maximum width of 1.5 feet near the northwest end. Secondary, thinner veins are present at the southeast end. The vein continues southeast onto the reservation and the BLM survey also shows the vein extending for over a mile onto the reservation. Two short prospect trenches occur on the middle of the northwest segment of the vein on outcrops about 1 foot wide.

We mapped the Willow #3 vein for approximately 0.5 mile, from a 5-inch-wide outcrop at the southeast end to an occurrence of gilsonite float chips next to the Uintah and Ouray Indian Reservation boundary at the northwest end. An isolated prospect trench striking N. 76° W. exposes a 2-foot-wide vein approximately 0.4 mile to the northwest and may be related to the Willow #3 vein. The Willow #3 vein generally strikes N. 68° W. and has a maximum width of 1.8 feet exposed in a short prospect trench near the northwest end. The vein appears to be short and narrow, containing one anomalous 1.8-foot-wide outcrop in a prospect trench. However, if the isolated 2-foot-wide deposit to the northwest is related to this vein, more than 0.5 mile of 2-foot-thick vein may exist under the alluvial cover that separates these two prospects.

We mapped the Willow #4 vein (the northernmost vein of the Willow group) for approximately 0.3 mile, from a

5-inch-wide outcrop at the southeast end to a 7-inch-wide outcrop next to the Uintah and Ouray Indian Reservation boundary at the northwest end. The vein generally strikes N. 60° W. and is short and narrow, appearing only on one low butte. It is possible that the isolated prospect trench containing a vein 2 feet wide, previously discussed in relation to the Willow #3 vein, may instead be related to this vein. The vein shows no other signs of mining or prospecting.

Turtle Shell

The Turtle Shell vein (plate 1) is located in N½ section 24, T. 9 S., R. 19 E. We mapped the vein for approximately 0.8 mile, from a 10-inch-wide outcrop a short distance west of Willow Creek at the southeast end to a 3-inch-wide outcrop a short distance east of the Green River at the northwest end. The vein generally strikes N. 66° W. and has a maximum width of 1 foot on an outcrop southeast of the vein's midpoint. Closely spaced, parallel veinlets are also present. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein appears to be short and narrow and shows no signs of mining or prospecting.

Petroglyph

The Petroglyph vein (plate 1) is located in section 30, T. 9 S., R. 20 E., and N½NE¼ section 25, T. 9 S., R. 19 E. We

mapped the vein for approximately 1.2 miles, from an 8-inch-wide outcrop a third of a mile east of Willow Creek at the southeast end to an outcrop at least 6 inches wide a half mile west of Willow Creek at the northwest end. The vein generally strikes N. 56° W. and has a maximum width of 1 foot, as measured near the southeast end. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The vein consists of one main vein that averages 9 inches wide along the well-exposed southeast segment east of Willow Creek, but then becomes less exposed to the northwest where it thins and then splits just before disappearing. The vein is unmined and only one or two superficial prospects are present. The Petroglyph vein gets its name from the numerous Native American rock carvings in the area.

Black Diamond

The Black Diamond vein (plate 1) is located mostly in S½NE¼ section 36, T. 9 S., R. 19 E. We mapped the N. 56° W.-striking vein for 0.4 mile, from an occurrence of gilsonite float chips at the southeast end to a 10-inch-wide outcrop at the northwest end. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). A historical mine site (figure 18) is located on the middle segment of the vein and appears fairly significant, having an open shaft and a ruined stone building present. A 1.4-foot-wide outcrop (the vein's maximum known width) is also present at the mine site. Pruitt (1961)



Figure 18. Black Diamond mine in SE¼NE¼ section 36, T. 9 S., R. 19 E., SLBLM, Uintah County.

referred to the site as the “Black Diamond mine” and further reported that two short veins of about the same length are also present, and that all three are located within a 200-foot-wide zone. We were unable to locate these two veins in the area where they are reported to crop out, and an extended search to the north and south was also unsuccessful.

Turkey Trail Hill

The Turkey Trail Hill vein (plate 1) is located in N½ section 9, T. 10 S., R. 20 E. The mapped extent of the vein is approximately 1 mile, striking N. 60° W. and mostly concealed in the Willow Creek drainage. At the southeast end, a 2-foot-wide exposure (the vein’s maximum known width) occurs in a collapsed shaft located next to an improved dirt road in the Willow Creek drainage, and at the northwest end a 4-inch-wide outcrop occurs just prior to the vein disappearing. Uinta Formation sandstone and mudstone host the vein at the surface (plate 1). The southeast segment of the vein is on the Uintah and Ouray Indian Reservation, so the extent could not be determined, but the northwest end appears to thin west of Willow Creek in a thick sequence of interbedded mudstone and channel sandstone in the middle Uinta Formation. Only one collapsed shaft was observed on the vein in SE¼NE¼ section 9. The Turkey Trail Hill vein may be related to the Black Diamond vein that is located approximately on trend 3 miles to the northwest.

Willow Creek #1 and #2

The Willow Creek #1 and #2 vein system (plate 1) is located in sections 7, 8, 17, and 18, T. 10 S., R. 20 E., and sections 1 and 12, T. 10 S., R. 19 E. The vein system consists of two main, subparallel veins separated by several hundred feet to the northwest but appearing to converge to the southeast; secondary, thinner split-off veins also exist in places. We mapped the Willow Creek #1 vein for 3.3 miles, from an occurrence of gilsonite float chips near the Uintah and Ouray Indian Reservation boundary at the southeast end to a 6-inch-wide outcrop near an abandoned well site at the northwest end. The vein generally strikes from N. 50 to 58° W. and has a maximum width of 1 foot, as measured on an outcrop near the northwest end.

We mapped the Willow Creek #2 vein for approximately 1.2 miles, from an occurrence of gilsonite float chips near the Uintah and Ouray Indian Reservation boundary at the southeast end to a 5-inch-wide outcrop at the northwest end. The vein generally strikes N. 54° W. and has a maximum width of 1.8 feet (the system’s maximum known width) measured on an outcrop near the northwest end. Uinta Formation sandstone and mudstone host the vein system at the surface (plate 1). Both veins are poorly

exposed and mostly traceable by sporadic gilsonite float and vein exposures. The vein system is reported by the BLM to continue for approximately 1 mile to the southeast in sections 16 and 21, T. 10 S., R. 20 E., on the Uintah and Ouray Indian Reservation. The veins show no signs of mining and only a few superficial prospects.

O.K., Original Owner, Black Dog, and Florence

A complex system of four veins (plate 1) is located 9 miles west of the town of Ouray in sections 26, 35, and 36, T. 4 S., R. 1 E., UBLM. Rocks hosting the veins belong to the Uinta Formation (plate 1) and consist of interbeds of some thin to thick beds of discontinuous channel sandstone, but predominantly thin- to thick-bedded mudstone. The vein system coincides with the east-west-trending Duchesne fault system (plate 1), and two of these veins are also east-west trending. The O.K. vein coincides with a fairly significant fault and was mapped for approximately 0.7 mile, striking from N. 80 to 90° W. The Original Owner vein lies approximately 0.3 mile south of the O.K. vein and was mapped for 0.25 mile, also striking from N. 80 to 90° W. The Black Dog vein is located near the west end of the O.K. vein, has a maximum width of 7 inches, and was mapped for approximately 0.4 mile, generally striking N. 50° W. These three veins appear to be relatively short and narrow, never reaching a width much greater than 0.5 foot. Copper mineralization is sporadically present in sandstone beds near the Duchesne fault and gilsonite veins in this area as well.

The Florence vein, which is located between the two east-west-striking veins, was mapped for approximately 0.8 mile. The vein generally strikes N. 47° W. in the middle and northwest segments, and at the northwest end it appears to abruptly terminate where it intersects the O.K. vein near its mapped midpoint. The Florence vein abruptly changes strike to between N. 75 and 80° W. at the southeast end where a caved shaft is present on the widest observed outcrop (1.8 feet) of the four-vein system. This vein dips very steeply (>70°) to the northeast where exposed in the shaft wall. The O.K., Original Owner, and Black Dog veins contain one or two prospects on each vein, but the Florence vein hosts the most significant prospecting and possibly some mining. The vein segment that strikes N. 47° W. averages 1 foot in width and was explored by significant trenching. The Florence vein is reported by Pruitt (1961) to continue for more than 3 miles to the southeast along the vein segment trending approximately N. 75° W. We briefly investigated outcrops that lie along this trend but could not trace the vein for more than 0.8 mile. The area where the continuation is reported to be located is mostly covered by alluvial and eolian deposits, but there are a few sporadic bedrock outcrops and the area may be worth further investigation.

Pariette Area Veins

Castle Peak, Baxter, and Pariette

The Castle Peak, Baxter, and Pariette vein system (plate 1) is located in sections 4, 5, and 9, T. 9 S., R. 17 E., and sections 19, 30, 31, and 32, T. 8 S., R. 17 E. The vein system is composed of the Castle Peak vein at the southeast, the Baxter vein in the middle area, and the Pariette vein at the northwest. We mapped the vein system for approximately 5 miles, from a prospect at the southeast end to a prospect at the northwest end. Uinta Formation sandstone and mudstone host the veins at the surface (plate 1). The Castle Peak segment generally strikes N. 40° W. and has a few outcrops about 2 feet wide exposed along its length. The Baxter segment is a broken area of gilsonite veins and veinlets in the central area of the system where the Castle Peak and Pariette veins meet. The Pariette segment generally strikes N. 30° W., and Pruitt (1961) reported that it averages 1.5 feet wide at the surface. Pruitt (1961) provided a detailed description of the vein system and its history and indicated that the vein system continues for more than 3 miles southeast of the last exposure we observed on the Castle Peak vein; however, we found no gilsonite deposits in the flat to low hilly topography that is mostly covered by alluvium. The vein system was one of the earliest gilsonite deposits to be mined in the basin, starting in 1890, and by 1960 a shaft on the Pariette vein had reached a depth of 1500 feet (Pruitt, 1961). Much of the Castle Peak, Baxter, and Pariette vein system mining disturbance has been reclaimed.

Gilsonite Draw

The Gilsonite Draw vein (plate 1) is located in NW $\frac{1}{4}$ NW $\frac{1}{4}$ section 31, T. 9 S., R. 16 E., and SE $\frac{1}{4}$ SE $\frac{1}{4}$ section 25, and partially in NE $\frac{1}{4}$ NE $\frac{1}{4}$ section 36, T. 9 S., R. 15 E. We mapped the vein for 0.25 mile in Gilsonite Draw, from an occurrence of gilsonite float chips at the southeast end to a fracture zone 6 inches wide containing veinlets at the northwest end. The vein generally strikes N. 35° W. and is poorly exposed. Thinly bedded Uinta Formation marlstone, sandstone, and mudstone host the vein at the surface (plate 1). A reclaimed prospecting or mining site in the bottom of the Gilsonite Draw drainage is reported by the UGS Utah Mineral Occurrence System to have contained shafts and 2-foot-wide trenches.

Fort Duchesne Area Veins

The first reports of gilsonite in the Uinta Basin came from the Fort Duchesne area in 1869, where two veins, the Carbon and Raven, are located in the Duchesne River Formation (Pruitt, 1961). The areas where these two veins are reported to be located are mostly covered by agriculture

and various other developments, and we were unable to locate any gilsonite outcrops associated with the Carbon or Raven veins. Therefore, these veins are not shown on plate 1. The following discussion of the two veins is taken from Pruitt's 1961 summary.

The Carbon vein is located in T. 2 S., R. 2 E., UBLM, crossing beneath U.S. Highway 40 near Gusher, Utah. This vein was the first gilsonite vein to be mined in the Uinta Basin, starting around 1888 by Samuel Gilson, and had a long history of production. The Carbon vein is more than 3 miles long, strikes N. 40° W., and is reported as having a maximum width of 3 to 4 feet that is maintained for 1.5 miles at the middle section of the vein. The Raven vein is located in the north half of T. 2 S., R. 1 E., UBLM, north of U.S. Highway 40 and Fort Duchesne, Utah. This vein was extensively mined, mainly from 1904 to 1939. The Raven vein is approximately 3 miles long, strikes N. 37° W., and is reported to average between 1.5 and 2 feet wide.

U.S. Bureau of Land Management Gilsonite Mapping

In the early 1980s, BLM employees conducted mapping of selected gilsonite veins on the Ouray, Ouray SE, Big Pack Mtn. NW, Big Pack Mtn. NE, Archy Bench, Asphalt Wash, and Rainbow 7.5-minute topographic quadrangles (Peter Sokolosky, BLM, written communication, April 2, 2004). The survey appears to have focused on veins in old BLM prospecting permit application areas. Many of the veins mapped by the BLM occur on the Uintah and Ouray Indian Reservation, which UGS geologists were unable to get permission to visit. UGS geologists did observe some of these veins in road cuts and outcrops along the public roads that pass through the reservation. In the areas covered by both field investigations, mapping mostly agrees except for small discrepancies. Plate 1 shows the significant veins mapped by the BLM, most of which occur on the Uintah and Ouray Indian Reservation.

Various veins and vein systems are reported by the BLM. The Pride of Utah vein system is reported in sections 14, 15, 16, 23, 24, and 25, T. 9 S., R. 20 E., and sections 29 and 30, T. 9 S., R. 21 E. The vein system is reported to be composed of several veins extending approximately 6 miles, generally strike N. 58° W., and have outcrops as wide as 2 feet. The Natural Buttes vein is reported in sections 15, 16, 22, and 23, T. 9 S., R. 20 E. The vein is reported to be approximately 2.2 miles long, strike N. 55° W., and have outcrops as wide as 8 inches. The Black Gnat vein is reported in sections 16, 21, and 22, T. 9 S., R. 20 E. The vein is reported to be approximately 2.4 miles long, strike N. 54° W., and have outcrops as wide as 2.5 feet. The Workman vein is reported in sections 17 and 20, T. 9 S., R. 20 E. The vein is reported to be approximately

0.9 mile long, strike N. 53° W., and have outcrops as wide as 14 inches. The Uintah and Ouray Indian Reservation southeast extension of the Willow #2 vein is reported in sections 20 and 29, T. 9 S., R. 20 E. This extension is reported to be approximately 1.3 miles long, generally strike N. 54° W., and have a maximum width of 1.2 feet. The Black Bridge vein group (possibly an extension of the Cottonwood vein system) is reported in sections 15, 16, 22, and 23, T. 10 S., R. 20 E. The vein group is reported to be composed of at least two veins, with the most significant vein extending approximately 2.2 miles, being variable in strike, and having a maximum width of 2.5 feet. A southeast extension of the Willow Creek #1 and #2 vein system is reported in sections 16 and 21, T. 10 S., R. 20 E. This extension is reported to be composed of two parallel, closely spaced veins less than a mile long generally striking N. 57° W., and ranging in width from 5 to 11 inches. Other veins of small dimensions are also reported to occur in the vicinity of the Black Bridge, Willow #2 southeast extension, Black Gnat, Natural Buttes, and Pride of Utah veins.

Unconfirmed Deposits

Gilsonite veins discussed in this section are veins mentioned in Pruitt's (1961) report but were not confirmed by our study. Some of the veins reported by Pruitt were not located in our field investigation. Other veins were not examined due to property access restrictions. Some reported veins are questionable because of vague location descriptions. Our inability to locate some veins reported by Pruitt does not necessarily mean they do not exist, but that they may be difficult to locate due to various factors. The gilsonite field is located in a remote and rugged region and access to some areas is difficult. Veins can be concealed by unconsolidated deposits and low topographic relief, and scarce bedrock outcrops can make locating gilsonite veins at the surface difficult. Vein width and stratigraphic location can also hinder locating veins. Narrow veins in thinly interbedded sandstone and mudstone are commonly discontinuous, meandering, and generally poorly exposed. Past maps of the gilsonite field are commonly simplified, general depictions of veins, and the veins may not have been located on accurate base maps. Other veins may exist in the Uinta Basin that were not located by this study or by any previous work. However, it is unlikely that thick veins having good surface exposures went unnoticed by early prospectors. In the gilsonite field, numerous veins of small dimensions are present near other more significant veins, or are found isolated.

A short, unnamed gilsonite vein is reported (Pruitt, 1961) to be located in section 32, T. 10 S., R. 23 E., approximately 0.3 mile southwest of the Harrison vein. During a brief field reconnaissance of the area we found only a few small gilsonite float chips. A gilsonite vein probably

exists in the area, perhaps covered by alluvial and colluvial deposits.

Pruitt (1961) reported that the Little Jack vein is located north of Ouray, Utah, and east of the Duchesne River in section 20, T. 4 S., R. 3 E., UBLM. The vein is reported to be 8 to 10 inches wide, 1.5 miles long, and strike N. 70° W. We did not investigate the area due to property restrictions.

Three gilsonite veins are reported (Pruitt, 1961) in the area near the southeast end of the Cottonwood vein system in and around Sand Wash. A short unnamed vein approximately 1 mile long is reported to lie a short distance north of the Cottonwood vein in Sand Wash, in SW¼ section 36, T. 10 S., R. 21 E. We were unable to locate any gilsonite deposits in Sand Wash where the vein was reported. The Red Rock vein is reported (Pruitt, 1961) to branch off the Cottonwood vein to the southeast in section 35, T. 10 S., R. 21 E., near two modern shafts, and continue for approximately 5 miles. Jim Lekas (Lexco Inc., verbal communication, 2004), who operated the two shafts on the Cottonwood vein, stated that he was unable to find that vein. Pruitt (1961) reported that he could not confirm the existence of the vein, and that its existence was doubtful. We were also unable to locate any gilsonite deposits in Sand Wash where the Red Rock vein is reported; therefore we believe its existence is questionable. This reported branch may be a misrepresentation of the vein split that is present on the Cottonwood vein in SW¼ section 28, T. 10 S., R. 21 E. Another unnamed vein is reported (Pruitt, 1961) to be located in section 2, T. 11 S., R. 21 E., approximately 1 mile southwest of the two modern shafts on the Cottonwood vein, and is reported to be about 1.7 miles long. We were unable to locate any gilsonite deposits where it is reported to crop out. The existence of these three veins located in the area near the southeast end of the Cottonwood vein system are in doubt; however, poorly exposed bedrock and the generally low topographic relief of the area makes locating gilsonite outcrops difficult.

Pruitt (1961) reported that the Canyon vein is located in section 26, T. 4 S., R. 2 E., UBLM, and that it is on trend with the Little Seam vein to the southeast and perhaps is part of the same vein. The vein is also reported to be 1.5 feet wide and strike N. 55° W. However, we were unable to locate any gilsonite deposits in the area where the vein is reported.

The Black Butte vein is reported (Pruitt, 1961) to be located in section 11, T. 5 S., R. 2 E., UBLM. We did not investigate the area, but believe the vein likely exists somewhere north of its reported location. We base this presumption on the existence of the Cliff Dweller and Green River veins that are located within a mile to the northeast of the reported location.

We investigated an area that may be the site of the April vein, which contained some small gilsonite chips and bitumen-impregnated sandstone outcrops. The area is in NW¼SE¼ section 22, T. 8 S., R. 17 E., where two northwest-trending shaft symbols appear on the Pariette Draw SW 7.5-minute quadrangle map. However, we did not observe any shafts or significant gilsonite deposits, and the surrounding area is now mostly covered by agriculture. The area where the April vein is reported (Pruitt, 1961) is approximately 1 mile southwest of the shaft symbols and 2 miles northeast of the Pariette mine. The vein is also reported striking parallel to the Pariette vein for 3000 feet. We briefly investigated the reported vein location area and were unsuccessful in locating any gilsonite deposits.

CONCLUSION

Gilsonite, a member of the asphaltite group of hydrocarbon bitumens, forms a swarm of subparallel, northwest-trending, near-vertical, laterally and vertically extensive veins in the Uinta Basin of Utah and Colorado. Gilsonite is sourced from the Mahogany oil shale zone in the Parachute Creek Member of the Green River Formation. Mechanisms for propagating the fractures that contain gilsonite appear to involve elevated pore pressure within the hydrocarbon source beds of the Green River Formation and regional stress fields. In this study we mapped and described in detail 59 veins, vein systems, and isolated vein outcrops, having a total combined vein length of over 120 miles. This report documents a total of 71 significant veins, vein systems, and vein extensions that were mapped by the UGS or previous investigators, having a total combined vein length of over 170 miles.

Gilsonite veins in the study area occur in Eocene Green River Formation lacustrine deposits and Uinta Formation marginal lacustrine and fluvial deposits. All but two (Weaver and Black Dragon) of the known significant veins are located above the Mahogany oil shale zone, and no gilsonite veins are known to cross-cut the Mahogany oil shale zone contained in the Parachute Creek Member of the Green River Formation. Gilsonite veins above the source beds in the eastern part of the study area are commonly eroded down near their roots, and the remaining veins generally can be expected to split and thin with depth. Gilsonite veins exposed beneath the source beds attain their greatest thickness in the Douglas Creek Member of the Green River Formation and the interfingering Renegade Tongue of the Wasatch Formation, and can be expected to thin both downward and upward from these units. Bedrock hosting gilsonite veins in the central and northwestern parts of the study area is generally less eroded, and the veins rooted in the Green River Formation oil shale have greater vertical extent and generally

can be expected to widen with depth from exposures in the middle and upper Uinta Formation.

The continuity of the veins can be impressive in some areas of the study area, having veins stretching in relatively long, straight ribbons across the hills of the Uinta Basin, and the longest vein system (Pride of the West-Rainbow-Black Dragon) extending more than 22 miles. The gilsonite veins can also be vertically continuous, ranging from hundreds to more than 3000 feet, commonly having only small variations in width. However, veins can be difficult to locate in some areas of the study area due to concealment by unconsolidated deposits, low topographic relief, and vein width and stratigraphic location. Narrow veins in thinly interbedded sandstone and mudstone are commonly discontinuous, meandering, and generally poorly exposed. Many of the veins in the study area appear to lie roughly along the same trend as other veins, but lack exposed gilsonite at the surface in extensive areas between the veins. All but two (O.K. and Original Owner) of the known significant gilsonite veins trend northwest; the two veins trend east-west and coinciding with the Duchesne fault system.

Gilsonite is a geologically interesting and economically significant resource, having an extremely wide range of uses that have changed over time with new technology and industrial needs. Gilsonite has a long, colorful mining history dating back to the late 1800s, and all of the veins mined today were discovered by early prospectors who located surface exposures. The original in-place gilsonite resource of all the veins is estimated by Cashion (1967) to be approximately 45 million short tons. The remaining resources tend to be in the deeper parts of the veins and in thinner, more remote veins that will likely be more expensive to mine.

ACKNOWLEDGMENTS

Funding for this study was provided by the Utah School and Institutional Trust Lands Administration and the Utah Geological Survey. We thank American Gilsonite Company; Lexco, Inc.; and Ziegler Chemical and Mineral Corp. for access to their properties and the information that they shared. We thank Pete Sokolosky of the U.S. Bureau of Land Management for access to unpublished BLM gilsonite vein maps. Dallas Rippey and Ammon McDonald, formerly of the UGS, contributed greatly to the field mapping of the veins, and Rippey also compiled much of the data in the appendix. We appreciate assistance from Buck Ehler (UGS) for additional GIS project support. Jeff Quick (UGS) provided some reference material and advice, and Roger Bon (UGS, retired) contributed some recent gilsonite production data. This report was greatly enhanced by peer reviews from Tom Faddies

(SITLA), Mark Longman (Questar Exploration and Production Company), Craig Morgan (UGS), Andrew Rupke (UGS), and Doug Sprinkel (UGS).

REFERENCES

- Abraham, H., 1960, *Asphalts and allied substances - their occurrence, modes of production, uses in the arts and methods of testing* (6th edition), Volume 1, Historical review and natural raw materials: Princeton, New York, D. Van Nostrand Company, Inc., p. 1–302.
- American Gilsonite Company, 2008, Home page: Online, www.americangilsonite.com/, accessed April 2008.
- Baker, J.H., 1950, Economics of gilsonite in the Uinta Basin, in Eardley, A.J., editor, *Petroleum geology of the Uinta Basin: Intermountain Association of Petroleum Geologists Guidebook to the Geology of Utah* 5, p. 119–120.
- Bell, K.G., and Hunt, J.M., 1963, Native bitumens associated with oil shales, in Breger, I.A., editor, *Organic geochemistry: New York, The Macmillan Company, and Oxford, Pergamon Press, Earth Science Series, Monograph* 16, p. 333–366.
- Bender, H.E., Jr., 1970, *Uintah railway—the gilsonite route: Berkeley, California, Howell-North Books*, 239 p.
- Blake, W.P., 1885, Uintahite—a new variety of asphaltum from the Uintah Mountains, Utah: *The Engineering and Mining Journal*, v. 40, no. 26, p. 431.
- Blake, W.P., 1890, Uintahite, albertite, grahamite, and asphaltum described and compared, with observations on bitumen and its compounds: *Transactions of the Society of Mining Engineers of American Institute of Mining, Metallurgical and Petroleum Engineers, Incorporated*, v. 18, p. 563–582.
- Boden, T., and Tripp, B.T., 2008, Gilsonite resources of the Uinta Basin, Utah, in Longman, M.W., and Morgan, C.D., editors, *Hydrocarbon systems and production in the Uinta Basin, Utah: Utah Geological Association, and Rocky Mountain Association of Geologists Publication* 37, p. 367–389, CD.
- Boleneus, D.E., 2007, Application of earth resistivity imaging to land management in Utah: U.S. Department of the Interior, Bureau of Land Management, unpublished draft manuscript, 19 p.
- Botbol, J.M., 1961, *Geochemical exploration for gilsonite: Salt Lake City, University of Utah, M.S. thesis*, 42 p.
- Bryant, B., 1992, Geologic and structure maps of the Salt Lake City 1° x 2° quadrangle, Utah and Wyoming: U.S. Geological Survey Miscellaneous Investigations Series Map I-1997, scale 1:125,000.
- Business Wire, 2008, American Gilsonite Company announces acquisition: Online, <http://www.reuters.com/article/pressRelease/idUS17981+22-Feb2008+BW20080222>, accessed April 2, 2008.
- Carey, G.A., and Roberts, I.C., 1949, *Dissertation on the history, occurrence, mining, and economics of gilsonite: Salt Lake City, University of Utah, B.S. thesis*, 89 p.
- Cashion, W.B., 1967, *Geology and fuel resources of the Green River Formation, southeastern Uinta Basin, Utah and Colorado: U.S. Geological Survey Professional Paper* 548, 48 p.
- Cashion, W.B., 1969, Other bituminous substances, in Hilpert, L.S., editor, *Mineral and water resources of Utah: Utah Geological and Mineralogical Survey Bulletin* 73, p. 63–70.
- Cashion, W.B., 1974, *Geologic map of the Southam Canyon quadrangle, Uintah County, Utah: U.S. Geological Survey Miscellaneous Field Studies Map* MF-579, scale 1:24,000.
- Cashion, W.B., 1977, *Geologic map of the Weaver Ridge quadrangle, Uintah County, Utah and Rio Blanco County, Colorado: U.S. Geological Survey Map* MF-823, scale 1:24,000.
- Cashion, W.B., 1978, *Geologic map of the Walsh Knolls quadrangle, Uintah County, Utah and Rio Blanco County, Colorado: U.S. Geological Survey Miscellaneous Field Studies Map* MF-1013, scale 1:24,000.
- Cashion, W.B., 1986, *Geologic map of the Bonanza quadrangle, Uintah County, Utah: U.S. Geological Survey Miscellaneous Field Studies Map* MF-1865, scale 1:24,000.
- Cornelius, C.D., 1987, Classification of natural bitumen—a physical and chemical approach, in Meyer, R.F., editor, *Exploration for heavy crude oil and natural bitumen: American Association of Petroleum Geologists Studies in Geology* 25, p. 165–174.
- Covington, R.E., 1964, A brief history of early mineral exploitation in the Uinta Basin, in Sabatka, E.F., editor, *Geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse: Intermountain Association of Petroleum Geologists Guidebook, 13th Annual Field Conference*, p. 1–16.
- Crawford, A.L., 1957, Gilsonite—its discovery and the early history of the industry, in Seal, O.G., editor, *Geology of the Uinta Basin: Intermountain Association of Petroleum Geologists, Eighth Annual Field Conference*, p. 149–151.
- Crawford, A.L., and Pruitt, R.G., Jr., 1963, Gilsonite and other bituminous resources of central Uintah County, Utah, in Crawford, A.L., editor, *Oil and gas possibilities of Utah, re-evaluated: Utah Geological and Mineralogical Survey Bulletin* 54, p. 215–224.
- Curiale, J.A., 1986, Origin of solid bitumens, with emphasis on biological marker results: *Organic Geochemistry*, v. 10, p. 559–580.

- Davis, L.J., 1951a, The characteristics, occurrences and uses of the solid bitumens of the Uinta Basin, Utah: Provo, Utah, Brigham Young University, M.S. thesis, 93 p.
- Davis, L.J., 1951b, Characteristics, occurrence and uses of the solid bitumens of the Uinta Basin, Utah: The Compass [of Sigma Gamma Epsilon], v. 29, no. 1, p. 32–39.
- Davis, L.J., 1957, Geology of gilsonite, in Seal, O.G., editor, Geology of the Uinta Basin: Intermountain Association of Petroleum Geologists Guidebook, 8th Annual Field Conference, p. 152–156.
- Dennis, E., 1930, A preliminary survey of Utah non-metallic minerals, exclusive of mineral fuels, with special reference to their occurrence and markets for them: Provo, Utah, Brigham Young University, M.A. thesis, p. 108–114.
- Dickie, J.P., and Yen, T.F., 1967, Macrostructures of the asphaltic fractions by various instrumental methods: Analytical Chemistry, v. 34, no. 14, p. 1847–1952.
- Eldridge, G.H., 1901, The asphalt and bituminous rock deposits of the United States, in Walcott, C.D., director: U.S. Geological Survey Twenty-second Annual Report of the United States Geological Survey to the Secretary of the Interior, pt. 1, p. 209–364.
- Erslev, E.A., 1993, Laramide basement tectonics, in Schmidt, C.J., Chase, R.B., and Erslev, E.A., editors, Laramide basement deformation in the Rocky Mountains foreland of the western United States: Geological Society of America Special Paper 280, p. 339–358.
- Garvin, R.F., compiler, 1966, A directory of the mining industry of Utah—1965: Utah Geological and Mineralogical Survey Bulletin 79, 94 p.
- Hatcher, H.J., Meuzelaar, H.L.C., and Urban, D.T., 1992, A comparison of biomarkers in gilsonite, oil shale, tar sand and petroleum from Threemile Canyon and adjacent areas in the Uinta Basin, Utah, in Fouch, T.D., Nuccio, V.F., and Chidsey, T.C., Jr., editors, Hydrocarbon and mineral resources of the Uinta Basin, Utah and Colorado: Utah Geological Association Guidebook 20, p. 271–287.
- Hawes, B., 1990, Uintaite, strange Utah mineral, has many uses: Pay Dirt Magazine, July, p. 4B-7B.
- Hays, W.W., Nuttli, O.W., and Scharon, L., 1967, Mapping gilsonite veins with the electrical resistivity method: Society of Exploration Geophysicists, Geophysics, v. 32, no. 2, p. 302–310.
- Henderson, J.H., Jr., 1957, The gilsonite refining project of the American Gilsonite Company, in Seal, O.G., editor, Geology of the Uinta Basin: Intermountain Association of Petroleum Geologists Guidebook, 8th Annual Field Conference, p. 157–160.
- Hunt, J.M., 1963, Composition and origin of the Uinta Basin bitumens, in Crawford, A.L., editor, Oil and gas possibilities of Utah, re-evaluated: Utah Geological and Mineralogical Survey Bulletin 54, p. 249–273.
- Hunt, J.M., 1979, Petroleum geochemistry and geology: San Francisco, W.H. Freeman Co., 617 p.
- Hunt, J.M., Stewart, F., and Dickey, P.A., 1954, Origin of hydrocarbons of Uinta Basin, Utah: Bulletin of the American Association of Petroleum Geologists, v. 38, no. 8, p. 1671–1698.
- Jackson, D., 1985, American Gilsonite—mining solid hydrocarbon, in Picard, M.D., editor, Geology and energy resources, Uinta Basin of Utah: Utah Geological Association Publication 12, p. 257–261.
- Jacob, H., 1983, Recent studies on the genesis of natural solid oil bitumens: Translated from the German, Geologisches Jahrbuch, Series D., no. 59, p. 3–61, RP: Report Number 286914.
- Keighin, C.W., 1977, Preliminary geologic map of the Rainbow quadrangle, Uintah County, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-893, scale 1:24,000.
- Kemmerer, J.L., Jr., 1934, Gilsonite: Salt Lake City, University of Utah, M.S. thesis, 61 p.
- Kilborn, G.R., 1964, New methods of mining and refining gilsonite, in Sabatka, E.F., editor, Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse: Intermountain Association of Petroleum Geologists, 13th annual field conference, p. 247–252.
- King, L.H., Goodspeed, F.E., and Montgomery, D.S., 1963, A study of sedimented organic matter and its natural derivatives: Ottawa, Canada Department of Mines and Technical Survey, Mines Branch Research Report R 114, 68 p.
- Kretchman, H.F., 1957, The story of gilsonite: Salt Lake City, Utah, American Gilsonite Company, 96 p.
- Ladoo, R.B., and Myers, W.M., 1951, Nonmetallic minerals, 2nd edition: New York, McGraw-Hill Book Company, Inc., p. 61–66.
- Lewis, H.D., 1994, Gilsonite, in Carr, D.D., editor, Industrial minerals and rocks, 6th edition: Littleton, Colorado, Society for Mining, Metallurgy and Exploration, Inc., p. 535–541.
- McGee, L.R., 1956, Porphyryns in gilsonite: Salt Lake City, University of Utah, Ph.D. thesis, 53 p.
- Monson, B., and Parnell, J., 1992a, The origin of gilsonite vein deposits in the Uinta Basin, Utah: Utah Geological Survey Contract Report 92-4, 20 p.
- Monson, B., and Parnell, J., 1992b, The origin of Gilsonite vein deposits in the Uinta Basin, Utah, in Fouch, T.D., Nuccio, V.F., and Chidsey, T.C., Jr., editors, Hydrocarbon and mineral resources of the Uinta Basin, Utah and Colorado: Salt Lake City, Utah Geological Association

- Guidebook 20, p. 257–270.
- Neel, K.R., 1980, Gilsonite, in Grayson, M., executive editor, Kirk-Othmer encyclopedia of chemical technology (3rd edition), v. 11: New York, John Wiley and Sons, Inc.
- Pruitt, R.G., Jr., 1961, The mineral resources of Uintah County: Utah Geological and Mineralogical Survey Bulletin 71, 101 p.
- Remington, N.C., 1959, A history of the gilsonite industry: Salt Lake City, University of Utah, M.S. thesis, 338 p.
- Rogers, M.A., McAlary, J.D., and Bailey, N.J.L., 1974, Significance of reservoir bitumens to thermal maturation studies, western Canada basin: Bulletin of the American Association of Petroleum Geologists, v. 58, no. 9, p. 1806–1824.
- Romney, M.P., 1963, The nonmetallics, Utah's Cinderella minerals, in Cooley, E., editor, Utah, treasure house of the nation—century of mining, 1863–1963: Utah State Historical Society, Utah Historical Quarterly, v. 31, no. 3, p. 229–230.
- Shushan, L., 1983, Bonanza—the only place on earth where everybody knows what gilsonite is: Standard Oiler, September–October, p. 1–5.
- Sprinkel, D.A., 2007, Interim geologic map of the Vernal 30' x 60' quadrangle, Uintah and Duchesne Counties, Utah, and Moffat and Rio Blanco Counties, Colorado: Utah Geological Survey Open-File Report 506DM, GIS data, scale 1:100,000, CD.
- Sprinkel, D.A., 2009, Interim geologic map of the Seep Ridge 30' x 60' quadrangle, Uintah, Duchesne, and Carbon Counties, Utah, and Garfield and Rio Blanco Counties, Colorado: Utah Geological Survey Open-File Report 549DM, GIS data, scale 1:100,000, CD.
- Stone, D.S., 1993, Tectonic evolution of the Uinta Mountains—palinspastic restoration of a structural cross section along longitude 109°15', Utah: Utah Geological Survey Miscellaneous Publication 93-8, 19 p.
- Tripp, B.T., 2004, Gilsonite—an unusual Utah resource: Utah Geological Survey, Survey Notes, v. 36, no. 3, p. 1-3, 7.
- Tripp, B.T., and White, E.R., 2006, Gilsonite, in Kogel, J.E., Trivedi, N.C., Barker, J.M., and Krukowski, S.T., editors, Industrial minerals & rocks - commodities, markets, and uses: Littleton, Colorado, Society for Mining, Metallurgy, and Exploration, Inc., p. 481–493.
- U.S. Mine Safety and Health Administration, 2008, Data retrieval system: Online, www.msha.gov/drs/drshome.htm, accessed April 2008.
- Utah Automated Geographic Reference Center, 2008, Utah GIS Portal: Online, <http://gis.utah.gov/naip2006>, accessed 2008.
- Utah Mining Association, 1955, Utah's mining industry—an historical, operational, and economic review of Utah's mining industry, 1st edition: Salt Lake City, Utah, p. 24–79.
- Utah Mining Association, 1959, Utah's mining industry—an historical, operational, and economic review of Utah's mining industry, 2nd edition: Salt Lake City, Utah, 132 p.
- Utah Mining Association, 1967, An historical, operational and economic review of Utah's mining industry, 3rd edition: Salt Lake City, Utah, 135 p.
- Verbeek, E.R., and Grout, M.A., 1992, Structural evolution of gilsonite dikes, eastern Uinta Basin, Utah, in Fouch, T.D., Nuccio, V.F., and Chidsey, T.C., Jr., editors, Hydrocarbon and mineral resources of the Uinta Basin, Utah and Colorado: Salt Lake City, Utah Geological Association Guidebook 20, p. 237–255.
- Verbeek, E.R., and Grout, M.A., 1993, Geometry and structural evolution of gilsonite dikes in the eastern Uinta Basin, Utah: U.S. Geological Survey Bulletin 1787-HH, 42 p., scale 1:250,000.
- Weiss, M.P., Witkind I.J., and Cashion W.B., 2003 (digital release), Geologic map of the Price 30' x 60' quadrangle, Carbon, Duchesne, Uintah, Utah, and Wasatch Counties, Utah (digitized from U.S. Geological Survey Miscellaneous Investigations Series Map I-1981 [1990]): Utah Geological Survey Map 198DM, CD, scale 1:100,000.
- Wen, C.S., Chilingarian, G.V., and Yen, T.F., 1978, Properties and structure of bitumens, in Chilingarian, G.V., and Yen, T.F., editors, Bitumens, asphalts, and tar sands: Amsterdam, Elsevier Science Publishing Company, p. 155–190.
- Wurtz, H., 1869, On the grahamite of West Virginia, and the new Colorado resinoid: American Association for the Advancement of Science Proceedings 18, p. 124–125.
- Ziegler Chemical & Mineral Corporation, 2008, Home page Online, <http://www.zieglerchemical.com>, accessed April 2008.

SUPPLEMENTAL BIBLIOGRAPHY

- Aizenshtat, Z., Miloslavski I., and Tannenbaum, E., 1986, Thermal behavior of immature asphalts and related kerogens: Organic Geochemistry, v. 10, no. 1 3, p. 537–546.
- Alturki, Y.I.A., Eglinton, G., and Pillinger, C.T., 1972, The petroporphyrins of gilsonite: Pergamon Press, Oxford, United Kingdom (GBR) Advances in Organic Geochemistry, International Series of Monographs on Earth Sciences, v. 33, p. 135–150.
- American Gilsonite Company, no date, Gilsonite: Salt Lake City, Utah, American Gilsonite Company brochure, 8 p.

- American Gilsonite Company, no date, Pipe insulation news from Gilsulate: Salt Lake City, Utah, American Gilsonite Company quarterly periodical published from 1958 to 1964.
- American Gilsonite Company, 1961, Gilsonite guidebook: Salt Lake City, Utah, American Gilsonite Company, 23 p.
- American Gilsonite Company, 1969, Gilsonite guidebook: Salt Lake City, Utah, American Gilsonite Company, 22 p.
- American Gilsonite Company, 1979, Gilsonite: Salt Lake City, Utah, American Gilsonite Company, 16 p.
- Anders, D.E., 1990, Thermal maturation in the Uinta Basin, Utah, in Carter, L.M.H., editor, U.S. Geological Survey research on energy resources—1990 program and abstracts, Sixth V.E. McKelvey Forum on Mineral and Energy Resources: U.S. Geological Survey Circular 1060, p. 2–3.
- Anders, D.E., and Gerrild, P.M., 1984, Hydrocarbon generation in lacustrine rocks of Tertiary age, Uinta Basin, Utah—organic carbon, pyrolysis yields and light hydrocarbons, in Woodward, J., Meissner, F.F., and Clayton, J.L., editors, Hydrocarbon source rocks of the greater Rocky Mountain region: Denver, Rocky Mountain Association of Geologists, p. 513–529.
- Anders, D.E., Palacas, J.G., and Johnson, R.C., 1992, Thermal maturity of rocks and hydrocarbon deposits, Uinta Basin, Utah, in Fouch, T.D., Nuccio, V.F., and Chidsey, Jr., T.C., editors, Hydrocarbon and mineral resources of the Uinta Basin, Utah and Colorado: Utah Geological Association Guidebook 20, p. 53–76.
- Anonymous, 1897, The Uinta and the Uncompahgre asphaltites of Utah: Engineering and Mining Journal 64, p. 10–11.
- Anonymous, 1905, Gilsonite: New York, McGraw Hill, Engineering and Mining Journal, v. 80, p. 608.
- Anonymous, 1945, The evolution of gilsonite transportation: Mining and Contracting Review, v. 47, no. 1, p. 5.
- Anonymous, 1946, Business Week (staff article), Develop gilsonite: Business Week no. 854, Jan. 12, 67 p.
- Anonymous, 1956, Industry, solid source of gasoline: Chemical Engineering News, v. 34, no. 30, p. 3546.
- Anonymous, 1956, Gilsonite: Mining Engineering News, v. 8, no. 18, p. 785–790.
- Arnold, A., and Potts, K., 1952, Gilsonite: Utah Economic and Business Review, v. 12, no. 6, p. 4–5.
- Aurand, H.A., 1920, Mineral deposits of the western slope: Colorado Geological Survey Bulletin 22, p. 15–19.
- Bakhshandeh, F., 1976, A study of the aromatic fraction of oil shales and other carbonaceous deposits from the Green River Formation in Utah: Golden, Colorado School of Mines, M.S. thesis, 68 p.
- Ball, J.O., 1944, Survey of bitumen analyses and extraction methods, in Hydrocarbons of the Uinta Basin of Utah and Colorado: Golden, Quarterly of the Colorado School of Mines, v. 39, no. 1, p. 71–115.
- Barb, C.F., 1942, Rubber from the Uinta Basin of Utah (investigation of bituminous sands): Mines Magazine, 32, p. 521–524.
- Barb, C.F., 1944, Review of geology and field work, in Hydrocarbons of the Uinta Basin of Utah and Colorado: Golden, Quarterly of the Colorado School of Mines, v. 39, no. 1, p. 7–65.
- Bardwell, C., 1913, Chemical properties of Utah hydrocarbons: Salt Lake City, University of Utah, B.S. thesis, 221 p.
- Bardwell, C., Berryman, B.A., Brighton, T.B., and Kuhre, K.D., 1913, The hydrocarbons of Utah: Industrial Engineering Chemistry, v. 5, issue 12, p. 973–976.
- Bardwell, C., and others, 1918, Chemical properties of Utah hydrocarbons: Transactions of the Utah Academy of Science, p. 78–95.
- Bereskin, S.R., and Richards, D.R., 1984, Structural control and fractured reservoirs in relation to oil production from Green River Formation, Pleasant Valley–Monument Butte fields, Uinta Basin, Utah: American Association of Petroleum Geologists Bulletin, v. 68, no. 7, p. 932.
- Berkey, C.P., 1904, Mineral resources of the Uinta Mountains: The Engineering and Mining Journal, v. 77, p. 841.
- Bezzant, H.A., 1949, The chromatographic separation of gilsonite: Provo, Utah, Brigham Young University, M.S. thesis, 60 p.
- Blake, W.P., 1889, Wurtzilite: The Engineering and Mining Journal, v. 48, p. 542–543.
- Bon, R.L., Gloyd, R.W., and Tabet, D.E., 1997, Summary of mineral activity in Utah: Utah Geological Survey Circular 98, 12 p.
- Bostwick, J.M., (revised by Bradbury, J.C.), 1983, Bituminous materials, in Lafond, S.J., editor-in-chief, Industrial minerals and rocks: New York, Society of Mining Engineers of the American Institute of Mining, Metallurgy, and Petroleum Engineers, Inc., p. 529–532.
- Breger, I.A., 1963, Origin and classification of naturally occurring carbonaceous substances, in Breger, I.A. editor, Organic geochemistry: New York, The Macmillan Company, and Oxford, Pergamon Press, Earth Science Series, Monograph 16, p. 50–86.
- Brewster, B., 1945a, Gilsonite mining goes modern: Mining and Contracting Review, v. 47, no. 5, p. 6–8.
- Brewster, B., 1945b, The evolution of gilsonite transport-

- tation: *Mining and Contracting Review*, v. 47, no. 1, p. 5–6.
- Bristol, J., 1929, Gilsonite deposits of the Uinta Basin: Phoenix, Arizona, Miller Freeman Publications, *Mining Journal*, v. 13, no. 3, p. 5–6 and 63–64.
- Buraneck, A.M., and Needham, C.E., 1949, Directory of the Utah mineral resources and consumers guide: Utah Department of Publicity and Industrial Development (now Utah Geological Survey) Bulletin 36, p. 15–16.
- Campbell, G.S., and Hebrew, Q., 1957, Oil and gas developments in Utah and Nevada in 1956: American Association of Petroleum Geologists Bulletin, v. 41, no. 6, p. 1245.
- Carman, E.P., and Bayes, F.S., 1961, Occurrences, properties, and uses of some natural bitumens: U.S. Bureau of Mines Information Circular 7997, 42 p.
- Cashdollar, K.L., and Chatrathi, K., 1993, Minimum explosive dust concentrations measured in 20-L and 1-M3 chambers: *Combustion Science and Technology*, v. 87, no. 1-6, p. 157–171.
- Cashion, W.B., 1973, Geologic and structure map of the Grand Junction quadrangle, Colorado and Utah: U.S. Geological Survey Miscellaneous Investigations Series Map I-736, scale 1:250,000.
- Cashion, W.B., 1982, Descriptions of four stratigraphic sections of parts of the Green River and Uinta Formations in the eastern Uinta Basin, Uintah County, Utah and Rio Blanco County, Colorado: U.S. Geological Survey Open-File Report 83-17, 42 p.
- Cashion, W.B., and Brown, J.H., 1956, Geology of the Bonanza-Dragon oil-shale area, Uintah County, Utah and Rio Blanco County, Colorado: U.S. Geological Survey Oil and Gas Investigations Map OM-153, scale 1:62,500.
- Cashion, W.B., and Donnell, J.R., 1974, Revision of nomenclature of the upper part of the Green River Formation Piceance Creek Basin, Colorado and eastern Uinta Basin, Utah: U.S. Geological Survey Bulletin 1394-G, 9 p.
- Chenoweth, W.L., 1985, The Uinta railway, in Picard, M.D., editor, *Geology and energy resources, Uinta Basin of Utah*: Utah Geological Association Guidebook 12, p. 17–18.
- Chesterman, C.W., and Main, F.H., 1947, Reconnaissance investigations for trace elements in Utah, Colorado, Nevada, California, and Oregon, preliminary report: U.S. Geological Survey, TEI 24, for U.S. Atomic Energy Commission, 45 p.
- Chilingarian, G.V., and Yen, T.F., editors, 1978, *Bitumens, asphalts, and tar sands*: New York, Elsevier Scientific Publishing Co., *Developments in Petroleum Science* 7, 331 p.
- Clark, A.F., 1924, *Utah hydrocarbons*: Salt Lake City, University of Utah, M.S. thesis, 59 p.
- Clark, A.F., and Newton, R.F., 1925, *Utah hydrocarbons*: Salt Lake City, Utah, Engineering Experiment Station Bulletin, v. 15, no. 2, p. 33–46.
- Clark, D.T., and Wilson, R., 1983, Some aspects of the surface chemistry of coal, kerogen and bitumen as revealed by ESCA: *Organic Geochemistry*, v. 6, p. 455–461.
- Clark, D.T., Wilson, R., and Quirke, J.M.E., 1983, An evaluation of the potential of ESCA (electron spectroscopy for chemical applications) (and other spectroscopic techniques) in the surface and bulk characterization of kerogens, brown coal and gilsonite: *Chemical Geology*, v. 39, no. 3-4, p. 215–239.
- Claypole, E.W., 1889, Gilsonite or uintahite: *The American Geologist*, v. 4, p. 386–387.
- Cohenour, R.E., 1966, Lands in T9, 10S, R20-21-22E, Salt Lake Meridian, Uinta Basin: Utah Geological and Mineralogical Survey Report of Investigation 27, 3 p.
- Crawford, A.L., 1941, Strategic minerals of Utah: *Mineral Society of Utah News Bulletin* 2, no. 2, p. 4–14.
- Crawford, A.L., 1942, Strategic minerals of Utah: *University of Utah Engineering Experiment Station Bulletin* 18, 19 p.
- Crawford, A.L., 1943, Utah gilsonite and related hydrocarbons: *Utah Industrial Development News*, July 1943, p. 7–8.
- Crawford, A.L., 1947, Gilsonite and related hydrocarbons from the Uinta Basin, Utah: *Mineralogical Society of Utah Bulletin* 7, no. 1, p. 44–48.
- Crawford, A.L., 1949, Gilsonite and related hydrocarbons of the Uinta Basin, Utah, in Hansen, G.H., and Bell, M.M., editors, *The oil and gas possibilities of Utah*: Utah Geological and Mineralogical Survey, p. 235–260.
- Crawford, A.L., 1959, Mineral potentials on approximately 30 townships for which Utah State Land Board selections are recommended: Utah Geological and Mineralogical Survey unpublished report, 22 p.
- Crawford, A.L., 1961, Utah raw materials of interest to the chemical engineer: Utah Geological and Mineralogical Survey Reprint 30, 2nd revision (reprinted from *Chemical and Engineering News*, American Chemical Society, 1949, v. 27, p. 3017).
- Cross, A.T., and Wood, G.D., 1976, Palynology and petrography of some solid bitumens of the Uinta Basin, Utah: *Brigham Young University Geology Studies*, v. 22, pt. 3, p. 157–173.
- Curiale, J.A., 1981a, Origin and geochemical correlation of near surface oil and asphaltite deposits of southeastern Oklahoma [abs.]: *American Association of Petroleum Geologists Bulletin*, v. 65, no. 5, p. 915–916.
- Curiale, J.A., 1981b, Source rock geochemistry and liq-

- uid and solid petroleum occurrences of the Ouachita Mountains, Oklahoma: Norman, University of Oklahoma, Ph.D. dissertation, 286 p.
- Curiale, J.A., 1983, Petroleum occurrences and source-rock potential of the Ouachita Mountains, southeastern Oklahoma: Oklahoma Geological Survey Bulletin 135, 65 p.
- Curiale, J.A., 1985, Origin of solid bitumens, with emphasis on biological marker results: *Organic Geochemistry*, v. 10, no. 1 3, p. 559–580.
- Curiale, J.A., 1987, Distribution and occurrence of metals in heavy crude oils and solid bitumens—implications for petroleum exploration, in Meyer, R.F., editor, *Exploration for heavy crude oil and natural bitumen: American Association of Petroleum Geologists Studies in Geology* 25, p. 207–219.
- Curiale, J.A., 1988, Biological markers in grahamites and pyrobitumens, in Yen, T.F., and Moldowan, J.M., editors, *Geochemical biomarkers: New York, Harwood Academic Publishers*, p. 1–24.
- Curiale, J.A., 1993, Occurrence and significance of metals in solid bitumens: an organic geochemical approach, in Parnell, J., Kucha, H., and Landais, P., editors, *Bitumens in ore deposits: New York, Springer-Verlag*, p. 461–474.
- Current, A.M., 1953a, Review of geology and activities in the Uinta Basin [Utah], in Moore, C.A., editor, *3rd Sub-surface Geological Symposium proceedings, March: Colorado School of Mines Quarterly*, v. 48, no. 3, p. 63–71.
- Current, A.M., 1953b, Review of geology and activities in the Uinta Basin: *Tulsa Geological Society Digest*, v. 21, p. 52–61; also *Colorado School of Mines Publication* 48, no. 3, 36 p.
- Current, A.M., 1954, It's rough going in the Uinta Basin: *World Oil*, v. 138, no. 5, p. 115–116, 118, 120, 122.
- Davis, H.C., 1952, *Geology of the Culmer gilsonite vein of Duchesne County, Utah: Provo, Utah, Brigham Young University, M.S. thesis*, 90 p.
- Davis, N., II, and Tooman, C.E., 1989, New laboratory tests evaluate the effectiveness of gilsonite as a borehole stabilizer: *Society of Petroleum Engineers of the American Institute of Mining, Metallurgical and Petroleum Engineers (AIME) Drilling Engineering*, v. 4, no. 1, p. 47–56.
- Day, W.C., 1895, Investigation of Utah gilsonite, a variety of asphalt: *Journal of the Franklin Institute*, v. 140, no. 837, p. 221–237.
- Denton, W., 1866, On a mineral, resembling albertite, from Colorado: *Boston Society of Natural History Proceedings*, v. 10, p. 305–306.
- Dewey, R.F., 1965, Gilsonite—hydraulic mining, tunnel boring, and shaft drilling operations, in Pope, B.J., Harry, J.V., and Lyon, L.B., editors, *Proceedings of the First International Symposium on Fossil Hydrocarbons: Brigham Young University Center for Continuing Education*, p. 98–113.
- Dickerson, R.P., Gaccetta, J.D., Kulik, D.M., and Kreidler, T., 1990, Mineral resources of the Coal Canyon, Spruce Canyon, and Flume Canyon Wilderness Study Areas, Grand County, Utah: *U.S. Geological Survey Bulletin* 1753-A, 25 p.
- Douglas, A.G., and Grantham, P.J., 1974, Fingerprint gas chromatography in the analysis of some native bitumens, asphalts and related substances, in Tissot, B. and Bienner, F., editors, *Advances in organic geochemistry 1973: Proceeding of the 6th International Congress on Organic Geochemistry*, p. 261–276.
- Douglass, E., 1921a, The oil problem in the Uinta Basin, Utah: *Salt Lake Mining Review*, v. 23, no.16, p. 12–13.
- Douglass, E., 1921b, Oil problems in the Uinta Basin, Utah: *Salt Lake Mining Review* 23, Nov. 30, p. 12-13; also Dec. 15, p. 14–15; Dec. 30, p. 9–11; Jan. 30, 1922, p. 13–14; Feb. 28, p. 9–11; Mar. 15, p. 11–14; Apr. 15, p. 17–18.
- Douglass, E., 1928, Final report on the gilsonite holdings of the Gilson Asphaltum Company in Utah and Colorado: Private report (as consultant) to the Gilson Asphaltum Company; manuscript on file in the Special Collections Department, University of Utah Libraries, Salt Lake City, Utah, 137 p.
- Eglinton, G., Hajibrahim, S.K., Maxwell, J.R., and Quirke, J.M.E., 1980, Petroporphyrins - structural elucidation and the application of HPLC fingerprinting to geochemical problems: *Physics and Chemistry of the Earth*, v. 12, p. 193–203.
- Eldridge, G.H., 1896a, Occurrence of uintahite in Utah: *Science*, v. 3 (new series), no. 75, p. 830–832.
- Eldridge, G.H., 1896b, The uintahite (gilsonite) deposits of Utah: *U.S. Geological Survey 17th Annual Report*, pt. 1, no. 2, 41 p.
- Eldridge, G.H., 1903, Origin and distribution of asphalt and bituminous rock deposits in the United States: *U.S. Geological Survey Bulletin* 213, p. 296–305.
- Eldridge, G.H., 1906, The formation of asphalt veins: *Economic Geology*, v. 1, p. 437–444.
- Elmore, R.D., 1987, Evidence for a relationship between hydrocarbons and secondary magnetizations that reside in authigenic magnetite, *International Union of Geodesy and Geophysics (IUGG), XIX General Assembly, Vancouver, BC, Canada, Aug. 9–22, 1987: International Union of Geodesy and Geophysics*, v. 19, p. 492.
- Elmore, R.D., Engel, M.H., Crawford, L., Nick, K., Imbus, S., and Sofer, S., 1987, Evidence for a relationship between hydrocarbons and authigenic magnetite: *Nature*, v. 325, no. 6103, p. 428–430.

- Epler, B., 1990, Uintah railway—crucial link for early-day gilsonite mines: *Pay Dirt*, July, p. 7b–8b.
- Farmer, V.E., Jr., 1952, Rocky Mountain "hot spot," Uinta Basin: *Oil and Gas Journal*, v. 51, no. 3, p. 199–201, 203–204.
- Fene, W.J., 1928, The mining of gilsonite in Utah: U.S. Bureau of Mines Information Circular 6069, 6 p.
- Fiack, H., 1929, Fort Duchesne's beginnings: *Utah Historical Quarterly*, v. 2, p. 31.
- Fleming, R.C., 1932, Mining gilsonite in Utah: *Mining and Metallurgy*, v. 13, no. 312, sec. 1, p. 529–530.
- Fouch, T.D., 1985, Oil- and gas-bearing Upper Cretaceous and Paleogene fluvial rocks in central and northeast Utah, in Flores, R.M., Galloway, W.E., Ethridge, F.G., Miall A.D., and Fouch, T.D., editors, Recognition of fluvial depositional systems and their resource potential: Society of Economic Paleontologists and Mineralogists Short Course 19, p. 241–271.
- Fouch, T.D., Nuccio, V.F., Anders, D.E., Rice, D.D., Pitman, J.K., and Mast, R.F., 1994, Green River(!) petroleum system, Uinta Basin, Utah, U.S.A., in Magoon, L.B., and Dow, W.G., editors, The petroleum system—from source to trap: American Association of Petroleum Geologists Memoir 60, p. 399–421.
- Franczyk, K.J., Pitman, J.K., Cashion, W.B., Chan, M.A., Donnell, J.R., Dyni, J.R., Fouch, T.D., Johnson, R.C., Lawton, T.F., and Remy, R.R., 1989, Evolution of resource-rich foreland and intermontane basins in eastern Utah and western Colorado: Washington, D.C., American Geophysical Union, 28th International Geological Congress, Field Trip Guidebook T-328, 53 p.
- Gill, J.P., Evershed, R.P., Chicarelli, M.I., Wolff, G.A., Maxwell, J.R., and Eglinton, G., 1985, Computerised capillary gas chromatographic mass spectrometric studies of the petroporphyrins of the gilsonite bitumen (Eocene, U.S.A.): *Journal of Chromatography*, v. 350, no. 1, p. 37–62.
- Gomez, V., 1994, Response to gilsonite plant blaze draws criticism: *Salt Lake Tribune*, October 4, p. C14.
- Guillotte, G.B., 1944, SOM (Uranium) in the hydrocarbons of the Uinta Basin of Utah and Colorado: U.S. Atomic Energy Commission, RMO-137, 21 p.
- Hager, D., 1952, Oil and gas in the Uinta Basin of Utah: *Petroleum Engineering*, v. 24, no. 3, p. A41–A45; also *Mines Magazine*, 1953, v. 43, no. 10, p. 81–83.
- Hajibrahim, S.K., Quirke, J.M.E., and Eglinton, G., 1981, Petroporphyrins—V, Structurally related porphyrin in bitumens, shales and petroleum—evidence from HPLC and mass spectrometry: *Chemical Geology*, v. 32, no. 3–4, p. 173–188.
- Harmer, E.W., 1939, Gilsonite mining in Utah expands: *Mining Journal (Arizona)*, v. 23, no. 7, p. 6–7.
- Harrison, A.G., and Thode, H.G., 1958, Sulphur isotope abundances in hydrocarbons and source rocks of Uinta Basin, Utah: *American Association of Petroleum Geologists Bulletin*, v. 42, no. 11, p. 2642–2649; also *Utah Geological and Mineralogical Survey*, 1959, Reprint 71.
- Hartzell, W.F., 1928, Natural solidified petroleum—its mining, treatment, and uses: *Engineering and Mining Journal*, v. 125, no. 6, p. 253–254.
- Hayes, A.A., 1866, Description and analysis of a new kind of bitumen: *Boston Society of Natural History Proceedings*, v. 10, p. 306–307.
- Hodgson, R.A., 1992, The Duchesne lineament—Utah and Colorado, in Mason, R., editor, *Proceedings of the 7th International Conference on Basement Tectonics*, Kingston, Ontario, 1987: Dordrecht, The Netherlands, Kluwer Academic Publishers, p. 353–362.
- Holten, R., 1994, Dayton streets get brick look without brick cost: *American City & County*, v. 109 (December 1994), p. 26.
- Hosterman, J.W., Meyers, R.F., Palmer, C.A., Doughten, M.W., and Anders, D.E., 1990, Chemistry and mineralogy of natural bitumens and heavy oils and their reservoir rocks from the United States, Canada, Trinidad and Tobago, and Venezuela: U.S. Geological Survey Circular 1047, 19 p.
- Hunt, J.M., 1978, Characterization of bitumens and coals: *American Association of Petroleum Geologists Bulletin*, v. 62, no. 2, p. 301–303.
- Hwang, R.J., Teerman, S.C., and Carlson, R.M., 1998, Geochemical comparison of reservoir solid bitumens with diverse origins: *Organic Geochemistry*, v. 29, p. 505–517.
- Jackson, D., 1981, American Gilsonite—mining solid hydrocarbon: *Engineering and Mining Journal*, v. 182, no. 7, p. 88–91.
- Jacob, H., 1985, Dispersed solid bitumens as an indicator for migration and maturity in prospecting for oil and gas: *Erdol & Kohle Erdgas Petrochemie*, v. 38, p. 365.
- Jacob, H., 1989, Classification, structure, genesis and practical importance of natural solid oil bitumen ("migrabitumen"): *International Journal of Coal Geology*, v. 11, no. 1, p. 65–79.
- Jacob, H., 1993, Nomenclature, classification, characterization, and genesis of natural solid bitumen (migrabitumen), in Parnell, J., Kucha, H., and Landais, P., editors, *Bitumens in ore deposits*: New York, Springer-Verlag, p. 11–27.
- Jacob, H., and Hiltmann, W., 1988, Disperse, feste Erdolbitumina als Maturitäts Indikatoren im Rahmen der Erdol-/Erdgas Prospektion: *Geologisches Jahrbuch, Reihe D*, 89, p. 3–37.

- Jacob, H., and Wehner, H., 1981, Mikroskopphotometrische analyse disperser festbitumina in sedimenten: Deutsche Gesellschaft für Mineralogische Wissenschaft und Kohlechemie, DGMK-Projekt 232, 257 p.
- Jiyang, S., Benshan, W., Luje, Z., and ZhiQuing, H., 1988, Study on diagenesis of organic matter in immature rocks, in Mattavelli, L., and Novelli, L., editors, *Advances in organic geochemistry 1987: Great Britain*, Pergamon Press, *Organic Geochemistry*, v. 14, no. 4-6, p. 867-874.
- Johnson, K.S., 1980, Energy fuels field course and workshop—exploring the Colorado Plateau and the Rocky Mountain area of Colorado and Utah, Grand Junction, Colorado, United States, July 27–Aug. 16: Norman, University of Oklahoma.
- Johnson, R.C., and Nuccio, V.F., 1993, Surface vitrinite reflectance study of the Uinta and Piceance basins and adjacent areas, eastern Utah and western Colorado—implications for the development of Laramide basins and uplifts: *U.S. Geological Survey Bulletin* 1787-DD, 38 p.
- Jones, F.R., 1939, *Gilsonite: Chemistry and Industry*, v. 58, no. 3, p. 800-801.
- Kelly, F.J., Kerns, W.H., Parker, B., and Ransome, A.L., 1959, *The mineral industry of Utah, 1955: Utah Geological and Mineralogical Survey Reprint* 61, 36 p.
- Khavari Khorasani, G., 1984, Free hydrocarbons in Uinta Basin, Utah: *American Association of Petroleum Geologists Bulletin*, v. 68, no. 9, p. 1193-1197.
- Khavari-Khorasani, G., and Michelsen, J.K., 1993, The thermal evolution of solid bitumens, bitumen reflectance, and kinetic modeling of reflectance—application in petroleum and ore prospecting, in Goodarzi, F., and Macqueen, R.W., editors, *Geochemistry and petrology of bitumen with respect to hydrocarbon generation and mineralization: Energy Sources*, v. 15, p. 181-204.
- Kirkpatrick, S.D., 1928, Marketing the natural hydrocarbons—how gilsonite, wurtzilite, elaterite, manjak and grahamite and the mineral waxes, ozokerite and cersine, are sold: *Engineering and Mining Journal Press*, v. 119, no. 8, 331 p.
- Kullman, G.J., Doak, C.B., and Keimig, D.G., 1989, Assessment of respiratory exposures during gilsonite mining and milling operations: *American Industrial Hygiene Association Journal*, v. 50 (August 1989), p. 413-418.
- Ladoo, R.B., 1920, The natural hydrocarbons—gilsonite, elaterite, wurtzilite, grahamite, ozokerite, and others: *U.S. Bureau of Mines Report of Investigation* 2121, 12 p.
- Lakes, A., 1909, Hydrocarbons in the United States— asphalt, gilsonite, and other hydrocarbons, their distribution, modes of occurrence, and methods used in mining them: *Mining Science*, v. 60, p. 340-342.
- Landis, C.R., and Castano, J.R., 1994, Maturation and bulk chemical properties of a suite of solid hydrocarbons: *Organic Geochemistry*, v. 22, no. 1, p. 137-149.
- Langston, H.M., 1929, Gilsonite and related bitumens: *Industrial Chemistry and Chemical Engineering Magazine*, v. 5, p. 324-326, 383-386.
- Lewis, R.Q., Sr., and Campbell, R.H., 1955, Elk Ridge area, Utah: *U.S. Geological Survey Trace Elements Investigations Report* TEI 590, p. 54-55.
- Lewis, R.S., and Varley, T., 1931, The mineral industry of Utah—1931: *U.S. Geological Survey Mineral Yearbook*.
- Lexco, Inc., 2008, Home page: Online, www.gilsonite.com/, accessed April 2008.
- Locke, J.M., 1887, Gilsonite or uintahite, a new variety of asphaltum from the Uintah Mountains, Utah: *American Institute of Mining Engineers Transactions* 16, p. 162-168.
- Lomando, A.J., 1992, The influence of solid reservoir bitumen on reservoir quality: *American Association of Petroleum Geologists Bulletin*, v. 76, p. 1137-1152.
- Luff, P., 1955, *The mineral industry of Utah, 1952: Utah Geological and Mineralogical Survey Reprint* 51.
- Maguire, D., 1900, The hydrocarbons of eastern Utah, with special reference to the deposits of ozokerite, gilsonite, and elaterite: *Scranton, Pennsylvania, Mines and Minerals*, v. 20, no. 9, p. 398-400.
- McCullough, T.F., 1955, Hydrocarbons and other compounds obtained from gilsonite: Salt Lake City, University of Utah, Ph.D. dissertation, 54 p.
- Marrow, J., 1957, Ozokerite at Soldier Summit, Utah, in Seal, O.G. editor, *Geology of the Uinta Basin: Intermountain Association of Petroleum Geologists Guidebook*, 8th Annual Field Conference, p. 161-164.
- Meuzelaar, H.L.C., Haverkamp, J., and Hileman, F.D., 1982, Curie-point pyrolysis mass spectrometry of recent and fossil biomaterials, in Meuzelaar, H.L.C., and Hileman, F.D., editors, *Compendium and atlas: Amsterdam, Elsevier Publishers*, 293 p.
- Meyer, R.F., and De Witt, W., Jr., 1990, Definition and world resources of natural bitumens: *U.S. Geological Survey Bulletin* 1944, 14 p.
- Miller, J.S., Jr., 1938, Native asphalts and bitumens—the science of petroleum, v. 4: London, Oxford University Press, p. 2710-2727.
- Murray, A.N., 1949, The gilsonite deposits of the Uinta Basin, Utah: *Tulsa Geological Society Digest*, v. 17, p. 104-106.
- Murray, A.N., 1950, The gilsonite deposits of the Uinta Basin, Utah, in Eardley, A.J., editor, *Petroleum geology of the Uinta Basin: Intermountain Association of Petroleum Geologists Guidebook to the Geology of*

- Utah No. 5, p. 115–118.
- Nackowski, M.P., Fisher, D., and Beer, L., 1963, Mineral resources of Duchesne County: Utah University Engineering Experiment Station Bulletin 125 (Bulletin of the University of Utah, v. 54, no. 20), 97 p.
- Needham, C.E., 1948, Utah's hundred years of mineral production [abs.]: Utah Academy of Sciences, Arts, and Letters Proceedings 25, p. 185.
- Neel, K.R., 1990, Over the years, gilsonite markets and applications evolve: Pay Dirt Magazine, July, p. 6B.
- Neilson, D.L., Chidsey, T.C., Jr., Morgan, C., and Wenzhi, Z., 1993, Fracturing in the Duchesne field, Utah—importance for horizontal drilling [abs.]: American Association of Petroleum Geologists 1993 annual convention, New Orleans, Louisiana, April 25–28, 1993, American Association of Petroleum Geologists and Society of Economic Paleontologists and Mineralogists, v. 1993, p. 156–157.
- Nelson, R.E., 1965, Gilsonite, in Pope, B.J., Harry, J.V., and Lyon, L.B., editors, Proceedings of the First International Symposium on Fossil Hydrocarbons: Provo, Utah, Brigham Young University Center for Continuing Education, p. 355–357.
- Newberry, J.S., 1887, Grahamite in Colorado: Golden, Colorado School of Mines Quarterly, v. 8, p. 327–335.
- Osmond, J.C., 1992, Greater Natural Buttes gas field, Uintah County, Utah, in Fouch, T.D., Nuccio, V.F., and Chidsey, T.C., Jr., editors, Hydrocarbon and mineral resources of the Uinta Basin, Utah and Colorado: Utah Geological Association Publication 20, p. 53–76.
- O'Sullivan, R.B., and Ging, T.G., 1987, Preliminary report on solid bitumens in Eocene rocks of Piceance Creek basin, northwestern Colorado: U.S. Geological Survey Open-File Report 87-0478, 13 p.
- Palacas, J.G., editor, 1984, Petroleum geochemistry and source rock potential of carbonate rocks: American Association of Petroleum Geologists Studies in Geology 18, 208 p.
- Palacas, J.G., Anders, D.E., King, J.K., and Lubeck, C.M., 1989, Use of biological markers in determining thermal maturity of biodegraded heavy oils and solid bitumens, in Meyer, R.F., and Wiggins, E.J., editors, International Conference on Heavy Crude and Tar Sands: Fourth UNITAR/UNDP International Conference on Heavy Crude and Tar Sands, v. 2, p. 575–592.
- Parnell, J., Carey, P.F., and Monson, B., 1996, Fluid inclusion constraints on temperatures of petroleum migration from authigenic quartz in bitumen veins: Chemical Geology, v. 129, no. 3–4, p. 217–226.
- Picard, M.D., 2002, Searching for gilsonite when I was young: Journal of Geoscience Education, v. 50, no. 4, p. 471–474.
- Pitman, J.K., Fouch, T.D., and Goldhaber, M.B., 1982, Depositional setting and diagenetic evolution of some Tertiary unconventional reservoir rocks, Uinta Basin, Utah: American Association of Petroleum Geologists Bulletin, v. 66, no. 10, p. 1581–1596.
- Raymond, R.W., 1889, Note on a specimen of gilsonite from Uintah County, Utah: Transactions of the American Institute of Mining Engineers, v. 17, p. 113–115.
- Redfield, A.H., 1937, Native bitumens, in Dolbear, S.H., and Bowles, O., editors, Industrial minerals and rocks (nonmetallics other than fuels), first edition: New York, The American Institute of Mining and Metallurgical Engineers, p. 527–532.
- Redfield, A.H., 1949, Native bitumens, in Dolbear, S.H., and Bowles, O., editors, Industrial minerals and rocks (nonmetallics other than fuels), second edition: New York, The American Institute of Mining and Metallurgical Engineers, p. 527–532.
- Reusser, R.E., 1949, An investigation of the organic bases in gilsonite: Salt Lake City, University of Utah, M.S. thesis, 37 p.
- Richardson, C., 1916, Gilsonite and grahamite, the result of metamorphism of petroleum under a particular environment: Journal of Industrial and Engineering Chemistry, v. 8, p. 493–494.
- Ritzma, H.R., 1974, Publications on hydrocarbons by the Utah Geological and Mineral Survey: Utah Geological and Mineralogical Survey Circular 56, 5 p.
- Rowley, P.D., Hansen, W.R., Tweto, O., and Carrara, P.E., 1985, Geologic map of the Vernal 1° x 2° quadrangle, Colorado, Utah, and Wyoming: U.S. Geological Survey Miscellaneous Investigations Series Map I-1526, scale 1:250,000.
- Ruble, T.E., 1990, Organic geochemical investigation of native bitumens from the Uinta Basin, Utah, U.S.A.: Norman, University of Oklahoma, M.S. thesis, 155 p.
- Ruble, T.E., Bakel, A.J., and Philp, R.P., 1994, Compound-specific isotopic variability in Uinta Basin native bitumens—paleoenvironmental implications, in Schoell, M., and Hayes, J.M., editors, Compound specific isotope analysis in biogeochemistry and petroleum research—proceedings of a symposium held at the spring 1992 national meeting of the American Chemical Society, San Francisco: Organic Geochemistry, v. 21, no. 6–7, p. 661–671.
- Ruble, T.E., Lewan, M.D., and Cardott, B.J., 1999, Uinta basin solid hydrocarbons—new data and new insights: TSOP Newsletter, v. 16, no. 3, p. 13–16.
- Ruble, T.E., and Philp, R.P., 1989, Geochemical characterization of some native bitumens from the Uinta Basin, Utah, U.S.A. [abs.]: Geological Society of America Abstracts with Programs, v. 21, no. 6, p. 6.
- Ruble, T.E., and Philp, R.P., 1991, Geochemical investiga-

- tion of native bitumens from the Uinta Basin, Utah, U.S.A.: The Compass of Sigma Gamma Epsilon, v. 68, no. 3, p. 135-150.
- Ruble, T.E., and Philp, R.P., 1992, A reevaluation of the geochemical characteristics of solid bitumens from the Ouachita Mountains, Oklahoma: Oklahoma Geological Survey Circular 93, p. 337-342.
- Ruble, T.E., Philp, R.P., and Bakel, A.J., 1992, Compound specific isotopic analysis of Uinta Basin native bitumens [abs.]: 203rd American Chemical Society national meeting, San Francisco, California, April 5-10, 1992, Abstracts of Papers, v. 203, p. 138.
- Sanford, S., and Stone, R.W., 1914, Useful minerals of the U.S.: U.S. Geological Survey Bulletin 585, p. 184-188.
- Schoell, M., Hwang, R.J., Carlson, R.M.K., and Welton, J.E., 1994, Carbon isotopic composition of individual biomarkers in gilsonites, in Schoell, M., and Hayes, J.M., editors, Compound specific isotope analysis in biogeochemistry and petroleum research—proceedings of a symposium held at the spring 1992 national meeting of the American Chemical Society, San Francisco: Organic Geochemistry, v. 21, no. 6-7, p. 673-683.
- Schrader, F.C., Stone, R.W., and Stanford, S., 1917, Useful minerals of the United States: U.S. Geological Survey Bulletin 624, 412 p.
- Schufle, J.A., 1963, Minerals and energy sources in the arid west, in Hodge, C. editor, Aridity and man—the challenge of the arid west in the United States: American Association for the Advancement of Science Publication 74, p. 173-213.
- Scott, R.W., and Pantea, M.P., 1985, Preliminary geologic map of the Dragon quadrangle, Uintah County, Utah, and Rio Blanco County, Colorado: U.S. Geological Survey Miscellaneous Field Investigations Map MF-1774, scale 1:24,000.
- Seegmiller, B.L., and Willson, J.E., 1968, Mining of fossil hydrocarbons: Chemical Engineering Progress Symposium Series, v. 64, no. 85, p. 51-56.
- Selker, A.H., Scott, A.H., and McPherson, A.T., 1943, Electrical and mechanical properties of the system bunsen-gilsonite: Journal of Research, National Bureau of Standards, v. 31, p. 141-161.
- Slagle, K.A., and Carter, L.G., 1959, Gilsonite—a unique additive for oil-well cements—Paper presented at the spring meeting of the Pacific Coast district, Division of Production, American Petroleum Institute, April 30-May 1.
- Smith, D.K., and Grant, H., 1989, Gilsonite leads fight against lost circulation: Petroleum Engineer International, v. 61, p. 42-46.
- Smith, W.S., 1905, Mineral resources of Uinta Reservation, Utah: Mining World, v. 23, p. 491-492.
- Sorensen, D.P., 1955, Acidic and basic compounds in Gilsonite: Salt Lake City, University of Utah, Ph.D. dissertation, 68 p.
- Sprinkel, D.A., 1999, Digital geologic resources atlas of Utah: Utah Geological Survey Bulletin 129DF, CD.
- Standard Oiler, 1983, The saga of Sam Gilson—from Pony Express rider to mining pioneer: Standard Oiler, September-October, p. 4-5.
- Stern, K., 1960, Native bitumens, pyrobitumens, and asphaltic type petroleum bitumens, in Gillson, J.J., editor-in-chief, Industrial minerals and rocks (non-metallics other than fuels), 3rd edition: New York, The American Institute of Mining, Metallurgy, and Petroleum Engineers, p. 631-637.
- Stone, G.H., 1891, Note on the asphaltum of Utah and Colorado: American Journal of Science, 3rd series, v. 42, no. 248, p. 148-159.
- Sweeney, J.J., Burnham, A.K., and Braun, R.L., 1987, A model of hydrocarbon generation from type I kerogen—application to Uinta Basin, Utah: American Association of Petroleum Geologists Bulletin, v. 71, no. 8, p. 967-985.
- Taff, J.A., and Smith, C.D., 1906, Asphalt, related bitumens, and bituminous rock, in Contributions to economic geology, 1905: U.S. Geological Survey Bulletin 285-H, 373 p.
- Taylor, J., 1928, Gilsonite deposits of the Uinta basin: Report of the Utah Industrial Commission (1926-1928), p. 42.
- Tetting, T.N., 1984, Origin of gilsonite fractures in Uinta Basin, Utah: American Association of Petroleum Geologists Bulletin, v. 68, no. 7, p. 951-952.
- Thomas, C.R., Huddle, J.W., and Bass, N.W., 1946, Rangely oil field, Rio Blanco County, Colorado: Bulletin of the American Association of Petroleum Geologists, v. 30, no. 5, p. 749-750.
- Untermann, G.E., and Untermann, B.R., 1964, Geology of Uintah County: Utah Geological and Mineralogical Survey Bulletin 72, 112 p.
- U.S. Bureau of Land Management, 1986, Draft environmental assessment, Gilsonite applications, Vernal District, Vernal, Utah: Environmental Assessment 1986-77, 65 p.
- U.S. Bureau of Land Management, 1999, Environmental assessment for American Gilsonite Company's Wagon Hound gilsonite mine shafts nos. 3 and 7, Uintah County, Utah: U.S. Department of Interior, Bureau of Land Management.
- U.S. Bureau of Mines, no date, Mineral resources of the United States, 1904-1932: U.S. Department of Commerce, Bureau of Mines.
- Utah Geological and Mineralogical Survey, 1975, Energy resources map of Utah: Utah Geological and Mineral-

- ogical Survey Map 36, scale 1:500,000.
- Utah Geological and Mineral Survey, 1977, Energy resources map of Utah: Utah Geological and Mineral Survey Map 44, scale 1:500,000.
- Vanderwilt, J.W., 1947, Metals, nonmetals, and fuels - mineral resources of Colorado: Colorado Mineral Resources Board, p. 258–259.
- Warner, M.M., 1963, Sedimentation of the Duchesne River Formation, Uinta Basin, Utah: Iowa City, State University of Iowa, Ph.D. dissertation, 493 p.
- Wavrek, D.A., Jarvie, D.M., and Burgess, J.D., 1999, Characterization of solid reservoir bitumen—insights to formation mechanism, timing, and correlation [abs.]: TSOP Abstracts and Program, v. 16, p. 7–10.
- Wells, L.F., 1958, Petroleum occurrence in the Uinta Basin: Habitat of Oil, a symposium of the American Association of Petroleum Geologists, p. 344–365.
- Wenger, W.J., and Ball, J.S., 1963, Characteristics of crude oils from Utah, *in* Crawford, A.L., editor, Oil and gas possibilities of Utah, re-evaluated: Utah Geological and Mineralogical Survey Bulletin 54, p. 497–510.
- Wernicke, H., 1974, Gilsonit, ein bemerkenswertes Kohlenwasserstoff "mineral" (Gilsonite, a noteworthy hydrocarbon "mineral"): Geographische Rundschau, v. 26, no. 7, p. 290–292.
- Weston, W., 1904, Gilsonite and elaterite, Routt County: Mining Investor, v. 34, p. 72–74.
- Weston, W., 1907, The hydrocarbon field of western Colorado and eastern Utah on the projected line of the Denver, Northwestern and Pacific Railway: Report on the above to Mr. D.H. Moffat, president, 42 p.
- Westover, J.D., 1965, Chromatographic examination of gilsonite: Provo, Utah, Brigham Young University, PhD. dissertation, 140 p.
- Whiting, D.L., 1974, Energy resources of the Uinta Basin, Utah: Utah Geological Association Publication 4, 73 p.
- Woodward Clyde Consultants, 1981, Elk Ridge area geologic map: Oakland, California, unpublished consultant's report, scale 1:62,500.
- Wurtz, H., 1889, Uintahite, a variety of grahamite: Engineering and Mining Journal, v. 48, no. 6, p. 114.
- Young, R.S., 1954, Preliminary X-ray investigations of solid hydrocarbons: American Association of Petroleum Geologists Bulletin, v. 38, no. 9, p. 2017–2020.
- Yushkin, N.P., 1990, Fibrous forms of natural solid hydrocarbons: International Geology Review, v. 32, p. 1041–1050.
- Ziegler, G.S., 1941, Gilsonite and stearin pitch: American Ink Maker, v. 19, no. 4, p. 23–26.

APPENDIX

Annual production and value of gilsonite produced in the Uinta Basin

Much of the data in table A1 was compiled from the mineral yearbooks of the U.S. Bureau of Mines (USBM, 1953–1970) (later published by the U.S. Geological Survey). Mineral yearbook data became less valuable in the mid-1950s because the federal government started withholding data to protect company confidentiality (indicated by a "W" in the "QUANT." or "VALUE" data field). The Utah Division of Oil, Gas, and Mining (DOGGM) started to collect and report production data in the late 1980s. DOGM collects production data from all mines that it regulates; these data were incorporated into this compilation for the years 1989 through 2009. Unfortunately, companies do not report valuation to DOGM. The Utah Geological Survey also collects production data through a questionnaire sent to mine operators; these data were used in the compilation for the years 1993 through 2009. Miscellaneous additional sources of data are noted in the "DATA SOURCE" column and are listed in a reference list at the end of this appendix.

Most of the Uinta Basin gilsonite production has been by Utah, although Colorado produced a substantial quantity early in the history of the gilsonite industry. Two companies have dominated gilsonite production over the past 50 years: American Gilsonite Co. and G.S. Zeigler and Co. American Gilsonite has produced continuously since it was organized in 1946 and its predecessor companies produced for decades earlier. G.S. Zeigler (now Zeigler Chemicals and Minerals) purchased some of the smaller gilsonite producers in the early 1950s and has produced gilsonite continuously since. American Gilsonite typically produces more gilsonite per year than Zeigler. Lexco began producing gilsonite around 1989 and produced a small amount continuously until being acquired by American Gilsonite in 2009.

Table A1. Annual production and value of gilsonite produced in the Uinta Basin.

| YEAR | QUANT. (st) | VALUE (\$) | DATA SOURCE | NOTES |
|------|-------------|------------|----------------|--|
| 1885 | - | - | 2 | Small quantity produced and marketed. |
| 1886 | - | - | - | - |
| 1887 | - | - | - | - |
| 1888 | - | - | - | - |
| 1889 | 2100 | 252,000 | 9 | From January to September, 1500 st was shipped by Gilsonite Manufacturing Co. Gilson Asphaltum Co. produced another 600 st. The gilsonite sold for \$120/st. |
| 1890 | - | - | - | - |
| 1891 | - | - | - | - |
| 1892 | - | - | - | - |
| 1893 | - | - | - | - |
| 1894 | - | - | - | - |
| 1895 | - | - | - | - |
| 1896 | - | - | - | - |
| 1897 | - | - | - | - |
| 1898 | - | - | - | - |
| 1899 | >2000 | >100,000 | 8 | More than 2000 st of gilsonite shipped from the "Strip"; average price was \$50/st. Production and hauling costs were \$21/st. |
| 1900 | - | - | - | - |
| 1901 | - | - | 11 | Cost of mining and shipping material to St. Louis was \$25/st. Price in Chicago and St. Louis was \$40-50/st. |
| 1902 | 7000 | ~602,000 | 13 | 7000 st of gilsonite sold in Chicago at \$72/st in carloads or \$100/st in 2-ton lots. |
| 1903 | - | - | 13 | Sold for \$50/st f.o.b. railroad at Price, Ut.; company production costs were \$20/st. Wages paid were \$4 per 12-hour day. |
| 1904 | 2978 | - | 15 | - |
| 1905 | 10,916 | - | 6, 15 | Price of gilsonite in St. Louis was about \$50/st of which \$20/st was transportation costs. |
| 1906 | - | - | 8 | Freight rates on the Uintah Railroad were raised from \$8 to \$10/st. |
| 1907 | 20,285 | 531,965 | 8, 16 | Miners wages were \$3 per 8-hr day. |
| 1908 | 18,533 | 61,824 | 6, 16 | Freight charge from Dragon, Utah to New York was \$22.35/st. |
| 1909 | 28,669 | 218,186 | 16 | - |
| 1910 | 29,832 | 372,900 | 16 | - |
| 1911 | 30,236 | 486,114 | 16 | - |
| 1912 | 31,478 | 573,069 | 16 | - |
| 1913 | 35,055 | 576,949 | 6 | Includes gilsonite, wurtzilite, and grahamite. |
| 1914 | 19,148 | 405,966 | 6, 16 | Includes gilsonite, wurtzilite, and grahamite. |
| 1915 | 21,157 | 352,257 | 6, 16 | USBM reported that 1916 production was 27% more than 1915 and 1915 price was \$13.11/st. Carey and Roberts (1949) reported 20,559 st worth \$275,252 which included gilsonite and wurtzilite. |
| 1916 | 26,870 | 626,640 | 6, 16 | - |
| 1917 | 35,468 | 511,803 | 6, 16 | USBM reported that production was 32% greater than 1916 and was valued at \$14.43/st. Carey and Roberts (1949) reported 35,049 st worth \$532,989 which included gilsonite and wurtzilite. |
| 1918 | 30,848 | 606,639 | 1, 6, 16 | Aurand (1920) reported 31,072 st produced worth \$663,257 from five producers. |
| 1919 | | | | |
| 1920 | 56,204 | 548,776 | 6, 16 | - |
| 1921 | 10,066 | 178,224 | 6, 16 | - |
| 1922 | 29,693 | 622,107 | 6, 10, 16 | - |
| 1923 | 34,425 | 681,622 | 6, 10, 16 | Average price was \$28.00/st. |
| 1924 | 35,907 | 603,620 | 6, 10, 16 | - |
| 1925 | 39,520 | 767,900 | 6, 10, 16 | - |
| 1926 | 42,190 | 863,840 | 6, 10, 16 | USBM noted that values may be inflated by some companies reporting a delivered value rather than at mine/railhead. |
| 1927 | 42,580 | 876,820 | 6, 10, 16 | USBM noted that values may be inflated by some companies reporting a delivered value rather than at mine/railhead. |
| 1928 | 47,023 | 1,037,679 | 6, 10, 16 | USBM noted that values may be inflated by some companies reporting a delivered value rather than at mine/railhead. Dennis (1930) wrote that the average price f.o.b. mine was \$22.07/st. Selects quality ore at |

| | | | | |
|------|---------|------------|---------------|--|
| | | | | Mack, Colorado was worth \$30–\$33/st. |
| 1929 | 54,987 | 1,235,920 | 6, 15 | Romney (1963) reported that this production is only for Uintah Co. Value was not totally based on f.o.b. price; f.o.b price was \$14.50–\$28.00/st. Carey and Roberts (1949) reported that the largest producers in 1929 were Gilson Asphaltum Co., Utah Gilsonite Co., Raven Mining Co. of Utah, Diamond Gilsonite Co., and the American Asphalt Association. |
| 1930 | 37,684 | 863,197 | 6, 16 | Dennis (1930) wrote that in January 1930, the Rainbow mine shipped 125 st daily which was about 80% of the total production from this area. About 20% of the gilsonite was shipped to foreign countries. Roughly 50% of the gilsonite was low-grade material used for roofing paper and other uses. Average price was \$22.91/st. |
| 1931 | 32,763 | 674,102 | 6, 16 | - |
| 1932 | 25,955 | 525,266 | 6, 17 | Drop in production due to reduced activity in manufacturing of paint, rubber, and insulated wire. |
| 1933 | 28,029 | 577,716 | 6, 17 | - |
| 1934 | 30,355 | 599,739 | 6, 17 | Increased production over 1933 due to greater paint demand and activity in the electrical industry. |
| 1935 | 33,277 | 696,601 | 6, 17 | - |
| 1936 | 33,694 | 833,966 | 17 | Higher prices due to increased demands for paint, varnish, and electrical fitting manufacture. |
| 1937 | 38,038 | 973,007 | 6, 17 | - |
| 1938 | 28,574 | 649,724 | 6, 17 | Ten percent increase in railroad freight rates and decreased demand for varnish and ink reduced domestic demand. Higher steamship rates and exchange difficulties reduced exports. |
| 1939 | 37,289 | 1,053,142 | 6, 14, 17 | Carey and Roberts (1949) reported, "Demand for better grades \$28.24/st during first part of year." |
| 1940 | 31,930 | 770,711 | 6, 14, 17 | Carey and Roberts (1949) reported, "Exports dropped off—Germany blockaded—f.o.b. \$24.14/ton." |
| 1941 | 36,407 | 851,623 | 6, 14, 17 | Price at mine was \$23.39/st. |
| 1942 | 40,041 | 909, 311 | 6, 14, 17, 20 | - |
| 1943 | 50,466 | 1,188,485 | 6, 14, 17 | \$23.56/st. |
| 1944 | 49,051 | 915,480 | 6, 14, 17 | \$18.73/st f.o.b. |
| 1945 | 61,273 | 1,250,546 | 6, 14, 17 | \$20.41/st f.o.b. |
| 1946 | 68,407 | 1,440,229 | 6, 14, 17 | All production was from Utah. Barber Asphalt Corp. sold half of its gilsonite business to Standard Oil of California (Chevron) and the two formed a joint partnership named the American Gilsonite Co. |
| 1947 | 67,165 | 1,746,228 | 6, 14, 17 | - |
| 1948 | 52,122 | 1,390,713 | 6, 14, 17 | - |
| 1949 | 51,462 | 1,303,584 | 6, 17 | From 1908 to 1949 prices fluctuated between \$20 and \$40/st f.o.b. at Craig, Colorado. |
| 1950 | 66,186 | 1,774,330 | 12, 17 | Garvin (1966) reported combined asphalt and gilsonite production of 66,186 st worth \$1,779,815. |
| 1951 | 65,521 | 1,895,374 | 15, 17, 19 | Production for Duchesne and Uintah Counties combined. |
| 1952 | 60,740 | 1,779,815 | 17, 19 | Production for Duchesne and Uintah Counties combined. |
| 1953 | 60,505 | 2,184,328 | 17, 19 | Production for Duchesne and Uintah Counties combined. |
| 1954 | 60,500 | 2,184,000 | 17, 19 | - |
| 1955 | W | W | 19 | American Gilsonite Co. was the largest producer, with mines and a plant at Bonanza. Several of the smaller operations were purchased a few years earlier by G.S. Ziegler & Co. Ziegler had a processing plant in Provo. Duchesne County's Castle Peak gilsonite district was active in 1955. |
| 1956 | W | W | 20 | - |
| 1957 | 206,041 | 4,259,120 | 13, 17, 20 | Price of gilsonite at the railhead is about \$35/st. Miners earned about \$20 for an 8-hour day. |
| 1958 | - | - | - | - |
| 1959 | - | - | 20 | Ziegler began construction of its processing plant in Uintah County (Little Bonanza?). |
| 1960 | - | - | - | - |
| 1961 | 470,000 | 14,500,000 | 7, 15, 17 | Gilsonite production was lumped by USBM into the native asphalt category but almost all was due to gilsonite. Two companies operated, American and Ziegler. Cashion (1964) reported that gilsonite production in 1961 was 422,294 st valued at \$9,916,000. |
| 1962 | - | - | - | - |
| 1963 | - | - | - | - |
| 1964 | W | W | 17, 21 | Value of production increased 22% from 1963. Active operators included American Gilsonite Company, Ziegler Chemical and Minerals Corp., Standard Gilsonite Company, Arthur Boren, and Alva L. Hatch. |
| 1965 | W | W | 17, 21 | Gilsonite production decreased 13% from 1964. American, Ziegler, and Standard operated mines in Uintah County. Ziegler also operated the Castle Peak mine in Duchesne County. Standard Gilsonite Co. became a division of Mesa Petroleum Co. on September 10, 1965. |
| 1966 | W | W | 17 | Gilsonite production decreased 24% from 1965. American, Ziegler, and Standard operated mines in Uintah County. Ziegler also operated the Castle Peak mine in Duchesne County. |
| 1967 | W | W | 17 | Gilsonite production from five mines in Uintah and Duchesne Counties was down 10%. The Castle Peak |

| | | | | |
|------|--------|---|----|--|
| | | | | mine in Duchesne County was closed after 30 years of production. Standard Gilsonite mined gilsonite in Duchesne County, processed it at a plant south of Myton, and made intermittent shipments. Other operators making occasional shipments include Arthur Boren and Alva L. Hatch. |
| 1968 | W | W | 17 | Yearbook contains no notes. |
| 1969 | W | W | 17 | Total output from the three gilsonite-producing companies was slightly higher than that of the previous year. |
| 1970 | W | W | 17 | Output from three gilsonite-producing companies rose 14% in tonnage and 12% in value in 1970. |
| 1971 | W | W | 17 | Total output from three gilsonite-producing companies fell 23% in tonnage and 19% in value |
| 1972 | W | W | 17 | Output from two gilsonite-producing companies dropped 15% but value rose 12% as a result of a higher unit value. |
| 1973 | W | W | 17 | Output from two gilsonite-producing companies fell 69% in quantity and 53% in value from that in 1972. The Fruita, CO, plant of American Gilsonite was sold in December 1973. |
| 1974 | W | W | 17 | Output from two gilsonite-producing companies rose 23% in quantity and as a result of increased unit price, value was more than triple that of 1973. |
| 1975 | W | W | 17 | Two companies produced gilsonite from properties in Uintah Co. Total production was 19% less than that during 1974 but as a result of higher average price for the product, total value was 2% greater. |
| 1976 | W | W | 17 | Output from the state's two gilsonite-producing properties increased 9% in quantity but decreased 12% in value as a result of lower average prices for this commodity. |
| 1977 | W | W | - | - |
| 1978 | W | W | 17 | Gilsonite was produced by American Gilsonite Co. at Bonanza in Uintah County, Utah, and by Ziegler Chemical and Minerals Corp. in Weber County. By mid-1979 American Gilsonite completed construction of its \$5.3 million consolidated processing plant at Bonanza. |
| 1979 | W | W | 17 | 1978 and 1979 yearbook reports were combined. |
| 1980 | W | W | - | - |
| 1981 | W | W | - | Chevron bought out Barber Oil's half ownership of American Gilsonite Co. |
| 1982 | W | W | 17 | American Gilsonite is listed as the principal producer. |
| 1983 | W | W | - | - |
| 1984 | W | W | - | - |
| 1985 | W | W | - | - |
| 1986 | W | W | - | - |
| 1987 | W | W | - | - |
| 1988 | W | W | 17 | American Gilsonite Co. and Ziegler Chemical and Minerals Corp. were listed as the two main producers. Lexco enters the gilsonite business. |
| 1989 | 52,873 | W | 17 | American Gilsonite Co. and Ziegler Chemical and Minerals Corp. were listed as the two producers. USBM withheld data but production is contained in DOGM files |
| 1990 | W | W | - | - |
| 1991 | W | W | - | Chevron sold American Gilsonite Co. to Stratford Enterprises Co. of Tulsa, OK. |
| 1992 | W | W | 17 | American Gilsonite Co. and Ziegler Chemical and Minerals Corp. were listed as the two principal producers. |
| 1993 | 47,124 | W | 3 | - |
| 1994 | 56,257 | W | 4 | - |
| 1995 | 57,819 | W | 5 | - |
| 1996 | 62,925 | W | 18 | Producers in decreasing order: American Gilsonite Co. (Bonanza mine), Ziegler Chemical and Minerals Corp. (Ziegler mine), and Lexco (Lexco mine). |
| 1997 | 66,032 | W | 18 | Producers in decreasing order: American Gilsonite Co. (Bonanza mine), Ziegler Chemical and Minerals Corp. (Ziegler mine), and Lexco (Lexco mine). |
| 1998 | 69,172 | W | 18 | Producers in decreasing order: American Gilsonite Co. (Bonanza mine), Ziegler Chemical and Minerals Corp. (Ziegler and Tom Taylor mines), and Lexco (Cottonwood mine). |
| 1999 | 59,000 | W | 18 | Producers in decreasing order: American Gilsonite Co. (Bonanza mine), Ziegler Chemical and Minerals Corp. (Ziegler and Tom Taylor mines), and Lexco (Cottonwood mine). |
| 2000 | 72,300 | W | 18 | Producers in decreasing order: American Gilsonite Co. (Bonanza mine), Ziegler Chemical and Minerals Corp. (Ziegler and Tom Taylor mines), and Lexco (Cottonwood mine). |
| 2001 | 69,450 | W | 18 | Producers in decreasing order: American Gilsonite Co. (Bonanza mine), Ziegler Chemical and Minerals Corp. (Cowboy, Neal State, and Hardaway mines), and Lexco (Cottonwood mine). |
| 2002 | 65,648 | W | 18 | Producers in decreasing order: American Gilsonite Co., Ziegler Chemical and Minerals Corp., and Lexco. |
| 2003 | 56,943 | W | 18 | Producers in decreasing order: American Gilsonite Co., Ziegler Chemical and Minerals Corp., and Lexco. |

| | | | | |
|------|--------|---|----|--|
| 2004 | 70,000 | W | 18 | Producers in decreasing order: American Gilsonite Co., Ziegler Chemical and Minerals Corp., and Lexco. Lexco production not included in quantity. |
| 2005 | 79,300 | W | 18 | Producers in decreasing order: American Gilsonite Co., Ziegler Chemical and Minerals Corp., and Lexco. |
| 2006 | 81,200 | W | 18 | Producers in decreasing order: American Gilsonite Co., Ziegler Chemical and Minerals Corp., and Lexco. |
| 2007 | 75,853 | W | 18 | Producers in decreasing order: American Gilsonite Co., Lexco, and Ziegler Chemical and Minerals Corp. |
| 2008 | 76,600 | W | 18 | Producers in decreasing order: American Gilsonite Co., Lexco, and Ziegler Chemical and Minerals Corp. Lexco portion of production estimated. Palladium Equity Partners bought American Gilsonite Co. in March. |
| 2009 | 63,500 | W | 18 | Lexco portion of production estimated. Lexco is acquired by American Gilsonite. |


Abbreviations: f.o.b., free on board; st, short tons; W, withheld.

Notes: values not inflation adjusted; free on board (f.o.b.) is an International Chamber of Commerce term to indicate the price of a commodity when all transportation, insurance, and other costs have been settled and it has been loaded onto the deck of a ship (technically the term should have been f.o.r.—free on rail—for gilsonite transportation); the title and risk pass to the buyer at this point.

SOURCES OF DATA

- 1 Aurand, H.A., 1920, Mineral deposits of the western slope: Colorado Geological Survey Bulletin 22, p. 15–19.
- 2 Baker, J.H., 1950, Economics of gilsonite in the Uinta Basin, *in* Eardley, A.J., editor, Petroleum geology of the Uinta Basin: Intermountain Association of Petroleum Geologists, Fifth Annual Field Conference, p. 119–120.
- 3 Bon, R.L., 1993, State activities 1993-Utah: Mining Engineering, vol. 46, no. 5, p. 418–420.
- 4 Bon, R.L., Gloyd, R.W., and Tabet, D.E., 1994, State activities 1994—Utah: Mining Engineering, vol. 47, no. 5, p. 447–452.
- 5 Bon, R.L., Gloyd, R.W., and Tabet, D.E., 1995, Utah mineral activity summary for 1995: Utah Geological Survey Circular 91, 15 p.
- 6 Carey, G.A., and Roberts, I.C., 1949, Dissertation on the history, occurrence, mining, and economics of gilsonite: Salt Lake City, University of Utah, B.S. thesis, 89 p.
- 7 Cashion, W.B., 1964, Other bituminous substances, *in* Hilpert, L.S., editor, Mineral and water resources of Utah: Utah Geological and Mineralogical Survey Bulletin 73, p. 63–70.
- 8 Covington, R.E., 1964, A brief history of early mineral exploration in the Uinta Basin, *in* Sabatka, E.F., editor, Geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse: Intermountain Association of Petroleum Geologists Guidebook, 13th Annual Field Conference, p. 1–16.
- 9 Crawford, A.L., 1957, Gilsonite - its discovery and the early history of the industry, *in* Seal, O.G., editor, Geology of the Uinta Basin: Intermountain Association of Petroleum Geologists, Eight Annual Field Conference, p. 149–151.
- 10 Dennis, E., 1930, A preliminary survey of Utah non-metallic minerals with special reference to their occurrence and markets for them: Provo, Utah, Brigham Young University, M.A. thesis, p. 108–114.
- 11 Eldridge, G.H., 1901, The asphalt and bituminous rock deposits of the United States, *in* Walcott, C.D., director, U.S. Geological Survey Twenty-Second Annual Report of the United States Geological Survey to the Secretary of the Interior, pt. 1, p. 209–364.
- 12 Garvin, R.F., compiler, 1966, A directory of the mining industry of Utah, 1965: Utah Geological and Mineralogical Survey Bulletin 79, 94 p.
- 13 Kretchman, H.F., 1957, The story of gilsonite: Salt Lake City, Utah, American Gilsonite Company, p. 37–38.
- 14 Ladoo, R.B., and Myers, W.M., 1951, Nonmetallic minerals, 2nd edition: New York, McGraw-Hill Book Company, Inc., p. 57–67.


- 15 Romney, M.P., 1963, Utah's Cinderella minerals—the nonmetallics: *Utah Historical Quarterly*, vol. 31, no. 3, p. 229–230.
- 16 U.S. Bureau of Mines, 1907 to 1931, *Minerals yearbooks*: Washington, D.C., U.S. Government Printing Office, variously paginated.
- 17 U.S. Bureau of Mines, 1932 to 1992, *Minerals yearbooks*: Washington, D.C., U.S. Government Printing Office, variously paginated, available online at <http://minerals.usgs.gov/minerals/pubs/usbmmyb.html>.
- 18 Utah Geological Survey, 1996 to 2009, *Summary of mineral activity in Utah*: Utah Geological Survey Circulars, variously paginated, available online at <http://geology.utah.gov/utahgeo/rockmineral/index.htm#minactivity>.
- 19 Utah Mining Association, 1955, *Utah's mining industry—an historical, operational, and economic review of Utah's mining industry*, 1st edition: Salt Lake City, Utah, p. 24–79.
- 20 Utah Mining Association, 1959, *Utah's mining industry—an historical, operational, and economic review of Utah's mining industry*, 2nd edition: Salt Lake City, Utah, 132 p.
- 21 Utah Mining Association, 1967, *Utah's mining industry—an historical, operational, and economic review of Utah's mining industry*, 3rd edition: Salt Lake City, Utah, 135 p.



Although this product represents the work of professional scientists, the Utah Department of Natural Resources, Utah Geological Survey, makes no warranty, expressed or implied, regarding its suitability for a particular use, and does not guarantee accuracy or completeness of the data. The Utah Department of Natural Resources, Utah Geological Survey, shall not be liable under any circumstances for any direct, indirect, special, incidental, or consequential damages with respect to claims by users of this product. For use at 1:100,000 scale only.

GEOLOGIC UNITS EXPLANATION


| | | |
|--|--|--|
| <ul style="list-style-type: none"> — Gilsonite vein, well located - - - Gilsonite vein, inferred - · - · - Gilsonite vein, uncertain — Mahogany oil-shale marker bed, well located — Contact, well located — Normal fault, well located - - - Normal fault, approximately located - - - Normal fault, concealed — Bar and ball on downthrown side where offset is known | <ul style="list-style-type: none"> Qal Stream Alluvium (Holocene) Qat Stream Terrace Deposits (Holocene and Pleistocene[?]) Qa Basin Alluvium (Holocene and Pleistocene[?]) Qaf Alluvial-Fan Deposits (Holocene and Pleistocene[?]) Qc Colluvium (Holocene) Qe Eolian Deposits (Holocene) Qms Landslides, Slumps, and Flows (Holocene and Pleistocene) Qac Mixed Alluvium and Colluvium (Holocene and Pleistocene[?]) Qae Mixed Alluvium and Eolian Deposits (Holocene) | <ul style="list-style-type: none"> Tdb Brennan Basin Member of Duchesne River Formation (Eocene) Tuc Member C of Uinta Formation (Eocene) Tub Member B of Uinta Formation (Eocene) Tua Member A of Uinta Formation (Eocene) Tgsl Sandstone and Limestone Facies of Green River Formation (Eocene) Tgp Paratcha Creek Member of Green River Formation (Eocene) Tgd Douglas Creek Member of Green River Formation (Eocene) Tg-Tw Green River–Wasatch Formations Transition Zone (Eocene) Tw Wasatch Formation (Eocene and Paleocene[?]) |
|--|--|--|



GILSONITE VEINS OF THE UINTA BASIN
 by
 Taylor Boden and Bryce T. Tripp
 2012



INDEX TO U.S. GEOLOGICAL SURVEY 30' x 60' QUADRANGLES AND SOURCES OF GEOLOGIC MAPPING



Topographic base from U.S. Geological Survey Duchesne (1982), Price (1980), Seep Ridge (1981), and Vernal (1982) 30' x 60' quadrangles. Elevation from 10m National Elevation Data. Geology compiled from Bryant (1992), Weiss and others (2003), and Sprinkel (2007, 2009). Projection: UTM, Zone 12. Units: Meters. Datum: NAD 1983.

Utah Geological Survey
 1594 West North Temple, Suite 3110
 P.O. Box 146100, Salt Lake City, UT 84114-6100
 (801) 537-3300
geology.utah.gov