


GEOLOGIC MAP OF THE MANTI 30' x 60' QUADRANGLE, CARBON, EMERY, JUAB, SANPETE, AND SEVIER COUNTIES, UTAH

by
Irving J. Witkind, Malcolm P. Weiss, and Terrence L. Brown

MANTI, UTAH

30X60 MINUTE SERIES (TOPOGRAPHIC)



Base from U.S. Geological Survey, 1981
Projection: UTM Zone 12
Units: Meters
Datum: NAD 83
Spheroid: Clarke 1866

Although this product represents the work of professional geologists, the Utah Department of Natural Resources, Utah Geological Survey, makes no warranty, expressed or implied, regarding the accuracy of this data. These data are not to be used for any purpose other than that for which they were collected, and the user assumes all responsibility for any use of these data, or consequential damages with respect to claims by users of this product.

This map is a cooperative project of the USGS and the Utah Geological Survey (UGS) funded through the STATEMAP component of the National Cooperative Geologic Mapping Program, Agreement No. G24AG0065.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.




SCALE 1:100 000
CONTOUR INTERVAL 50 METERS
SUPPLEMENTARY CONTOUR INTERVAL 25 METERS



Caution to data user: The original published topographic map from which these GIS data files were prepared contains problems that affect the accuracy of this data. These include: 1) the topographic base map was not properly registered, as a result the geology does not properly fit the base topography; 2) color separations for the original topographic base map were not properly registered with each other and 3) some quadrangles for 7.5' quadrangle tickmarks are not in the correct locations. While preparing this GIS database, the USGS made considerable effort to better fit the digital geology with the other data by using digital 7.5' topographic maps as guides, but the results were not entirely satisfactory and the eastern part of the digital 1:100,000-scale geologic and topographic maps do not match. Discrepancies were noted across the map, but may locally exceed 100 meters on the ground. Therefore, we recommend that the digital geology be used only at a scale of 1:100,000, and with the digital topographic map included on this CD. These inaccuracies will be accumulated if this data is "stretched" over other 1:100,000-scale topographic base maps or "blown up" to scales larger than 1:100,000.


Project Manager: Jon R. King
GIS Data Preparation: Baska Matjask

CORRELATION OF MAP UNITS¹



The Utah Geological Survey modified this explanation sheet from the original publication to match the GIS database project files (Utah Geological Survey Map 212DM). Changes include modifying map unit colors to match colors used in the project file, and slight wording changes to clarify meaning of a few unit descriptions. Care was taken to preserve the authors' original intent.

¹(Unlabeled units, shown uncolored on correlation, are not shown on map)



REFERENCES CITED

(References shown by colors or patterns are keyed to the sources of geologic data)

American Society for Testing and Materials, 1978. Standard specifications for classification of coals by rank (ASTM designation D 388-77). Annual Book of ASTM standards, Pt. 26, p. 214-218.

Anonymous, 1961. Gordon Creek field, Carbon County, Utah: A symposium of the oil and gas fields of Utah.

Babisak, Julius, 1949. The geology of the southeastern portion of the Gunnison Plateau, Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:24,000.

Baer, J. L., 1976. Structural evolution of central Utah—Late Permian to Recent, in Geology of Cordilleran hinge-line, 1976. Rocky Mountain Association of Geologists Guidebook, 27th Annual Field Conference, p. 37-45.

Baker, A. A., 1923. Geology and oil possibilities of the Moab district, Grand and San Juan Counties, Utah. U.S. Geological Survey Bulletin 841, 95 p.

Baughman, R. L., 1959. The geology of the Musina graben area, Sevier and Sanpete Counties, Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:24,000.

Bonar, C. M., 1948. Geology of the Ephraim area, Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:31,680.

Brown, T. L., and Sanchez, D. L., 1987. Stratigraphic framework and coal resources of the Upper Cretaceous Blackhawk Formation in the East Mountain and Gentry Mountain areas, Mantt 30' x 60' quadrangle, Wasatch Plateau coal field, Emery and Carbon Counties, Utah. U.S. Geological Survey Coal Investigations Map C-94-2.

Detterman, J. S., 1956. Photogeologic map of Castle Dale-16 [Moine] quadrangle, Emery County, Utah. U.S. Geological Survey Miscellaneous Investigations Map I-124, scale 1:24,000.

Doelling, H. H., 1972a. Coal in the Sevier-Sanpete region: Utah Geological Association Publication 2, Plateau-Basin and Range transition zone, central Utah, p. 81-89.

—1972b. Central Utah coal fields: Utah Geological and Mineralogical Survey, Monograph Series No. 3, 571 p.

Duncan, D. C., 1944. The Mount Pleasant coal field, Sanpete County, Utah. U.S. Geological Survey map, scale 1:36,000.

Ellis, E. G., 1979. Geologic map of the Ferron Canyon quadrangle, Emery and Sanpete Counties, Utah. Unpublished map, scale 1:24,000. (U.S. Geological Survey, Box 25046, Phoenix, Arizona 85025)

—1980. Geologic map of the Cap quadrangle, Emery County, Utah. U.S. Geological Survey, unpublished map, scale 1:24,000.

Faulk, N. R., 1948. The Green River Formation in the Mantt-Spring City area of central Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:31,680.

Fleming, R. W., Schuster, R. L., Johnson, R. B., and Robinson, S. L., 1977. Recent movement of the Mantt, Utah, Landslide: Idaho State University, Pocatello, Idaho, Proceedings 15th Annual Engineering Geology and Soils Engineering Symposium, April 6-8, 1977, p. 161-178.

Gilliland, W. N., 1948. Geology of the Gunnison quadrangle, Utah. Columbus, Ohio State University, Ph. D. thesis, 178 p.

—1951. Geology of the Gunnison quadrangle, Utah: University of Nebraska Studies, no. 8, 101 p.

—1952. Another Tertiary crustal disturbance in central Utah: American Association of Petroleum Geologists Bulletin, v. 36, no. 7, p. 1461-1464.

Hardy, C. T., 1949. Stratigraphy and structure of the Arapian Shale and the Twist Gulch Formation in Sevier Valley, Utah. Columbus, Ohio State University, Ph. D. thesis, 85 p.

—1952. Eastern Sevier Valley, Sevier and Sanpete Counties, Utah: Utah Geological and Mineralogical Survey Bulletin 43, 41 p., scale 1:187,500; pl. 5, scale 1:62,500.

Hardy, C. T., and Zeller, H. D., 1953. Geology of the west-central part of the Gunnison Plateau, Utah. Geological Society of America Bulletin, v. 64, no. 11, p. 1, scale 1:187,500.

Heyl, A. V., 1963. Oxidized zinc deposits of the United States: U.S. Geological Survey Bulletin 1135-B, 104 p.

—1978. Redmond silver mine, Redmond District, Sevier County, Utah, in Guidebook to mineral deposits of southwestern Utah: Utah Geological Association Publication 7, p. 73-75.

Hunt, R. E., 1950. Geology of the northern part of the Gunnison Plateau, Utah. Columbus, Ohio State University, Ph. D. thesis, pl. 1, scale 1:31,680.

John, C., 1972. Petrology and petrography of the intrusive igneous rocks of the Levan area, Juab County, Utah, in Plateau-Basin and Range transition zone, central Utah, 1972: Utah Geological Association Publication 2, p. 97-107.

Johnson, M. S., 1949. Geology of the Twelve Mile Canyon area, central Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:31,680.

Kirkland, D. W., and Evans, Robert, 1981. Source-rock potential of evaporitic environment: American Association of Petroleum Geologists Bulletin, v. 65, no. 2, p. 181-191.

Kucera, R. E., 1954. Geology of the Joes Valley and North Dragon area, central Utah. Columbus, Ohio State University, M.S. thesis, pl. 30, scale 1:31,680.

Millen, T. M., 1982. Stratigraphy and petrology of the Green River Formation (Eocene), Gunnison Plateau, Utah. DeKalb, Illinois, Northern Illinois University, M.S. thesis, 220 p.

Moulton, F. C., 1975. Lower Mesozoic and Upper Paleozoic petroleum potential of the hinge-line area, central Utah, in Deep drilling frontiers in the central Rocky Mountains, 1975: Rocky Mountain Association of Geologists Guidebook, p. 87-97.

Muessig, S. J., 1951. Geology of a part of Long Ridge, Utah. Columbus, Ohio State University, Ph. D. thesis, pl. 1, scale 1:31,680.

Niehaus, J. W., 1956. Geology of the northeastern Valley Mountains and adjacent Sevier Valley area, Utah. Columbus, Ohio State University, Incomplete M.S. thesis, pl. 1, scale 1:31,680.

Pashley, E. F., Jr., 1956. The geology of the western slope of the Wasatch Plateau between the Arapian Shale and the Gunnison Plateau, Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:62,500.

Peterson, V. E., 1961. Farham dome field, Emery County, Utah: A symposium of the oil and gas fields of Utah.

Picard, M. D., 1980. Stratigraphy, and origin of evaporites, Jurassic Arapian Shale, central Utah, in Henry Mountains Symposium: Utah Geological Association Publication 8, p. 129-150.

Pratt, A. R., and Callaghan, E., 1970. Land and mineral resources of Sanpete County, Utah. Utah Geological and Mineralogical Survey Bulletin 55, 69 p.

Pratt, A. R., Heylman, E. B., and Cohonour, R. E., 1966. Salt deposits of Sevier Valley, Utah, in Second Symposium on salt: Northern Ohio Geological Society, Pt. 1, 48 p.

Price, 1983. Map showing selected surface-water data for the Mantt 30' x 60' quadrangle, Utah. U.S. Geological Survey Minerals Investigations Map I-1482.

Sanchez, J. D., Blanchard, L. F., and August, L. L., 1983. Stratigraphic framework and coal resources of the Upper Cretaceous Blackhawk Formation in the Muddy Creek and Nelson areas, Mantt 30' x 60' quadrangle, Wasatch Plateau coal field, Emery and Sanpete Counties, Utah. U.S. Geological Survey Coal Investigations Map C-93-B.

Sanchez, J. D., and Brown, T. L., 1983. Stratigraphic framework and coal resources of the Upper Cretaceous Blackhawk Formation in the Muddy Creek and Nelson areas, Mantt 30' x 60' quadrangle, Wasatch Plateau coal field, Emery and Sanpete Counties, Utah. U.S. Geological Survey Coal Investigations Map C-94-A.

—1987. Stratigraphic framework and coal resources of the Upper Cretaceous Blackhawk Formation in the central Utah area, Mantt 30' x 60' quadrangle, Wasatch Plateau coal field, Emery and Sanpete Counties, Utah. U.S. Geological Survey Coal Investigations Map C-94-B.

—1988. Stratigraphic framework and coal resources of the Upper Cretaceous Blackhawk Formation in the Trail Mountain and East Mountain areas, Mantt 30' x 60' quadrangle, Wasatch Plateau coal field, Emery County, Utah. U.S. Geological Survey Coal Investigations Map C-94-C.

Sanchez, J. D., and Hayes, P. T., 1979. Geology and coal resources of the Flagstaff Peak quadrangle, Emery, Sanpete, and Sevier Counties, Utah: U.S. Geological Survey Coal Investigations Map C-83, scale 1:24,000.

Speiker, E. M., 1931. The Wasatch Plateau coal field, Utah: U.S. Geological Survey Bulletin 1178, scale 1:62,500.

—1946. Late Mesozoic and early Cenozoic history of central Utah. U.S. Geological Survey Professional Paper 205-D, p. 117-161.

—1948. Geology of the transition between the Colorado Plateau and the Great Basin in central Utah. Utah Geological Society Guidebook 4, pt. 1, scale 1:125,000.

Sprinkel, D. A., 1982. Twin Creek Limestone-Arapian Shale relations in central Utah, in Overthrust salt belt of Utah: Utah Geological Association Publication 10, p. 169-179.

Stokes, W. L., 1952. Salt-generated structures of the Colorado Plateau and possible analogies [abs.]. American Association of Petroleum Geologists Bulletin, v. 36, no. 1, p. 961.

—1956. Tectonics of Wasatch Plateau and near-by areas [abs.]. American Association of Petroleum Geologists Bulletin, v. 40, no. 4, p. 790.

Stokes, W. L., and Holmes, C. N., 1954. Jurassic rocks of south-central Utah in Geology of the High Plateaus and adjacent Canyonlands, central and south-central Utah. Intermountain Association of Petroleum Geologists Guidebook, 5th Annual Field Conference, p. 34-41.

Taylor, D. A., 1948. The geology of the Gunnison Plateau front in the vicinity of Wales, Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:24,000.

Taylor, J. M., 1980. Geology of the Sterling quadrangle, Sanpete County, Utah: Brigham Young University Geology Studies, v. 27, pt. 1, fig. 4, scale 1:24,000.

Vogel, J. W., 1957. Geology of the southernmost Juab Valley and adjacent highlands, Juab County, Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:31,680.

Volker, D. G., 1980. Stratigraphy and petrology of the Colton Formation (Eocene), Gunnison Plateau, central Utah. DeKalb, Illinois, Northern Illinois University, M.S. thesis, 132 p.

Walton, P. T., 1963. Gas fields of the Wasatch Plateau area, central Utah and Mineralogical Survey Bulletin 54, p. 341-352.

Washington, C. R., 1945. Geology of the Mantt Canyon area, central Utah. Columbus, Ohio State University, M.S. thesis, pl. 1, scale 1:31,680.

Weiss, M. P., 1969. Oncolites, paleoecology, and Laramide tectonics, central Utah: American Association of Petroleum Geologists Bulletin, v. 53, no. 5, p. 1105-1120.

—1982. Structural variation on east front of the Gunnison Plateau, central Utah in Overthrust belt of Utah: Utah Geological Association Publication 10, p. 49-63.

Wilson, M. D., 1949. The geology of the upper Six Mile Canyon area, central Utah. Columbus, Ohio State University, M.S. thesis, pl. 10, scale 1:31,680.

Witkind, I. J., 1979. Reconnaissance geologic map of the Wellington quadrangle, Carbon County, Utah: U.S. Geological Survey Miscellaneous Investigations Series Map I-178, scale 1:24,000.

—1981. Reconnaissance geologic map of the Redmond quadrangle, Sanpete and Sevier Counties, Utah. U.S. Geological Survey Miscellaneous Investigations Series Map I-1304-A, scale 1:24,000.

—1982. Salt diapirs in central Utah in Overthrust belt of Utah: Utah Geological Association Publication 10, p. 13-30.

—1983. Overthrusts and salt diapirs, central Utah, in Tectonic and stratigraphic studies in the eastern Great Basin region, Geological Society of America Memoir 157, p. 45-59.

Young, R. A., and Carpenter, C. H., 1965. Ground-water conditions and storage in the central Sevier Valley, Utah. U.S. Geological Survey Water-Supply Paper 1787.

DESCRIPTION OF MAP UNITS

- QUATERNARY DEPOSITS**
- Oal** ALLUVIUM (HOLOCENE)—Dark brown to gray, thin to thick bedded, locally massive, crossbedded in places. Unconsolidated. Consists of clay, silt, sand, granules, pebbles, and coarse cobbles. Fluvial sediments. Forms broad, even surfaces of low relief. Locally includes higher patches of "older alluvium". Thickness ranges widely; commonly less than 15 m (50 ft) thick.
- Oao** OLDER ALLUVIUM (HOLOCENE AND PLEISTOCENE)—Much like alluvium in color, bedding, and composition. Fluvial sediments. Forms broad, rounded to irregular masses generally 15-45 m (50-150 ft) above adjacent valley floor. Thickness varies greatly, ranging from about 3-60 m (10-200 ft).
- Ocl** COLLUVIUM (HOLOCENE)—Brown to dark-brown heterogeneous mixture of fragments of many sizes and shapes which locally mantles lower valley walls and accumulates at the base of some steep cliffs. Unconsolidated to semi-consolidated debris. Thickness ranges from a few centimeters (one inch) to as much as 17 m (50 ft).
- Osw** SLOPE WASH (HOLOCENE)—Light to dark gray; thin to thick bedded, laminated locally, faintly crossbedded. Unconsolidated to weakly cemented locally. Consists of clay, silt, sand, granules, and some pebbles. Fluvial sediments. Forms broad, gently sloping sheets. Thickness ranges from a thin film to as much as 8 m (25 ft).
- Oq** DUNE SAND (HOLOCENE AND PLEISTOCENE)—Light-brown to brown unconsolidated, well-sorted frosted grains of fine quartzose sand. Eolian sediments. Locally forms small active dune fields. Thickness ranges from a thin veneer to about 8 m (25 ft).
- Oqf** ALLUVIAL FAN DEPOSITS (HOLOCENE)—Light brown to brown, locally gray; unconsolidated to semi-consolidated; moderately well sorted silt, sand, granules, pebbles, or cobbles at the mouths of streams. Fluvial deposits. Commonly lobate. Thickness uncertain, probably as much as 15 m (50 ft) locally.
- Oq1** LANDSLIDE DEPOSITS (HOLOCENE AND PLEISTOCENE)—Brown to dark-brown and gray heterogeneous mixture of fragments of diverse sizes and shapes. Debris irregular to lobate masses of bedrock that have downslope to form chaotic accumulation of rubble. Mantt landslide, in Mantt Canyon, is one of the larger landslides that has moved recently and threatened man-made features (Fleming and others, 1977). Hummocky topography, locally some conchoidal ridges. Thickness varies widely; may be as much as 45 m (150 ft) thick locally.
- Oe** EARTHFLOW DEPOSITS (HOLOCENE AND PLEISTOCENE)—Brown to dark brown, sand, granules, pebbles, cobbles, and boulders in an unsorted matrix of clay and silt. Unconsolidated to semi-consolidated debris. Masses of debris that have flowed downslope to form elongate, hummocky, lobate land forms. Thickness varies widely; probably as much as 45 m (150 ft) thick locally.
- Ors** ROCK SLIDE DEPOSITS (HOLOCENE AND PLEISTOCENE)—Light-gray to brown unconsolidated and unsorted chaotic accumulation of angular boulders on steep slopes. Debris, hummocky, locally lobate. Ranges in thickness from a few meters (about 10 ft) to as much as 45 m (150 ft) locally.
- Omw** MASS-WASTING DEPOSITS (HOLOCENE AND PLEISTOCENE)—Brown to dark-brown heterogeneous masses of mixed country rock, of diverse sizes and shapes that have slid downslope to form chaotic accumulation of rubble and large debris flows. Debris. Locally includes small earthflows and rock falls. Thickness ranges widely; probably does not exceed 61 m (200 ft) locally.
- Ot** TILL (PLEISTOCENE)—Brown to dark-brown masses of unsorted, unconsolidated to semi-consolidated debris. Fragments range in size from clay to boulders. Glacial deposits. Characterized by lobate outlines and knob-and-kettle topography. Chiefly in Joes Valley although other deposits, here mapped as mass-wasting deposits, may have been deposits of till that have been saturated with water, mobilized, and then flowed downslope. Thickness as much as 61 m (200 ft).
- Oof** OLDER ALLUVIAL FAN DEPOSITS (PLEISTOCENE)—Gray to dark gray; thin to thick bedded; prominently crossbedded. Consists of silt, sand, granules, pebbles, and small cobbles. Includes small thin interbedded lenses of crossbedded coarse sand. Fluvial sediments. As much as 61 m (200 ft) thick. Includes the "volcanic gravels" of Gilliland (1951, p. 53).
- QUATERNARY AND TERTIARY DEPOSITS**
- OTr** RUBBLE (HOLOCENE TO OLIocene?)—Fragments, mostly of mudstone, light brown to grayish orange; unconsolidated to semi-consolidated; unbedded, well compacted debris. Composed chiefly of deposits of Quaternary age, but may include some material of Tertiary age. Thickness uncertain; may be as much as 30 m (100 ft) locally.
- OTrf** COALESCED ALLUVIAL FAN DEPOSITS (HOLOCENE TO Pliocene?)—Brown to dark brown or gray; thin to thick bedded, commonly crossbedded; unconsolidated to semi-consolidated. Consist of silt, sand, granules, pebbles, cobbles, and sparse boulders. Fluvial sediments. Formed as a result of the overlap and interfingering of adjacent alluvial fans; form broad, low sloping aprons at foot of adjacent highlands. Includes Sevier River Formation of late Pliocene and Middle Pliocene (Young and Carpenter, 1965, p. 20). Thickness uncertain; possibly as much as 30 m (100 ft) locally.
- OTrpm** PEDIMENT MANTLE (HOLOCENE TO Pliocene?)—Light brown to brown, gray, or locally reddish brown; massive to crudely bedded, unconsolidated to well-cemented. Consists of silt, sand, granules, pebbles, and small cobbles, pebbles, and boulders derived from adjacent uplands. Fluvial sediments. Surfaces are even and slope gently away from the uplands, but are somewhat deformed locally. In a few places includes gray, pink, and dark-gray to black, thin, dense limestone beds. Salt-and-pepper sandstone at base. Conglomerate lenses contain distinctive black, well-rounded chert pebbles. Fluvial deposits. Ranges in thickness from 0-50 m (160 ft) thick in this area, but as much as 305 m (1,000 ft) thick farther south.
- Tertiary Detachment Blocks**
- T(Tch)** DISPLACED BLOCK OF THE OLIocene(?) CRAZY HOLLOW FORMATION (TERTIARY)—Unbroken block of the Crazy Hollow Formation (Tch) carried "piggyback" to its present position, during either Oligocene(?) or Miocene time, on a block of the Green River Formation that slid valleyward.
- T(TTg)** DISPLACED BLOCK OF THE Eocene GREEN RIVER FORMATION (TERTIARY)—Unbroken block of the Eocene Green River Formation (TTg) that slid into its present position, during either Oligocene(?) or Miocene time, along a westward-sloping glide plane. This block has rotated and tilted somewhat in its downward movement and now overlies the country rocks discordantly.
- Tertiary Sedimentary Rocks**
- Tbk** BALD KNOLL FORMATION OF GILLILAND (1951) (OLIGOCENE?)—Light-gray to tan mudstone, claystone, and interbedded siltstone, sandstone, and a few thin dense, finely crystalline limestone beds. Locally, limestone beds are soft. Fresh-water lacustrine deposits. Probably ranges in thickness from about 244 m (800 ft) to 305 m (1,000 ft).
- Tch** CRAZY HOLLOW FORMATION OF SPIEKER (1949) (OLIGOCENE?)—Red to reddish-brown, light yellow-brown, and locally white sandstone, shaly siltstone, and some conglomerate. In a few places includes gray, pink, and dark-gray to black, thin, dense limestone beds. Salt-and-pepper sandstone at base. Conglomerate lenses contain distinctive black, well-rounded chert pebbles. Fluvial deposits. Ranges in thickness from 0-50 m (160 ft) thick in this area, but as much as 305 m (1,000 ft) thick farther south.
- Tg** GREEN RIVER FORMATION (Eocene)—Consists of a limestone unit underlain by a shale unit. Fresh-water lacustrine deposits. Millen (1982) has discussed details of the stratigraphy and petrology as exposed on the Gunnison Plateau. Total thickness of formation varies widely; probably ranges from about 150-365 m (500-1,200 ft).
- Limestone unit—Pale-yellowish-gray to yellow-brown to light-brown limestone; thin to thick bedded; even bedded. Contains thin sandstone and tuff layers. Limestones are dense, thinly laminated, and commonly oolitic. Includes thin stromatolitic limestone beds rich in ostracods. Forms resistant ledges and low cliffs.
- Shale unit—Light-green to grayish-green shale; thin bedded; fissile, somewhat calcareous. A few interbedded micritic limestones. Forms gentle slopes.
- COLTON FORMATION (Eocene)**—Commonly claystone and mudstone variegated in shades of reddish brown, light gray, or light greenish gray. Locally includes beds of yellowish-gray to yellowish-brown siltstone and sandstone, and reddish-brown conglomerate. Sparse, interlayered thin beds of clay, light gray, dense, finely crystalline limestone. Fluvial deposits. Volker (1980) has discussed details of the stratigraphy and petrology as exposed on the Gunnison Plateau. Ranges in thickness from 100-260 m (325-850 ft).

- TERTIARY AND MESOZOIC SEDIMENTARY ROCKS**
- Tkn** NORTH HORN FORMATION (PALEOCENE AND UPPER CRETACEOUS)—Mudstone, claystone, sandstone, conglomeratic sandstone, conglomerate, and sparse limestone; units alternate irregularly. Mudstones are thick bedded to massive; sandstones range from thin to thick bedded; commonly crossbedded; fine to medium grained. Limestone beds are thin and dense, locally arenaceous. Contains minor coal beds along east flank of Gunnison Plateau near Wales. Fluvial and some marine lacustrine deposits. Unstable, marked by many slumps, landslides, earthflows, and other mass-wasting deposits. Ranges in thickness from about 152-915 m (500-3,000 ft).
- MESOZOIC SEDIMENTARY ROCKS**
- Kpr** PRICE RIVER FORMATION (UPPER CRETACEOUS)—Gray to light gray conglomerate, conglomeratic sandstone, and sandstone, minor shale. Thin to thick bedded; locally massive; commonly well cemented. Coarse conglomerates contain well-rounded clasts of light-brown and purple quartzite, light-gray quartz, light-gray and black chert, and sparse dark-blue limestone. Sandstones are fine to coarse grained. Fluvial deposits. Forms steep slopes and low cliffs. In western part of area consists chiefly of conglomerate beds that probably contain equivalents of Castlegate Sandstone. Ranges in thickness from a pinch out to about 366 m (1,200 ft) locally.
- Kc** CASTLEGATE SANDSTONE (UPPER CRETACEOUS)—Sandstone, light brown to brownish gray; locally conglomeratic; irregularly bedded; massive; fine to coarse grained. Locally includes some thin, dark-gray, shaly siltstone units and some carbonaceous material. Fluvial deposits. Ranges in thickness from about 15 m (50 ft) to about 150 m (500 ft).
- Ki** INDIANOLA GROUP, UNDIVIDED (UPPER CRETACEOUS) West of Sanpete Valley—Conglomerate, reddish brown and gray, thick bedded to massive, well cemented. Consists of sand, granules, pebbles and well-rounded cobbles of white, purple, green, grayish-green, and light-brown quartzite, light-brown to light-gray chert, white quartz, and some gray to dark-blue limestone. More carbonate clasts than in Price River Formation, and less than in North Horn Formation. Fluvial (synorogenic) deposits. Ranges in thickness from 2,440-4,570 m (8,000-15,000 ft) (Speiker, 1949, p. 13).
- East of Sanpete Valley—Light-brown to yellowish-brown sandstone, shale, coal, and some conglomerate. Chiefly marine deposits. Includes four named units (in descending order) as follows: Sixmile Canyon Formation (sandstone, coal, and minor conglomerate), Funk Valley Formation (sandstone and shale), Allen Valley Shale (gray, water-lacustrine deposits), and Sanpete Formation (sandstone). About 2,250 ft thick (Speiker, 1949, p. 13, 14).
- Kbh** BLACKHAWK FORMATION (UPPER CRETACEOUS)—Sandstone, shaly siltstone, shale, carbonaceous shale, and coal. Sandstones, light gray, light brown, and brownish gray; locally reddish brown; thin to medium bedded; crossbedded; fine to medium grained. Many thin to thick coal zones in lower part; major thick coal zone at base directly overlies Star Point Sandstone. Continental and delicate deposits. Ranges in thickness from about 213 m (700 ft) to about 305 m (1,000 ft).
- Ksp** STAR POINT SANDSTONE (UPPER CRETACEOUS)—Light-brown to brown sandstone, shale, and shaly siltstone; thin to medium bedded; fine to medium grained. Consists of three sandstone units, and Sanpete Formation, of Spring Canyon, Stors, and Panther Tongues. Sandstone units separated by beds of shale and shaly siltstone. Fluvial deposits. Formation ranges in thickness from about 61-305 m (200-1,000 ft); generally about 107 m (350 ft) thick.
- Km** MANCOS SHALE (UPPER CRETACEOUS)—Consists of five members (in descending order) as follows: Masuk, Emery Sandstone, Blue Gate, Ferron Sandstone, and Tununk Member. Formation ranges in thickness from 700 m (2,300 ft) to 1,860 m (6,100 ft).
- Kmm** Masuk Member—Shale and shaly siltstone, light gray, bluish gray, and dark gray; thin to thick bedded; low thin interlayered brown sandstone beds. Includes sparse discontinuous ledges of siltified shale. About 305 m (1,000 ft) thick.
- Kme** Emery Sandstone Member—Consists of upper and lower sandstone units separated by a middle shale unit. Where exposed, member is about 87 m (280 ft) thick.
- Upper unit—Sandstone, light brown, yellowish brown, and grayish brown; thin to medium bedded, locally crossbedded; very fine grained to fine grained. Quartzose. Weathers to angular thin plates. Stands as low cliffs. Where exposed, member is about 107 m (350 ft) thick.
- Middle unit—Light-gray to gray shale, shaly siltstone, and sandstone; fissile, sandy locally; thin bedded, even bedded. Few thin interlayered sandstone beds that weather as stippled prominences. About 61 m (200 ft) thick.
- Lower unit—Sandstone, light brown, yellowish brown, and grayish brown; thin to medium bedded; very fine grained to fine grained. Quartzose. Stands as low cliff. About 15 m (50 ft) thick.
- Kmbg** Blue Gate Member—Shale and shaly siltstone, light gray, bluish gray, and gray, fissile; thin to medium bedded, sparsely interlayered thin sandstone beds. Forms low rounded hills. Resembles the Masuk and Tununk Members. As much as 61 m (200 ft) thick.
- Kmf** Ferron Sandstone Member—Consists of upper and lower sandstone units separated by a middle shale unit. Where exposed, member is about 49 m (160 ft) thick.
- Upper unit—Sandstone, buff, brown, and brownish gray; medium to thick bedded, locally massive; even bedded; very fine to fine grained. Forms cliffs. This northward. Zero to as much as 61 m (200 ft) thick in this area.
- Middle unit—Shale, light gray to gray; fissile; locally somewhat silty and carbonaceous; thin bedded; even bedded. Interleaved thin lenticular sandstone beds. Contains important minable coal beds south of this quadrangle. Ranges in thickness from 0-30 m (0-100 ft).
- Lower unit—Sandstone and shale, light brown and grayish brown; thin bedded and even bedded; crossbedded; very fine to fine grained. Consists of two informally named sandstone units (in descending order) as follows: Washboard and Clason units and a thin shale unit. Upper sandstone (Washboard unit) contains many very large, 1.5-2.4 m (5-8 ft), rounded, sandstone concretions. The lower unit ranges in thickness from about 3-13 m (10-40 ft).
- Kmt** Tununk Member—Shale and shaly siltstone, light gray to dark gray; fissile; thin to medium bedded; even bedded. Locally contains discontinuous ledges of siltified shale. Few sandstone concretions. Forms low rounded hills. Ranges in thickness from 122-198 m (400-650 ft).
- Kdc** DAKOTA SANDSTONE AND CEDAR MOUNTAIN FORMATION UNDIVIDED Dakota Sandstone (Upper Cretaceous)—Sandstone, tan to light brown; fine to medium grained; crossbedded. Contains thin discontinuous carbonaceous seams. Beach to marginal marine deposits. Thickness ranges from 0.9 m (3-30 ft).
- Cedar Mountain Formation (Lower Cretaceous)—Consists of two units. Fluvial deposits. Thickness of Cedar Mountain Formation ranges from 49-100 m (160-330 ft).
- Upper unit—Dominantly mudstone, variegated in shades of purple, red, gray, and green; massive to thick bedded. Few discontinuous sandstone lenses. Thickness ranges from 46-76 m (150-250 ft).
- Lower unit—Conglomerate and conglomeratic sandstone, gray, massive to thick bedded; crossbedded. Consists of well-rounded clasts of white quartz, black, brown, and light-gray chert, and white quartzite. Forms resistant ledges. Thickness ranges from 3-24 m (10-80 ft).
- Kju** CRETACEOUS AND JURASSIC STRATA, UNDIVIDED—Includes units assigned in the past to the qualified Morrison(?) Formation (Speiker, 1946, p. 125; 1949, p. 18). Includes conglomerate, sandstone, mudstone, and limestone, but all lithologies are not necessarily represented at any one exposure. Fluvial deposits.
- Conglomerate, reddish brown; medium bedded to massive; crossbedded. Clasts are well rounded; include light-brown quartzite, light-gray quartz, and dark-blue limestone.
- Sandstone, reddish brown to very light gray; thin to medium bedded; crossbedded; fine- to coarse-grained; quartzose.

- Mudstone**, variegated in shades of pale red, violet, and light gray; unbedded. Deposits weathered to low rounded hills and knolls. Bentonitic. Locally contains small lenses of conglomerate and sandstone, and light-gray, dense, finely crystalline freshwater limestone. Contains gastroliths which suggest that the mudstone facies is correlative with the upper member of the Cedar Mountain Formation of Early Cretaceous age (W. L. Stokes, oral communication, 1981).
- Jm** MORRISON FORMATION (UPPER JURASSIC)—Consists of Brushy Basin Member (Jmbb) overlain by the Salt Wash Sandstone Member (not exposed in this area). Fluvial deposits. Formation ranges in thickness from 107-122 m (350-400 ft).
- Jmbb** Brushy Basin Member—Claystone and mudstone, variegated, dominantly bluish gray and gray with bands of yellow, green, maroon, brick red, and purple. Bentonitic. Contains few thin limestone and sandstone beds. Limestone beds locally gray and nodular. Ranges in thickness from 46-76 m (150-250 ft).
- Jtg** TWIST GULCH FORMATION (MIDDLE JURASSIC)—Reddish brown sandstone, shaly siltstone, and shale; locally some thin interbedded light-gray laminated or crossbedded sandstone. Thin to medium bedded, even bedded; fine grained. On Gunnison Plateau, from Chats Canyon northward, thin orange-red conglomerate beds in upper part grade to red and gray massive conglomerates of Indiana Group. Marine deposits. Thickness estimated at about 914 m (3,000 ft) (Hardy, 1952, p. 23).
- Ja** ARAPIEN SHALE (MIDDLE JURASSIC)—Mudstone, calcareous, commonly light gray marked by pale-red blotches. In places wholly drab gray, elsewhere wholly reddish brown. Thin to medium bedded, even-bedded; locally amorphous. Interbedded thin, lenticular beds and seams of yellowish-gray to light-brown siltstone and sandstone. Sparse limestone beds. Contains thin beds of rock salt (halite), gypsum, and other evaporites. Selenitic crystals abundant on many outcrops. Marine saline-basin deposits. Formation is complexly deformed, and shows signs of intense compression. Weathers to badland topography. Thickness uncertain because of intense deformation; estimates range from about 1,220 m (4,000 ft) to as much as 3,960 m (13,000 ft) (Speiker, 1949, p. 17; Gilliland, 1948, p. 30; Hardy, 1949, p. 16, 17).
- Jtc** TWIN CREEK LIMESTONE (MIDDLE JURASSIC)—Includes seven members (in descending order) as follows: Giffels Creek, Leeds Creek, Watton Canyon Boundary Ridge, Rich, Slidereck, and Gypsum Spring. Only Leeds Creek Member is believed to be exposed in this quadrangle, chiefly north of Little Salt Creek along the west flank of the Gunnison Plateau. Marine deposits. Light to dark-gray, argillaceous limestone; dense; thin to medium bedded; even bedded. Locally complexly folded, and badly fractured. Complex relations between the Arapian Shale and the Twin Creek Limestone have been discussed by Sprinkel (1982). Thickness uncertain, possibly 150-305 m (500-1,000 ft).
- QUATERNARY, TERTIARY, AND MESOZOIC SEDIMENTARY ROCKS**
- T(Ja)** INTRUSIVE MASSES OF THE ARAPIEN SHALE (QUATERNARY?) TO MIDDLE JURASSIC?—Probably the salt component (and possibly other evaporites) in the Arapian Shale (see above) has been moving ever since they were deposited during the Middle Jurassic (Witkind, 1982). Some of this movement has been a slow, almost imperceptible upwelling. Every now and then, however, the salt seemingly has surged upward rapidly, and in so doing has forced the overlying mudstones of the Arapian Shale to bow up the country rock to form elongate, narrow diapiric folds. Subsequent removal of the salt resulted in collapse of the upwells. Thus this major diapiric upwelling of the salt may have occurred during the Late Cretaceous, early Paleocene, and the late(?) Oligocene or Miocene. A localized upward surge of the salt likely deformed semi-consolidated sediments probably during the Pleistocene. The formation, thus, has several ages; its depositional age is Middle Jurassic. Its emplacement age (age(s)) has changed repeatedly. Symbol (T(Ja)) reflects both the depositional age (Ja), and the time of major emplacement (T), that is, Tertiary.
- EXTRUSIVE IGNEOUS ROCKS AND THEIR PRODUCTS**
- TOb** OLDER BASALT FLOW(?) (MIOCENE?)—Basalt, dark gray to black, weathers reddish brown; fine grained. Only a small remnant preserved south of Willow Creek Reservoir, about 1.8 m (6 ft) thick.
- Tvg** GOLDENS RANCH FORMATION OF MUESSIG (1951) (OLIGOCENE?)—Light-gray to gray, fine grained, silty porphyritic. Phenocrysts (10 percent) of rock tuff plus a lenticular, light-gray limestone. Conglomerate, brown to dark-brown; thin to thick bedded; locally massive; well-rounded cobbles of andesite and quartzite; includes a few sandstone beds. Interbedded volcanic flows in western part of area. Fluvial sediments rich in clasts of volcanic origin. Radiometric ages of about 32 m.y. determined on tuffs. Thickness uncertain; probably about 305 m (1,000 ft).
- INTRUSIVE IGNEOUS ROCKS OF THE LEVAN AREA (JOHN, 1972)**
- Tim** MONZONITE (OLIGOCENE TO UPPER Eocene?)—Dike(?) Light gray to gray; fine grained, silty porphyritic. Phenocrysts (10 percent) consist of plagioclase (oligoclase-andesine) (about 8 percent), hornblende (about 1 percent), and biotite (about 1 percent). Groundmass includes plagioclase, orthoclase, and sparse quartz. Accessory minerals are apatite, zircon, and magnetite.
- Tmp** MONZONITE PORPHYRY (OLIGOCENE TO UPPER Eocene?)—Dikes and two small stocks. Light gray to gray; aphanitic groundmass, porphyritic. Phenocrysts (32 percent) consist of plagioclase (andesine) (about 20 percent), hornblende (about 10 percent), and biotite (about 2 percent). Groundmass consists of plagioclase, orthoclase, and sparse quartz. Accessory minerals consist of apatite, magnetite, sphene, and zircon.
- TThp** HORNBLENDE MONZONITE PORPHYRY (OLIGOCENE TO UPPER Eocene?)—Small irregular-shaped intrusions. Light gray to reddish gray; aphanitic groundmass, porphyritic. Phenocrysts (63 percent) consist of plagioclase (andesine-labradorite) (about 50 percent), augite (about 15 percent), hornblende (about 3 percent) and biotite (about 1 percent); locally some olivine. Groundmass consists of essentially the same minerals. Accessory minerals include apatite, sphene, and magnetite.
- Td** PORPHYRYTIC DIABASE (OLIGOCENE TO UPPER Eocene?)—Dikes(?) Greenish gray; porphyritic. Phenocrysts (70 percent) of rock consist of plagioclase (andesine-labradorite) (about 50 percent), augite (about 15 percent), hornblende (about 3 percent) and biotite (about 1 percent); locally some olivine. Groundmass consists of essentially the same minerals. Accessory minerals include apatite

INTRODUCTION

The U.S. Geological Survey is engaged in a broad program of field studies designed to present the geologic framework of the United States on easily read topographic maps. The maps selected as a base for these geologic data are part of the Army Map Service (AMS) series of 1° x 2° quadrangles at a scale of 1:250,000. The Price, Utah, AMS quadrangle is one of these maps (see fig. 1). For certain areas, however, chiefly those sectors of the country involved in the U.S. Geological Survey's Coal Exploratory Program, the geologic data are being compiled on newly developed base maps at a scale of 1:100,000. On these new maps the configuration of the land is shown by metric contours having a 50-meter contour interval. One of these new maps has been used as a base for this geologic map of the Manti 30' x 60' (1:100,000) quadrangle, the southwestern quadrangle of the four quadrangles that together make up the Price 1° x 2° quadrangle.

THE PRICE, UTAH, 1° x 2° AMS (1:250,000) QUADRANGLE

The geologic data compiled on the Manti quadrangle are but a part of a much larger geologic pattern best displayed on the Price, Utah, 1° x 2° quadrangle. The Price quadrangle includes within its borders parts of three major physiographic elements: the Colorado Plateaus, the Basin and Range, and the Middle Rocky Mountains Provinces. Most of the quadrangle, including the central and eastern parts, overlies the west margin of the Colorado Plateaus. Within this part of the plateaus are the southern edge of the Uinta Basin (expressed as the southward-facing, sinuous escarpments formed by the Book and Roan Cliffs), the northern part of the Canyonlands section (expressed by the northeast-trending San Rafael Swell), and the northernmost of the High Plateaus of Utah (the Wasatch Plateau). The western part of the sheet includes the east edge of the Basin and Range Province (the Great Basin). The join between the Colorado Plateaus and the Basin and Range Provinces trends northward through western Sanpete County and is a transition zone that is gradational in both geology and topography. A small wedge of the Middle Rocky Mountains Province—the southern Wasatch Mountains—dominates the northwest corner of the Price quadrangle.

THE MANTI, UTAH, 30' x 60' (1:100,000) QUADRANGLE

About half the Manti quadrangle (see fig. 2) is underlain by the Wasatch Plateau, a high, forested tableland composed of near-horizontal to gently warped Cretaceous and Tertiary strata. The east edge of the Wasatch Plateau overlooks Castle Valley, an arid, relatively barren, sage-covered lowland cut into poorly resistant strata on the west flank of the San Rafael Swell. By contrast, the west edge of the Wasatch Plateau looks across Sanpete Valley, a rich agricultural and ranching area, toward another tableland known to geologists as the Gunnison Plateau, to local inhabitants as the West Mountain, and to some Federal agencies as the San Pitch Mountains. The fact that the Gunnison Plateau, consists of the same stratigraphic units as are exposed in the Wasatch Plateau, and is essentially a gently warped upland much like the Wasatch Plateau, suggests that it is an integral part of the Colorado Plateau. The Gunnison Plateau, however, is marked locally by structural complexities that seem to relate it to the Basin and Range structures so common farther west. We are uncertain, thus, just where the join between the two provinces should be in this part of central Utah. West of the Gunnison Plateau, and separated from it by the broad valley of the Sevier River, are the Valley Mountains, a low, elongate, north-trending tilted fault block that is much like the block-faulted ranges that make up the Basin and Range Province of western Utah and eastern Nevada. Because both the Wasatch and Gunnison Plateaus are plateaus, whereas the Valley Mountains are a tilted fault block, we tend, arbitrarily, to include the Wasatch and Gunnison Plateaus with the Colorado Plateaus Province.

We tentatively propose, thus, that the physiographic join between the Colorado Plateaus and the Basin and Range Provinces extends southward along the west flank of the Gunnison Plateau, flexes sharply to the southeast at the south end of the Gunnison Plateau, and continues southward along the west flank of the Wasatch Plateau.

In summary, the Manti quadrangle can best be visualized as consisting of parts of four physiographic elements. From east to west, these are: (1) the west flank of the San Rafael Swell, (2) the south half of the Wasatch Plateau, (3) the south half of the Gunnison Plateau (all three within the Colorado Plateaus Province), and (4) the east half of the Valley Mountains (essentially the easternmost range of the Basin and Range Province).

COLORADO PLATEAUS SEGMENT

San Rafael Swell

West-dipping shale and sandstone beds of Cretaceous age (Mancos Shale), exposed along the east edge of the Manti quadrangle, reflect the northwest flank of the San Rafael Swell. The swell is an elongate, oval, domelike structure, about 115 km (70 mi) long, and 50 km (30 mi) across (at its maximum width) that trends about N. 30° E. through Emery and Wayne Counties. The major axis of the San Rafael Swell is about 32 km (20 mi) east of the Manti quadrangle. The influence of the swell is strong: former near-horizontal Cretaceous and Tertiary strata were domed as the swell developed, and these units, now much eroded, express the general outline of the upwarp. The northern half of the San Rafael Swell (eastward beyond the Manti quadrangle) is bordered by a lowland underlain by Cretaceous shale beds, which are truncated in many places by pediment surfaces mantled by extensive deposits of sand and gravel. Rising abruptly above this sage-covered lowland are sandstone and shaly siltstone beds, younger but also of Cretaceous age, that crop out as steep, sinuous escarpments. On the east and north (beyond the Manti quadrangle) these cliffs are known as the Book Cliffs; on the west (and partly within the Manti quadrangle) they are unnamed but form the east flank of the Wasatch Plateau. Thick, well-developed coal beds (in the Blackhawk Formation of Late Cretaceous age) crop out along the cliffs and are mined extensively.

Wasatch Plateau

The Wasatch Plateau is a flat-topped mass about 130 km (80 mi) long and 40 km (25 mi) wide, and it appears as a huge upland that trends about N. 20° E. separating Sanpete Valley on the west from Castle Valley on the east. The top of the plateau is at an altitude of about 3,050 m (10,000 ft). In striking contrast to the clifflike aspect of the east flank of the Wasatch Plateau, the west flank of the plateau is a continuous westward-facing downwarp—the Wasatch monocline—that extends for almost the full length of the plateau. Westward-flowing consequent streams on the monocline have locally cut through the mantle of tilted beds to form deep, serpentine canyons that extend far back toward the crest of the plateau.

The Wasatch Plateau is underlain by flat-lying Cretaceous and Tertiary beds, most of which are well exposed either in the cliffs along its east flank or in the canyons cut into the Wasatch monocline. These rocks are broken by north and northeast-trending high-angle normal faults many of which are paired to form narrow, elongate grabens. Of these, one of the more spectacular is the Joes Valley graben, which traverses the full extent of the Manti quadrangle.

Gunnison Plateau

The Gunnison Plateau is a north-trending, oval upland composed of sedimentary rocks that have been warped into a southward-plunging syncline. Completely encircled by valleys, the plateau is about 61 km (38 mi) long, and about 23 km (14 mi) wide near its midpoint. The crest of the Gunnison Plateau reaches altitudes of about 2,450 m (8,000 ft), and thus is somewhat lower than the Wasatch Plateau. The southern end of the Gunnison Plateau merges with the lowlands of the Sanpete and Sevier Valleys, but the plateau gradually gains height northward, and its northern end, marked by steep, precipitous cliffs, towers above the adjacent valley floors.

The rocks that form the plateau are much faulted, although on the whole they are but gently warped. Only along the east edge of the plateau are these rocks intensely deformed, in a zone of complex structures that extends as a narrow welt, 1–2 km (1/2–1 mi) wide, from near Wales southward to the southern edge of the plateau (Spieker, 1949; Witkind, 1982). Comparable large-scale structures have not been found along the west flank of the plateau, although locally the rocks there are much deformed (Witkind, 1983).

BASIN AND RANGE SEGMENT

Valley Mountains

If the Gunnison Plateau is correctly assigned to the Colorado Plateaus Province, only a small part of the Basin and Range province is represented in the Manti quadrangle—essentially the east flank of the north-trending Valley Mountains. The Valley Mountains are an eastward-titled fault block whose west flank (west of the quadrangle boundary) is bounded by a high-angle normal fault that trends north and dips steeply to the west. West of the Valley Mountains are other ranges, such as the Canyon and Pavant Ranges, which also trend north and which also are bounded along one or both margins by high-angle normal faults. From the Valley Mountains westward, therefore, the geologic pattern is one of alternating north-trending basins and parallel block-faulted ranges—typical Basin and Range topography.

THE TRANSITION ZONE

Structurally, the south-trending transition zone between the Colorado Plateaus and the Basin and Range Provinces dominates the Manti quadrangle. Structures transitional between the two provinces extend from the crest of the Wasatch Plateau westward to and possibly beyond the Valley Mountains.

The same sedimentary sequence that underlies the Wasatch Plateau is found in the Gunnison Plateau and in the Valley Mountains. Locally, in all three areas, the rocks are intensely deformed; complex structures abound. In the Wasatch Plateau these structures are best exposed at and near the mouth of Sixmile Canyon, near Sterling (Spieker, 1949, p. 49–51). The east flank of the Gunnison Plateau is marked by an extensive belt of complexly deformed rocks (Spieker, 1949, p. 72–74). Comparably deformed rocks are exposed along the northeast flank of the Valley Mountains. Beds are vertical to steeply overturned; locally, one or more angular unconformities break the sedimentary sequence. In several places two profound angular unconformities are exposed in a single outcrop, but surprisingly these unconformities are not extensive laterally. Strata that are separated by a striking angular unconformity, when traced laterally, rapidly approach parallelism and become conformable in distances as short as three-fourths of a kilometer (half a mile). Overturned beds become vertical, then upright, decrease in dip and in unusually short distances are near-horizontal and conform with both overlying and underlying units. The intense deformation is extremely localized.

Interpretations

This remarkable deformation has been interpreted in different ways. The late Professor E.M. Spieker and his many graduate students have attributed the deformation to multiple episodes of tectonic disturbance. In all, some 14 such episodes have been postulated (Spieker, 1949, p. 78). Subsequent work has suggested an additional two episodes to make a total, then, of 16 tectonic pulses (Gilliland, 1948, p. 74; 1951, p. 72; 1952, p. 1461). Of these, eight were orogenic (compressive) movements that have occurred since Jurassic time (Spieker, 1949, p. 40). The remainder were chiefly episodes of normal faulting, warping, or monoclinical flexing. Other workers in the area, however, chiefly Stokes (1952, 1956), Moulton (1975), Baer (1976), and Witkind (1982), have suggested that much of the deformation is more reasonably explained by multiple episodes of salt diapirism.

Regrettably, the field evidence is equivocal; strong arguments can be made for either interpretation. Weiss, one of the co-authors of this map, tends toward multiple tectonism as an explanation for most of the deformation. He sees much of the structural complexity as being thrust related and considers that salt diapirism played a minor role. Weiss believes the

question is moot on the basis of surface geology, and that multiple tectonism cannot be fully discredited until diapirism can be demonstrated by subsurface data. Weiss attributes the intense deformation that marks the east flank of the Gunnison Plateau to an antiformal welt that lay just east of the Gunnison Plateau in latest Cretaceous and Paleocene time, and which subsequently bent the edge of the plateau block toward the west (Weiss, 1982). The welt may have been caused by compression from the east, diapirism, or both. Witkind, another co-author of this map, is a strong advocate of salt diapirism, and accordingly he attributes the bulk of the deformation, as well as the extreme localization of the deformed structures, to recurrent salt movement—in essence, the repeated growth and collapse of many linear salt diapirs. In his view, eastward-directed compressive impulses probably played a significant role in that, initially, they triggered the salt into upward movement. He sees this upwelling of the salt as responsible for most of the local deformation. Thus, he attributes the westward bend of the east edge of the Gunnison Plateau to the repeated upwelling of a salt diapir concealed beneath Sanpete Valley (Witkind, 1982, p. 26–27). As the diapir welled upward, vertical stresses were translated laterally into horizontal stresses, and it is the westward-directed component of these horizontal stresses that flexed the beds to the west.

FAULTS

A series of high-angle normal faults, of diverse orientation, cut and offset the rocks in the Manti quadrangle. In the Wasatch Plateau these faults form a simple pattern of breaks that trends generally north and northeast. By contrast, in the Gunnison Plateau, the pattern is much more complex. In addition to the north- and northeast-trending faults are a series of northwest- and westward-trending cross faults that break the rocks of the plateau into small, almost disjointed blocks. This same complex fault pattern persists in the Valley Mountains but is not as intense.

WASATCH PLATEAU

The crest and west flank of the Wasatch Plateau are broken by a system of high-angle normal faults that range in trend from about N. 10° W. to about N. 30° E. This fault system does not appear to extend eastward into the San Rafael Swell, for only a few northerly trending faults break the rocks of the swell; there, the fault pattern is dominated by west and northwest-trending faults.

The faults on the Wasatch Plateau are remarkably straight, and commonly persist as single breaks, or very narrow fault zones, traceable for very long distances. Some, such as those that bound the Joes Valley graben, are narrow fault zones that extend for the full length of the Manti quadrangle. The continuity of the faults is masked on the ground, however, by the fact that the faults are concealed in the valleys beneath mass-wasting deposits. In most places the faults are visible only along valley walls and on the interfluvies between the valleys.

A few of the faults show opposite directions of displacement along their lengths. Thus, for example, at the northern end of such a fault, commonly known as a "scissors fault", the block west of the fault is downthrown. As the fault is traced southward the amount of displacement lessens, reaches a null point, and then changes, with the end result being that at the fault's southern end the block east of the fault is downthrown.

Many of the high-angle normal faults are paired to form grabens that exceed 65 km (40 mi) in length. In general, the grabens maintain a relatively constant width of about 3 km (2 mi) although locally they narrow to about half a kilometer (a quarter of a mile). Structural relief differs from graben to graben. In some grabens it is as low as 100 m (several hundred feet); elsewhere it is hundreds of meters. Spieker (1949, p. 43) suggested stratigraphic separation of 610–915 m (2,000–3,000 ft) in the Joes Valley graben, one of the major structural elements in the quadrangle, and in excess of 1200 m (4,000 ft) in the nearby Musinia graben.

In many grabens the downthrown blocks are unbroken, but locally they are cut by a series of small internal faults that trend parallel to the major faults that bound the graben. In places, the many faults have so disrupted and broken the coal beds that underlie much of the Wasatch Plateau that these beds cannot be mined (Doelling, 1972a, p. 83).

The origin of the faults, and thus, of the grabens, is uncertain. The faults may reflect widespread crustal spreading stemming from an episode of extensional tectonics that has dominated the western interior of the United States since Miocene time. If so, the faults are tectonic in origin. By contrast, the faults may be related, in one way or another, to the salt that underlies much of the Sanpete-Sevier Valley area (Stokes, 1952; Stokes and Holmes, 1954, p. 40). Moulton (1975, fig. 19) implies that many of the faults that break the crest of the Wasatch Plateau do not extend below the base of the salt-bearing beds. This suggests that at least some of the faults and grabens may be genetically related to withdrawal of the salt. Not all faults, however, are so shallow; several appear to extend into the presalt rocks (Moulton, 1975, fig. 19).

If some of the grabens are indeed salt-induced structures, possibly they formed in response to an outward (valleyward) flowage of the salt, much as postulated by Baker (1933, p. 74) to explain the development of the grabens that are so common in the Paradox Basin of southeastern Utah and southwestern Colorado.

Differential flowage of salt may also explain the anomalous pattern of displacement that characterizes the scissors faults.

GUNNISON PLATEAU

As on the Wasatch Plateau, many of the north-trending faults that break the rocks on the Gunnison Plateau are paired to form grabens. The linear trends of the grabens, however, are masked by a maze of smaller faults that break and disrupt the downdropped blocks in the grabens. Two conspicuous examples of this are at the southern end of the plateau. Along the southwestern flank of the Gunnison Plateau, essentially between Mellor and Hells Kitchen Canyons, the downdropped block in a large north-trending graben consists of a chaotic

jumble of blocks. A comparable graben, although one in which the downdropped block is not as badly broken, begins near the southeastern end of the plateau and extends northwesterly across the crest of the plateau. In sharp contrast to these north-trending grabens, however, is a transverse graben that trends west and breaks the rocks along the east flank of the Gunnison Plateau between Maple and Rock Canyons (directly west of Ephraim).

All three grabens and many of the other faults in the Gunnison Plateau may be salt related. Possibly the causative salt originally bowed up the rocks to form elongate, linear folds; subsequent removal of the salt caused the axial parts of folds to collapse along boundary faults and form the grabens.

The striking difference in intensity of faulting, when the Gunnison and Wasatch Plateaus are compared, may reflect the fact that the Gunnison Plateau appears to have been subjected to compressive forces from both the west and east, possibly by salt diapirs, which co-author Witkind believes about its west, east, and north flanks. The synclinal aspect of Gunnison Plateau, and its southward plunge, may stem from the upward movement of these diapirs.

VALLEY MOUNTAINS

The crest of the Valley Mountains is broken by a north-trending graben known as Japanese Valley. Both east and west of the graben are other high-angle normal faults that also trend north. In general, those faults east of the graben are marked by downthrow to the west—toward the graben; those few faults west of the graben (beyond the confines of the Manti quadrangle) are marked by downthrow to the east—again toward the graben. The implication is strong that the graben is a collapse feature possibly related to withdrawal of salt. Test wells have indicated that the Valley Mountains are underlain by the Arapian Shale (the salt-bearing unit), but as yet none of the wells has penetrated salt. It may well be that whatever salt did underlie the Valley Mountains has been dissolved and removed.

MODERN FAULT SCARPS

A series of modern fault scarps are along both flanks of the Gunnison Plateau. In all localities, these scarps cut and offset surficial deposits, implying that the causative faults have moved in the recent past. The fault scarps trend northerly, dip valleyward, and range in height from 0.3–8 m (1–25 ft). Those scarps along the west flank of the Gunnison Plateau clearly reflect movement along the Wasatch fault zone, one of the major, young, active fault zones in the United States. The fault along the east flank of the Plateau, known as the Gunnison fault, cuts and offsets an old alluvial fan that heads in the mouth of Wales Canyon. The new scarp, in turn, is now partly concealed by a younger alluvial fan which also heads in Wales Canyon.

The origins of the faults are uncertain. The Wasatch fault has always been considered to be an extensional fault; hence, it may have formed in response to tectonic impulses. Movement along the fault would likely result in a major earthquake that would cause much damage to manmade features. By contrast, the close parallelism between the trace of the Gunnison fault and the suspected salt diapir believed to be concealed beneath Sanpete Valley, suggests that withdrawal of salt may have played some role in the development of that fault. If so, the Gunnison fault, although a geologic hazard, may not pose as much of a risk as the Wasatch fault.

WATER RESOURCES

Details about the surface-water resources of the Manti quadrangle are contained in a companion publication, U.S. Geological Survey Miscellaneous Investigations Series Map I-1482 (Price, 1983).

ECONOMIC DEPOSITS

Materials of economic interest in and near the Manti quadrangle include mineral fuels such as coal and hydrocarbon gas. Pools of oil and gas also may be in the area. Several fields producing carbon-dioxide (CO₂) gas are northeast of the quadrangle. Nonmetallic mineral deposits in the quadrangle include gypsum, salt and brines, sand and gravel, building stone, and limestone. Metallic mineral deposits are sparse; some zinc ore was mined near Redmond.

MINERAL FUELS

Coal

Vast amounts of coal underlie the Wasatch Plateau with some of it accessible along the Plateau's east flank (Spieker, 1931; Doelling, 1972a, 1972b). Of the six coal fields in and near the Price 1° x 2° quadrangle (the Wasatch Plateau, Book Cliffs, Salina Canyon, Mount Pleasant, Wales, and Sterling fields), only the Book Cliffs and Salina fields are beyond the boundaries of the Manti quadrangle. Of those fields, or parts of fields, within the quadrangle, apparently only the Wasatch field contains deposits of coal that are economically significant at present. Although seams and beds of coal crop out west of the Plateau, in Sanpete Valley near Wales, Mount Pleasant, and Sterling, these beds, for the most part, are thin (generally about 0.3 m (1 ft) thick), discontinuous, and of poor quality. The early settlers mined these beds until the thicker, higher quality, and more continuous beds that are exposed along the east edge of the Wasatch Plateau were discovered. These latter beds are considered to contain some of the best coal in Utah (Doelling, 1972b, p. 554). Some test holes by the U. S. Bureau of Mines, however, indicated that at least three minable coal beds, ranging from 1–2 m (4–6 ft) in thickness, are in the subsurface in the Mount Pleasant area at depths between 290 m (955 ft) and 350 m (1150 ft) (Doelling, 1972b, p. 30).

Coal in the Manti quadrangle is contained chiefly within five units of Tertiary and Cretaceous age: the North Horn, Blackhawk, and South Flat (probably correlative with the Blackhawk) Formations, the Sixmile Canyon Formation of the Indianola Group, and the Ferron Sandstone Member of the Mancos Shale. Of these, only the Blackhawk Formation

contains thick (as much as 8 m (25 ft) thick) coal beds that are either exposed or are buried beneath an overburden of 305 m (1,000 ft) or less. The Blackhawk Formation crops out all along the east flank of the Wasatch Plateau, and it is likely that its coal beds underlie much of the plateau. Comparable coal beds may be contained within the Ferron Sandstone Member, but if so these are so deeply buried they probably will be difficult to mine.

Surface investigations and drilling have determined that as many as 22 coal beds in the Blackhawk Formation are thick enough to be named (Doelling, 1972a, p. 85). Some of the thickest and most persistent coal beds are the Hiawatha, Castlegate A, Bear Canyon, Blind Canyon, Wattis, Upper Ivie, and Ivie beds (Doelling, 1972a, p. 85). Recent work by U. S. Geological Survey geologists, notably J. D. Sanchez, L. F. Blanchard, E. E. Ellis, and T. L. Brown, has suggested that these coal beds, rather than being laterally uniform masses, are best viewed as coal zones—groups of alternating discontinuous beds of coal separated by sandstone and mudstone units. In general, each zone is separated from overlying and underlying coal zones by sandstone-rich strata, 2–6 m (5–20 ft) thick, that contain thin coal seams and carbonaceous material but are free of thick coal beds. Sanchez and his colleagues (Sanchez, Blanchard, and August, 1983) have grouped the various coal beds into a series of discrete, named coal zones. Within the Manti quadrangle, these workers recognize six such discontinuous coal zones in the lower part of the Blackhawk Formation; from oldest to youngest, these are (1) the Hiawatha, (2) Acord Lakes (which is correlative with the Muddy No. 1 and Muddy No. 2 coal zones), (3) Axel Anderson, (4) Cottonwood, (5) Blind Canyon, and (6) Bear Canyon coal zones. Within each coal zone the coal beds range in thickness from several centimeters (one inch) to as much as 8 m (25 ft).

The rank of most of the coal beds is high-volatile B bituminous as calculated from the Parr formulas of the American Society for Testing and Materials (1978) (J. D. Sanchez, oral commun., 1983).

Details about these coal zones, including analyses of the moisture, fixed carbon, ash, and volatile contents of the contained coal beds, and the relations between the various coal zones are contained in a series of U.S. Geological Survey Coal Investigations maps (Sanchez and Brown, 1983; 1986; 1987; Brown and Sanchez, 1986).

Large, modern mines dug into those coal beds of the Blackhawk exposed along the east flank of the Wasatch Plateau have just begun to tap the large coal reserves that underlie the plateau. In the Manti quadrangle, most of the more productive mines are along the north and east flanks of East Mountain (northwest of Castle Dale). The bulk of the coal mined is transported via truck to coal-fired power plants constructed at Huntington (Huntington Plant) and Castle Dale (Hunter Plant); much of the electrical power produced by these two plants is transmitted out of state.

The presence of these rich coal beds has led to tentative plans for the construction of other coal-fired power plants in and near the Price quadrangle; five are planned for areas east of the Wasatch Plateau, and three for areas west of the Plateau (Witkind, 1979). The Intermountain Power Plant (IPP), now under construction near Lynndyl (west of the Wasatch Plateau), is one of these plants; when completed it will be the largest coal-fired power plant in the world.

Reserves

As a consequence of the increased interest in coal as a fuel, the east edge of the Wasatch Plateau has been drilled extensively in an attempt to determine reserves in the Wasatch Plateau coal field. On the basis of these data, coal reserves identified in that part of the Wasatch Plateau coal field within the Manti quadrangle total about 2 billion short tons (J.D. Sanchez, oral commun., 1983).

Principal coal reserves in the combined Salina Canyon, Mount Pleasant, Wales, and Sterling fields are about 350 million short tons (Doelling, 1972a, p. 88).

Hydrocarbon gas

Significant amounts of hydrocarbon gas have been produced from small structures that underlie the Wasatch Plateau (Walton, 1963). The most important is the Clear Creek structure, a fault segment, near Scofield Reservoir, north of the Manti quadrangle. Domelike structures, from which minor amounts of hydrocarbon gas have been produced, are in and near the northeast corner of the Manti quadrangle, but of these only the Flat Canyon structure (directly east of Upper Joes Valley) is within the quadrangle. Production is from both the Dakota Sandstone, and the Ferron Sandstone Member of the Mancos Shale (Walton, 1963, p. 345).

Carbon-dioxide gas

Pure carbon-dioxide gas has been produced from wells drilled into structures northeast of the Manti quadrangle. The Iarnham dome, east of Wellington, contains CO₂ gas with major production from the Navajo Sandstone (Peterson, 1961). Somewhat closer to the Manti quadrangle is the Gordon Creek field southeast of Scofield Reservoir. The field is on the faulted Gordon Creek anticline, and production is from a series of beds that include the boundaries of the Manti quadrangle. Of those fields, or parts of fields, within the quadrangle, apparently only the Wasatch field contains deposits of coal that are economically significant at present. Although seams and beds of coal crop out west of the Plateau, in Sanpete Valley near Wales, Mount Pleasant, and Sterling, these beds, for the most part, are thin (generally about 0.3 m (1 ft) thick), discontinuous, and of poor quality. The early settlers mined these beds until the thicker, higher quality, and more continuous beds that are exposed along the east edge of the Wasatch Plateau were discovered. These latter beds are considered to contain some of the best coal in Utah (Doelling, 1972b, p. 554). Some test holes by the U. S. Bureau of Mines, however, indicated that at least three minable coal beds, ranging from 1–2 m (4–6 ft) in thickness, are in the subsurface in the Mount Pleasant area at depths between 290 m (955 ft) and 350 m (1150 ft) (Doelling, 1972b, p. 30).

Coal in the Manti quadrangle is contained chiefly within five units of Tertiary and Cretaceous age: the North Horn, Blackhawk, and South Flat (probably correlative with the Blackhawk) Formations, the Sixmile Canyon Formation of the Indianola Group, and the Ferron Sandstone Member of the Mancos Shale. Of these, only the Blackhawk Formation

contains thick (as much as 8 m (25 ft) thick) coal beds that are either exposed or are buried beneath an overburden of 305 m (1,000 ft) or less. The Blackhawk Formation crops out all along the east flank of the Wasatch Plateau, and it is likely that its coal beds underlie much of the plateau. Comparable coal beds may be contained within the Ferron Sandstone Member, but if so these are so deeply buried they probably will be difficult to mine.

Source rocks

Possible source rocks for oil and gas include the Mancos Shale of Late Cretaceous age, the Manning Canyon Shale of Late Mississippian and Pennsylvanian age, and the Arapian Shale of Middle Jurassic age.

Three exploratory wells drilled near Moroni in Sanpete Valley penetrated the entire thickness of the Mancos Shales, composed chiefly of black, marine, sedimentary rocks. The easternmost test well, Hanson Oil Corporation well No. 1 A-X Moroni (SE, NW, sec. 14, T. 15 S., R. 3 E.) cut about 2,135 m (7,000 ft) of Mancos strata. About 2.4 km (1.5 mi) to the southwest the Tennessee Gas Transmission Company's J.W. Irons well No. 1 (C, SE, NE, sec. 16, T. 15 S., R. 3 E.) penetrated about 1,700 m (3,500 ft) of Mancos. And the westernmost well, the Phillips Petroleum Company's well No. 1 Price-N (SE, SE, sec. 29, T. 15 S., R. 3 E.) cut only about 610 m (2,000 ft) of Mancos Shale. Seemingly, the Mancos Shale thins markedly westward, implying a shoaling of the Mancos sea near the east edge of the Gunnison Plateau. How much of the Manti quadrangle is underlain by the Mancos is uncertain; we suspect that it underlies most of the Wasatch Plateau and Sanpete Valley, at the very least.

The Manning Canyon Shale is another black, marine sedimentary unit that may underlie the area at depth. This shale, like the Mancos, is rich in an oil-generative type of organic material, and so a most suitable source rock. Although not exposed in the Manti quadrangle, the Manning Canyon is a conspicuous unit farther to the north where it is part of the Charleston-Nebo thrust plate that forms the Mt. Nebo massif.

The Arapian Shale may be a source rock. Recently Kirkland and Evans (1981) suggested that calcareous mudstones deposited in highly saline, marine evaporitic basins may be rich source rocks. Considerable uncertainty exists, however, about the suitability of the Arapian Shale as a source rock for oil and gas because of its apparent low content of total organic carbon (TOC) (R. J. Coskey, Forest Oil Corporation, oral commun., 1982).

Reservoir rocks

Appropriate reservoir rocks are believed to be within the Mancos Shale, chiefly the Ferron Sandstone Member and sandstone units within the Emery Shale Member. Among the other beds in the stratigraphic section that have served as reservoir rocks for either hydrocarbon or carbon dioxide gas are the Dakota Sandstone, the Navajo Sandstone, the Sinbad Limestone Member of the Moenkopi Formation, and the "Coconino Sandstone" (probably the Cedar Mesa Sandstone of the Cutler Formation). Significantly, these beds have served as good reservoir rocks mainly because they are strongly fractured, and apparently not because of unusually high primary porosity. How severely a unit has been fractured may determine its suitability as a reservoir rock.

Structural traps

If the diapiric concept as proposed by Witkind (1982) is valid, structural traps may be adjacent either to the salt diapirs or to the mudstone sheaths (composed of Arapian Shale strata) that are integral parts of the diapiric folds. Presumably as the salt surged upward it raised the enveloping mudstones of the Arapian Shale, and these, in turn, bowed up the adjacent country rocks. The traps formed between the upturned country rocks and either the intrusive salt or the mudstones would seem suitable sites for the accumulation of oil and gas.

NONMETALLIC MINERAL DEPOSITS

Gypsum deposits

Extensive gypsum deposits are in the Arapian Shale; most of the commercial deposits are concentrated either south of the Manti quadrangle between Salina and Sigurd, or north of the quadrangle at or near the mouth of Salt Creek near Nephi. The mined gypsum is transported by truck to several gypsum plants at Sigurd where it is fabricated into plaster board. Small pods and lenses of gypsum are found, however, in most outcrops of the Arapian Shale; selenite crystals are common on outcrops.

In the Manti quadrangle the Arapian Shale crops out in three localities: along the west flank of the Gunnison Plateau, along the east side of Sevier Valley south of Sterling, and along the west side of Sevier Valley near and north of Redmond. Only farther north, however, near Nephi (beyond the Manti quadrangle) and in adjacent areas south of Nephi, is much gypsum seen in outcrop, where it occurs in disconnected pods and sheets. This scattered occurrence regionally may be misleading for extreme irregularity of thickness, width, and length is characteristic of many gypsum deposits. What may appear as a thin pod of gypsum on the outcrop may thicken and widen rapidly in the subsurface to become a significant deposit. In general, we estimate that the range in thickness of a gypsum lens is from about 15 m (50 ft) to as much as 105 m (350 ft). In like fashion the width and length are also highly irregular; we estimate that some lenses are as much as 305 m (1,000 ft) wide, and about 210 m (700 ft) long.

Salt and brines

Very little rock salt (halite) is exposed at the surface, although much salt probably underlies the Sanpete-Sevier Valley area. So, for example, the Phillips Price-N well, near Moroni in Sanpete Valley, penetrated about 610 m (2,000 ft) of salt. Farther west (beyond the Manti quadrangle), near Yuba Dam in Sevier Valley, Placid Oil Company's Monroe 13-7 well (SW, NE, sec. 13, T. 16 S., R. 2 W.) cut about 215 m (700 ft) of salt. Chevron Oil Company's Chris Canyon well (NE, SW, sec. 33, T. 16 N., R. 1 E.), near the southern end of the Gunnison Plateau, cut through about 245 m (800 ft) of salt (L. A. Standlee, Chevron Oil Co., oral commun., 1982). The salt is contained within the Arapian Shale, and where exposed and

mined near Redmond, appears as near-vertical, much-contorted beds interleaved with reddish-brown calcareous mudstone. It is likely that several salt beds are in the Redmond area, but the Arapian Shale strata have been so contorted that correlation of beds from one salt mine to the next is uncertain. At Redmond the main salt bed being mined is about 60 m (200 ft) thick (Picard, 1980, p. 145). Pratt, Heylmun, and Cohenour (1966, p. 54) estimate that the salt may range from 180 to 300 m (590 to 980 ft) in thickness. About 9,000 metric tons of salt are mined per year (Pratt and Callaghan, 1970, p. 48), with most of it being used chiefly for livestock and de-icing roads.

Considerable brine is formed in the salt mines near Redmond as a result of ground-water seepage, and drillers have used this brine to drill through the salt beds in the Arapian Shale. The drillers add the brine to their drilling mud, gradually increasing the salinity until the mud becomes supersaturated. At that stage they can drill through a bed of salt without the salt either caving in or dissolving.

The salt is extremely mobile and plastic. Chevron Oil Company was forced to abandon