

INTERIM GEOLOGIC MAP OF THE SUMMIT QUADRANGLE, IRON COUNTY, UTAH

by

Tyler R. Knudsen

Suggested citation:

Knudsen, T.R., 2026, Interim geologic map of the Summit quadrangle, Iron County, Utah: Utah Geological Survey Open-File Report 775DM, 19 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/OFR-775DM>.

Disclaimer

This open-file release makes information available to the public that has undergone only minimal peer review and may not conform to Utah Geological Survey technical, editorial, or policy standards. The map may be incomplete, and inconsistencies, errors, and omissions have not been resolved. The Utah Department of Natural Resources, Utah Geological Survey, makes no warranty, expressed or implied, regarding the suitability of this product for a particular use, and does not guarantee accuracy or completeness of the data. The Utah Department of Natural Resources, Utah Geological Survey, shall not be liable under any circumstances for any direct, indirect, special, incidental, or consequential damages with respect to claims by users of this product. For use at 1:24,000 scale.

This geologic map was funded by the Utah Geological Survey and the U.S. Geological Survey National Cooperative Geologic Mapping Program under STATEMAP award number G21AC10880 (2020–2022). The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.



OPEN-FILE REPORT 775DM
UTAH GEOLOGICAL SURVEY
UTAH DEPARTMENT OF NATURAL RESOURCES
2026

Blank pages are intentional for printing purposes.

SUMMARY

The Summit 7.5' quadrangle is centered on the town of Summit, Iron County, in southwestern Utah. The quadrangle covers the precipitous Hurricane Cliffs along the northwestern margin of the Markagunt Plateau and part of southern Parowan Valley. The northwestern part of the map area covers the southern Red Hills. The quadrangle is bisected from northeast to southwest by Interstate 15. Bedrock in the Summit quadrangle consists of a vertical sequence of sedimentary and volcanic rocks ranging in age from Late Cretaceous to Early Pleistocene. Along much of the Hurricane Cliffs, Cretaceous through Eocene strata dip moderately to steeply northwest as part of the Cedar City-Parowan monocline. The newly described Summit thrust fault places gently north-dipping Cretaceous rocks over west-dipping Cretaceous rocks of the Cedar City-Parowan monocline. Structural relations show that the Summit thrust fault and Cedar City-Parowan monocline are genetically linked and represent the eastward progression of Sevier-age deformation in this area that was active into the Eocene. The modern landscape is dominated by northeast-southwest-trending Basin and Range normal faults that form a series of horsts and grabens. The largest graben, Parowan Valley, is bound by the Parowan fault on the east and the Red Hills fault on the west. The Parowan fault and nearby intrabasin faults in Parowan Valley have locally displaced Late Pleistocene to Holocene alluvial-fan deposits, indicating that the faults should be considered hazardous. This horst-graben system extends onto the Markagunt Plateau, where the Summit Mountain graben preserves Oligocene and Miocene volcanic rocks and Miocene regional-scale gravity-slide deposits derived from the Marysvale volcanic field about 45 miles (72 km) to the north. Extensive Quaternary landslide complexes conceal much of the bedrock near Elliker Basin, Braffits Creek, and Summit Creek.

MAP UNIT DESCRIPTIONS

QUATERNARY

Human-derived deposits

Qh **Artificial fill** (Historical) – Borrow material and engineered fill used to construct flood-control dams, retaining ponds, and other uses; minor fill associated with roadways and building pads generally not mapped; typically less than 20 feet (6 m) thick.

Alluvial Deposits

Qa₁ **Stream alluvium** (Late Holocene) – Stratified, moderately to well-sorted gravel, sand, silt, and clay-size sediment deposited in channels, floodplains, and minor river terraces of Parowan Creek, which no longer receives water/sediment flow in the Summit quadrangle due to diversions; maximum thickness about 20 feet (3 m) thick.

Qay **Young alluvium** (Holocene to Late Pleistocene) – Moderately sorted sand, silt, clay, and pebble to boulder gravel deposited in low-gradient stream channels, floodplains, and adjacent low river terraces; includes small alluvial-fan deposits from adjacent minor drainages and colluvium from adjacent slopes too small to map separately; locally incised by narrow, active stream channels; includes historical debris-flow and debris-flood deposits; probably less than 20 feet (6 m) thick, but deposits in Summit Canyon and Winn Hollow may locally exceed 30 feet (9 m) thick.

Qaf₂ **Level-2 fan alluvium** (Early Holocene to Late Pleistocene) – Poorly to moderately sorted, subangular to subrounded, clay- to boulder-size sediment deposited principally by debris flows and debris floods; forms mostly inactive, moderately incised surfaces cut by younger stream and fan deposits; fault scarps locally prominent on these deposits; equivalent to the older, lower part of younger fan alluvium (**Qafy**); less than 30 feet (9 m) thick.

Qafy **Younger fan alluvium** (Holocene to Late Pleistocene) – Poorly to moderately sorted, non-stratified, subangular to subrounded, boulder- to clay-size sediment deposited at the mouths of streams and washes; forms both active depositional surfaces and low-level mostly inactive surfaces incised by small streams (**Qaf₂** equivalent) that are undivided here; deposited principally as debris flows and debris floods, but colluvium locally constitutes a significant part of the deposits; small isolated deposits are typically less than a few tens of feet thick, but large, coalesced deposits are probably as much as 200 feet (60 m) thick.

Qafc Coalesced fan alluvium (Holocene to Late Pleistocene) – Poorly to moderately sorted, non-stratified, subangular to subrounded, boulder- to clay-size sediment that forms coalesced alluvial fans in Parowan Valley; deposited principally by debris flows and debris floods; similar to younger fan alluvium (**Qafy**) but typically exhibits a lower overall slope and a larger percentage of fine-grained deposits; locally includes small eolian-sand deposits too small to map separately; unconsolidated to poorly consolidated Neogene basin-fill deposits are as much as 2000 feet (610 m) thick in southern Parowan Valley (Hurlow, 2002); only the uppermost part of this basin fill is included in map unit **Qafc**; likely less than 200 feet (60 m) thick.

Qafo Older alluvium (Early? to Late Pleistocene) – Poorly to moderately sorted, non-stratified, subangular to subrounded, clay- to boulder-size sediment with moderately developed pedogenic carbonate; forms broad, gently sloping, deeply dissected surfaces; deposited principally as debris flows and debris floods; exposed thickness is generally less than 40 feet (12 m), but **Qafo** deposits on the floor of Parowan Valley may be more than 100 feet (30 m) thick.

Colluvial Deposits

Qc Colluvium (Holocene to Late? Pleistocene) – Poorly to moderately sorted, angular to subrounded, clay- to boulder-size, locally derived sediment deposited principally by slope wash and soil creep on moderate slopes and in shallow depressions; locally includes talus and alluvial deposits too small to map separately; typically less than 20 feet (6 m) thick.

Qco Older colluvium (Late? Pleistocene) – Poorly to moderately sorted, angular, clay- to boulder-size, locally derived sediment deposited principally by slope wash and soil creep; forms gently sloped, moderately resistant caps on Claron Formation exposed in upper Red and Dalley Canyons; locally includes talus and alluvial deposits too small to map separately; typically less than 20 feet (6 m) thick.

Mass-movement deposits

The Summit quadrangle contains extensive mass-wasting deposits that span in age from older Miocene megabreccia deposits (mapped and described separately) of the Marysvale gravity slide complex (Biek et al., 2019, 2020, 2022) to younger, historically active landslides in West Fork Braffits Creek. The youngest landslide deposits (**Qms**) are locally derived from adjacent slopes and exhibit youthful landslide features such as hummocky topography, main and internal scarps, and lobate morphologies indicating a Late Pleistocene age or younger. Older Quaternary landslides (**Qmso**) are mapped where deposits are perched high above modern drainages, and/or where source areas have been removed by erosion, indicating a Middle Pleistocene or older age. Quaternary landslide deposits are generally unfaulted by underlying high-angle normal faults, although the Parowan fault appears to displace **Qmso** deposits near Elliker Basin.

Qms Landslide deposits (Historical to Late? Pleistocene) – Very poorly sorted, locally derived material deposited principally by rotational and translational movement; locally includes flow-style deposits with lobate morphology; composed of clay- to boulder-size debris as well as large, partly intact bedrock blocks; composition and color depends on local source; characterized by hummocky topography, internal scarps, chaotic bedding attitudes, small ponds, marshy depressions, and meadows; where possible, monolithologic and polyolithologic landslide debris composed of recognized map units are mapped separately, with the bedrock units identified in parentheses within the label—for example, **Qms(Qbe)**; parts of the landslide complex filling the West Fork Braffits Creek drainage are historically active; age and stability determinations require detailed geotechnical investigations; thickness highly variable, some larger landslide complexes are likely at least 200 to 300 feet (60–90 m) thick.

Qms(Qbe), Qms?(Qbe)

Landslide deposits of the Elliker Basin lava flow (Holocene to Late? Pleistocene) – Back-rotated slump blocks of Elliker Basin lava flow southeast of Elliker Basin; queried where a mostly coherent, moderately northwest-dipping sheet of Elliker Basin lava flow may be in place on the footwall of the Parowan fault; see **Qms** unit description for more information; several tens of feet thick.

Qmso Older landslide deposits (Late? to Early? Pleistocene) – Very poorly sorted, locally derived material deposited principally by rotational and translational movement; locally includes flow-style deposits; composed of clay- to boulder-size debris as well as large, partly intact bedrock blocks; deposits are deeply dissected by modern drainages and source areas have locally been removed by erosion, indicating a likely Pleistocene age; composition and color depends on

local source; characterized by hummocky topography and chaotic bedding attitudes, generally lack youthful landslide features such as scarps and lobate morphology; even landslides that have subdued morphology (suggesting that they are older, weathered, and have not experienced recent, large-scale movement) may continue to exhibit slow creep or are capable of renewed movement if stability thresholds are exceeded (Ashland, 2003); age and stability determinations require detailed geotechnical investigations; where possible, monolithologic landslide debris composed of a recognized map unit is mapped separately, with the bedrock unit identified in parentheses within the label—for example, Qmso(Ti); many Qmso deposits mapped in the Summit quadrangle may be equivalent to older Quaternary-Tertiary landslide deposits (QTms) mapped in the adjacent Parowan quadrangle (Knudsen, 2024a), but are not differentiated here due to low confidence in that correlation; thickness highly variable, but locally exceeds several hundred feet.

The hummocky topography and the apparent complexly folded and faulted volcanic units that characterize the Elliker Basin area have previously been explained by the effects of Neogene monoclinial folding (e.g., Anderson and Bucknam, 1979; Anderson and Christenson, 1989, p. 10) or are the result of faulted and juxtaposed megabreccia and lower-plate autochthonous volcanic rocks (Maldonado et al., 1997, plate 1; Biek et al., 2015). Interpretation of lidar (Utah Geospatial Resource Center, 2020) shows hummocks, closed basins, scarps, and lobate topography that are typical of landslide complexes. The unusual topography and geological complexity at Elliker Basin are here interpreted as the result of the Early Pleistocene deposition of landslide blocks derived from the ancestral Hurricane Cliffs. The 0.9 Ma Elliker Basin lava flow (Qmso[Qbe]) is present within the landslide complex, although it is less deformed than underlying landslide blocks. Therefore, most of the landsliding at Elliker Basin occurred prior to the emplacement of the Elliker Basin lava flow, with only minor remobilization of the deposits and overlying lava flow since 0.9 Ma. The source area for landslide deposits was likely a now-eroded series of cliffs located about where lower West Fork Braffits Creek is today. North-northeast of Elliker Basin, distal blocks of Cretaceous strata (Qmso[Ksj]) have no plausible source in the modern landscape, indicating that the current Hurricane Cliffs at this latitude are significantly more subdued today than they were in the Early Pleistocene. The Elliker Basin landslide complex largely conceals the Parowan fault, although subtle scarps aligned with the inferred trace of the fault indicate that the landslide deposits are likely displaced by recurrent movement on the fault.

The large landslide complexes preserved in the Summit Mountain graben contain several recognizable blocks of the Isom and Leach Canyon Formations, and Bauers Tuff. Brian Head strata are dispersed throughout many of the deposits and likely hosted detachment surfaces in the landslides' source areas. Some monolithic landslide blocks in the Summit Mountain graben are likely remobilized gravity-slide deposits, although differentiating between slump blocks derived from gravity-slide deposits versus slump blocks derived from autochthonous Tertiary volcanic rocks was not always possible.

Qmso(Qbe), Qmso(Qbe?)

Older landslide deposits of Elliker Basin lava flow (Late? to Early? Pleistocene) – Mapped where the Elliker Basin lava flow has slumped and been transported as large blocks within the Elliker Basin landslide complex; queried north-northeast of Elliker Basin where large distal slump blocks of lava flow are tentatively correlated with the Elliker Basin lava flow but could also be correlative with the Summit lava flow; see Qmso unit description for more information; several tens of feet thick.

Qmso(Tqh)

Older landslide deposits of Harmony Hills Tuff (Late? to Early? Pleistocene) – Discontinuous, sheared blocks of Harmony Hills Tuff mapped as part of the Elliker Basin landslide complex; see Qmso unit description for more information; less than 50 feet (15 m) exposed.

Qmso(Tqcb)

Older landslide deposits of Bauers Tuff Member of the Condor Formation (Late? to Early? Pleistocene) – Small, structurally isolated slivers of Bauers Tuff Member mapped as part of the Elliker Basin landslide complex; includes some larger partly intact slump blocks mapped in the Summit Mountain graben; see Qmso unit description for more information; thickness highly variable, but deposits in the Summit Mountain graben likely exceed 150 feet (45 m).

Qmso(Tql)

Older landslide deposits of Leach Canyon Formation (Late? to Early? Pleistocene) – Structurally isolated slivers of Leach Canyon Formation mapped as part of the Elliker Basin landslide complex; includes larger, partly intact slump

blocks mapped in the Summit Mountain graben; see **Qmso** unit description for more information; thickness highly variable, but deposits in the Summit Mountain graben likely exceed 200 feet (60 m).

Qmso(Ti)

Older landslide deposits of Isom Formation (Late? to Early? Pleistocene) – Structurally isolated slivers of Isom Formation mapped as part of the Elliker Basin landslide complex; includes larger, partly intact slump blocks mapped in the Summit Mountain graben; see **Qmso** unit description for more information; thickness highly variable, but deposits in the Summit Mountain graben likely exceed 200 feet (60 m).

Qmso(Tcwl)

Older landslide deposits of lower white member of the Claron Formation (Late? to Early? Pleistocene) – Single rotated block of white micritic limestone exposed within an unnamed drainage between Red and Dalley Canyons near the base of the Hurricane Cliffs; see **Qmso** unit description for more information; about 80 feet (25 m) exposed.

Qmso(Tcp)

Older landslide deposits of pink member of the Claron Formation (Late? to Early? Pleistocene) – Mapped where highly sheared slivers of pink member are interspersed within older landslide deposits of Isom Formation between Dalley and Red Canyons near the base of the Hurricane Cliffs; see **Qmso** unit description for more information; less than 150 feet (45 m) exposed.

Qmso(Ksj)

Older landslide deposits of John Henry Member of the Straight Cliffs Formation (Late? to Early? Pleistocene) – Back-rotated slump blocks of John Henry Member mapped at the distal northern edge of the Elliker Basin landslide complex; see **Qmso** unit description for more information; less than 100 feet (30 m) exposed.

Qmt Talus (Holocene to Late Pleistocene) – Poorly sorted, angular cobbles and boulders and finer-grained interstitial sediment deposited principally by rockfall on or at the base of steep slopes and cliffs; talus is common at the base of steep slopes across the map area, but is only mapped where it conceals contacts or forms broad aprons below cliffs of resistant bedrock units; typically grades downslope into colluvium and combined where impractical to differentiate the two; may also include alluvium in the bottom of washes where too small to map separately; typically less than 30 feet (9 m) thick.

Mixed-environment deposits

Qac Alluvium and colluvium (Holocene to Late Pleistocene?) – Poorly to moderately sorted, generally poorly stratified, clay- to boulder-size, locally derived sediment deposited in swales and small drainages by fluvial, slope-wash, and creep processes; gradational with alluvial and colluvial deposits; generally less than 20 feet (6 m) thick.

Qacf Alluvium, colluvium, and fan alluvium (Holocene to Late Pleistocene) – Poorly to moderately sorted, non-stratified, clay- to boulder-size sediment deposited principally by debris flows, debris floods, and slope wash at the mouths of active drainages and the base of steep slopes; locally reworked by small, ephemeral streams; forms coalesced apron of fan alluvium and colluvium impractical to map separately; typically 10 to 40 feet (3–12 m) thick.

Qae Alluvium and eolian sand (Holocene to Late Pleistocene) – Moderately to well-sorted, mostly light-reddish-brown silt and sand deposited by sheetwash and ephemeral streams and locally reworked by eolian processes on the Red Hills lava flow; probably less than 15 feet (5 m) thick.

Qea Eolian sand and alluvium (Holocene to Late Pleistocene) – Moderately to well-sorted, light-brown, fine- to medium-grained eolian sand locally reworked by alluvial processes in Parowan Valley; locally includes minor gravel, sand, and silt deposited in small stream channels; generally less than 20 feet (6 m) thick.

Qmtc Talus and colluvium (Holocene to Late Pleistocene) – Poorly sorted, angular to subangular, cobble- to boulder-size and finer-grained interstitial sediment deposited principally by rockfall and slope wash on steep slopes; includes minor alluvial sediment at the bottom of washes and locally contains small landslides; generally less than 30 feet (9 m) thick.

Qmsc Landslides and colluvium (Holocene to Late Pleistocene) – Unsorted, locally derived, clay- to boulder-size material deposited by slope wash and shallow landslides; mapped where landslide deposits are difficult to identify and inter-mixed with colluvium; most deposits are less than 40 feet (12 m) thick.

Basaltic lava flows

Qbs, Qbsc

Summit lava low and cinder cone (Early Pleistocene) – Medium- to dark-gray, fine-grained olivine basalt ([Table 1](#)) that Maldonado et al. (1997) referred to as the Cinder Hill cone and flow; lava flow (**Qbs**) erupted from a vent at a cinder cone (**Qbsc**) at the base of the Hurricane Cliffs, about 2 miles (3 km) southwest of Summit; the cinder cone has been displaced downward and folded by west-dipping normal faults associated with the Parowan fault so that the cone now sits at a lower elevation than much of the flow exposed higher in the Hurricane Cliffs; **Qbs** also crops out at the southeast margin of the Red Hills near Braffits Creek (Nealey et al., 1997) and is presumed to underlie the southern part of the Parowan Valley graben, where it is displaced by graben-bounding faults; yielded K-Ar ages of 1.00 ± 0.16 Ma and 0.94 ± 0.14 Ma (Anderson and Mehnert, 1979; Table 2); lava flow is as much as 60 feet (18 m) thick above East Fork Braffits Creek.

Qbe Elliker Basin lava low (Early Pleistocene) – Medium- to dark-gray, fine-grained olivine basaltic trachyandesite (shoshonite; [Table 1](#)) mapped near the southern end of the Red Hills and near Elliker Basin; lava flow remnants at Elliker Basin are not continuous and commonly rotated to disparate attitudes and are here interpreted to be part of the Elliker Basin landslide complex; vent area is uncertain, but outcrop patterns and discovery of basaltic dikes in West Fork Braffits Creek indicate the vent may have been centered there; yielded K-Ar ages of 1.00 ± 0.16 Ma and 1.11 ± 0.11 Ma (Anderson and Mehnert, 1979); lava flow is generally less than 20 feet (6 m) thick.

Qid Basaltic dikes (Early Pleistocene?) – Medium-gray, fine-grained olivine basalt dikes exposed in the north canyon wall of West Fork Braffits Creek; appears to be of similar composition to nearby Summit and Elliker Basin lava flows, but based on location, is here interpreted to be feeder dikes to the Elliker Basin vent area that has since been removed by erosion and landslides; dikes intruded into Grand Castle Formation and the capping sandstone member of the Wahweap Formation; most dikes trend northeast and are subvertical, although the northernmost exposed dike dips 48° to the west, concordant with Wahweap bedding; thickness ranges from 3 to 10 feet (1–3 m).

Qbrh, Qbrhc

Red Hills lava low and cinder cone (Early Pleistocene) – Medium- to dark-gray, fine-grained basaltic andesite ([Table 1](#)) with small olivine and plagioclase phenocrysts; lava flows (**Qbrh**) erupted from vents at three cinder cones (**Qbrhc**) along the central axis of the Enoch graben in the Red Hills—two in the Summit quadrangle and one in the Enoch quadrangle (Rowley and Threet, 1976; Knudsen, 2014; Biek et al., 2015); lava flow partially fills the Enoch graben and is cut by graben-bounding faults; yielded K-Ar ages of 1.28 ± 0.4 Ma (Anderson and Mehnert, 1979) and 1.30 ± 0.4 Ma (Best et al., 1980); lava flow is typically about 25 feet (8 m) thick.

unconformity

QUATERNARY-TERTIARY

QTaf Basin-fill deposits (Pleistocene? to Late Miocene?) – Poorly to moderately sorted, non-stratified, subangular to sub-rounded, clay- to boulder-size sediment with moderately to well-developed calcic soils (caliche); forms deeply dissected surfaces with no remaining fan morphology; deposited principally as debris flows; widely exposed in the southern Red Hills where these deposits are stranded on exhumed, fault-bounded horst blocks; presumed to underlie much of Parowan Valley; prominent clasts include Tertiary volcanic rocks, pale-reddish-orange and light-pinkish-gray limestone and calcareous mudstone of the Claron Formation, and yellowish-brown Cretaceous siltstone and sandstone; includes lesser amounts of chalcedony and recycled quartzite from the Grand Castle Formation, Drip Tank Member of the Straight Cliffs Formation, and Claron Formation; locally overlain by the ~1 Ma (Anderson and Mehnert, 1979) Summit lava flow; texture and clast lithology is similar to **QTaf** deposits mapped in the Cedar City quadrangle (Knudsen, 2024b), where the unit is interpreted as basin-fill deposits shed off of the ancestral Hurricane Cliffs (Averitt, 1962, 1964; Knudsen, 2024b); maximum exposed thickness in map area is about 200 feet (60 m).

unconformity

TERTIARY

Marysvale gravity slide complex (Early Miocene to Late Oligocene) – Gravitational collapse of the south flank of the Oligocene to Miocene Marysvale volcanic field produced three gigantic catastrophic gravity slides—the Sevier (SGS), Markagunt (MGS), and Black Mountains (BGS)—that form an overlapping contiguous complex covering an area $>2800 \text{ mi}^2$ ($>7200 \text{ km}^2$) (Biek et al., 2019, 2020, 2022). Each mega-slide travelled southward away from the volcanic edifice with runout over the former land surface of at least 20 miles (32 km). Features such as basal cataclastic breccias and shears, clastic dikes (injectites), jigsaw puzzle fracturing of clasts, and rare pseudotachylyte (frictionite) indicate high-velocity movement aided by over-pressured fluids (Biek et al., 2019, 2022; Braunagel et al., 2023). The principal zone of failure was in mechanically weak, clay-rich, Brian Head Formation sedimentary strata at the base of the volcanic section.

The gravity slide masses become younger westward, possibly following westward migration of volcanism in the field. SGS emplacement was at about 25 Ma, MGS emplacement was at about 23 Ma, and BGS emplacement was at about 21 Ma (Biek et al., 2019; Holliday et al., 2022; Mayback et al., 2022; Stevens et al., 2023; Rivera et al., 2025).

Biek et al. (2019, 2022) provided a history of discovery and current understanding of the gravity slides along with a guide to locations of particularly instructive exposures wherein they drew their conclusions about size, distinctive structural features, emplacement ages, and interpreted emplacement mechanisms.

The BGS is the most recently discovered part of the Marysvale gravity slide complex and is currently the most poorly understood (Rowley et al., in review). The Summit quadrangle lies within the southernmost extent of the BGS as currently understood (Biek et al., 2022), and I tentatively assign gravity-slide (megabreccia) deposits in the Summit quadrangle to the BGS. Components of the Black Mountains Megabreccia (Tmb) are listed below; for their descriptions, see the named pre-existing (undeformed) rock unit elsewhere in the text and in the correlation chart.

Tmb Black Mountains Megabreccia deposits, undivided (Early Miocene) – Chaotic mix of mostly Isom Formation blocks interspersed with sheared blocks of Leach Canyon Formation and calcedony-rich Brian Head Formation with lesser amounts of Bauers Tuff Member; deposits rest on a seemingly autochthonous Tertiary volcanic section well-exposed in the Summit Narrows that includes rocks as young as the ~22 Ma Harmony Hills Tuff; as much as 400 feet (120 m) thick.

Tmb(Tql)

Black Mountains Megabreccia deposits of Leach Canyon Formation (Early Miocene) – Mapped where monolithic piles of intensely brecciated Leach Canyon Formation are present in the Summit Mountain graben, along a ridge east of Elliker Basin, and north of Winn Hollow; typically rests on similarly brecciated Isom Formation, but appears to rest on autochthonous Leach Canyon Formation near Elliker Basin; as much as 400 feet (120 m) thick in the Summit Mountain graben.

Tmb(Ti)

Black Mountains Megabreccia deposits of Isom Formation (Early Miocene) – Mapped where monolithic piles of intensely brecciated Isom Formation are present in the Summit Mountain graben, along a ridge east of Elliker Basin, and at Winn Hollow; typically rests on highly sheared Brian Head Formation or highly brecciated Leach Canyon Formation, and is locally repeated with intervening Leach Canyon breccia; maximum exposed thickness is about 450 feet (135 m) in Dalley Canyon.

Tmb(Tn)

Black Mountains Megabreccia deposits of Needles Range Group, undivided (Early Miocene) – Pale-red to grayish-orange-pink, moderately welded, crystal-rich, poorly exposed, dacitic ash-flow tuff (Mackin, 1960) that crops out between underlying Brian Head Formation and overlying Isom Formation near Winn Gap, but is here interpreted to be part of the Black Mountains Megabreccia; phenocrysts of plagioclase, hornblende, and biotite (plus minor quartz, Fe-Ti oxides, and sanidine) constitute about 40% of the rock; derived from the 27 to 32 Ma Indian Peak caldera complex

that straddles the Utah-Nevada border (Best et al., 1989a, 1989b, 2013); Lund Formation unconformably overlies the similar Wah Wah Springs Formation, but is undivided here due to structural complications; today, the Wah Wah Springs covers at least 8500 square miles (22,000 km²) with an estimated volume of as much as about 720 cubic miles (3000 km³), making it one of the world's most voluminous ash-flow tuffs (Best et al., 1989a); the Lund Formation is of similar volume; Lund Formation is about 27.9 Ma and the Wah Wah Springs Formation is about 29.5 Ma on the basis of many K-Ar and ⁴⁰Ar/³⁹Ar age determinations (Best and Grant, 1987; Best et al., 1989a, 1989b; Rowley et al., 1994a; Best et al., 2013); about 100 to 300 feet (30–90 m) thick in the southern Black Mountains (Rowley, 1976), but only about 25 feet (8 m) of structurally thinned Tn is preserved in the Summit quadrangle.

Tmb(Tbh)

Black Mountains Megabreccia deposits of Brian Head Formation (Early Miocene) – Slivers of sheared white to light-gray sandstone and mudstone and multi-hued chalcedony of the Brian Head Formation mapped in Dalley Canyon and Winn Hollow; underlies gravity-slide deposits of Isom Formation and Needles Range Group; maximum exposed thickness is about 100 feet (30 m) in Dalley Canyon.

structural unconformity

Quichapa Group (Early Miocene to Late Oligocene) – Consists of three regionally distinctive ash-flow tuffs: in ascending order, the Leach Canyon Formation, Condor Canyon Formation, and Harmony Hills Tuff (Mackin, 1960; Williams, 1967; Anderson and Rowley, 1975; Rowley et al., 1995).

Tqh Harmony Hills Tuff (Early Miocene) – Pale-pink to grayish-orange-pink, crystal-rich, moderately welded dacitic ash-flow tuff; contains about 50% phenocrysts that include plagioclase (~60%), biotite (~15%), hornblende (~10%), quartz (~10%), pyroxene (~5%), and sanidine (trace) (Williams, 1967); weathers to rounded outcrops and glittery, sandy soils; the only in-place exposure of Harmony Hills Tuff in the map area is in the northwest corner of section 8, T. 35 S., R. 9 W. in Summit Canyon where it overlies the Bauers Tuff Member and underlies deposits of the Black Mountains Megabreccia (Tmb); rotated blocks of Tqh elsewhere in the map area are interpreted to be Quaternary landslide blocks; source of the Harmony Hills Tuff is unknown, but isopach thickness is greatest near Bull Valley (Williams, 1967), suggesting that it was derived from the eastern Bull Valley Mountains, probably from an early, much more voluminous eruptive phase of the Bull Valley/Hardscrabble Hollow/Big Mountain intrusive arch, as suggested by Blank (1959), Williams (1967), and Rowley et al. (1995, 2006); a preliminary ⁴⁰Ar/³⁹Ar age of 22.023 ± 0.093 Ma (hornblende) and 22.14 ± 0.03 Ma (biotite) for the unit in Parowan Canyon (written communication from Tiffany Rivera, University of Missouri, as reported by Rowley et al. [in review]) agrees well with an ⁴⁰Ar/³⁹Ar plateau age of 22.03 ± 0.15 Ma by Cornell et al. (2001); less than 40 feet (12 m) is exposed in Summit Canyon.

Tqcb Bauers Tuff Member of Condor Canyon Formation (Early Miocene) – Resistant, crystal-poor, light-brownish-gray to pinkish-gray, densely welded, rhyodacitic ash-flow tuff; contains about 10% to 20% phenocrysts of plagioclase (40%–70%), sanidine (25%–50%), biotite (2%–10%), Fe-Ti oxides (1%–8%), and pyroxene (<3%), but lacks quartz phenocrysts (Rowley et al., 1995); bronze-colored biotite present in the upper part of the unit, light-gray flattened lenticles are conspicuous throughout the unit, and a basal vitrophyre 10 to 20 feet (3–6 m) thick is normally present; overlies the Leach Canyon Formation between Summit and Dalley Canyons, and appears as rotated Quaternary landslide blocks elsewhere in the map area; derived from the northwest part (Clover Creek caldera) of the Caliente caldera complex; at the time of its eruption, covered an area of at least 8900 square miles (23,000 km²) (Best et al., 1989b; Rowley et al., 1995) with an estimated volume of 740 mi³ (3200 km³) (Best et al., 2013); the preferred ⁴⁰Ar/³⁹Ar age of the Bauers Tuff Member is 22.7 Ma (Best et al., 1989a) or 22.8 Ma (Rowley et al., 1995), which is the ⁴⁰Ar/³⁹Ar age of its intracaldera intrusion exposed just north of Caliente, Nevada (Rowley et al., 1994b); about 150 feet (45 m) thick in Summit Canyon but thins to about 50 feet (15 m) to the north in Parowan Canyon (Maldonado and Moore, 1995; Biek et al., 2015; Knudsen, 2024a).

Tql Leach Canyon Formation, undivided (Late Oligocene) – Light-pinkish- to orangish-gray, poorly to moderately welded, crystal-rich rhyolite ash-flow tuff that contains abundant white to pale-yellow pumice fragments and lithic clasts, many of which are reddish-brown; contains as much as 25% to 35% total phenocrysts of plagioclase and sub-equal amounts of quartz and sanidine, with minor biotite, hornblende, Fe-Ti oxides, and a trace of pyroxene; disconformably overlies the Isom Formation and is locally overlain by Black Mountains Megabreccia; source is uncertain, but it is probably the Caliente caldera complex because isopach thickness increases toward the complex spanning the Utah-Nevada border (Williams, 1967; Rowley et al., 1995); total volume of the Leach Canyon is estimated to be 830

cubic miles (3600 km³), representing the largest eruption of the Caliente caldera complex (Best et al., 2013); widely agreed to be about 23.8 Ma (Best et al., 1993; Rowley et al., 1995; Biek et al., 2015); as much as 250 feet (75 m) thick in Summit Canyon.

Tda Mount Dutton Formation, alluvial facies (Early Miocene to Early Oligocene) – Resistant to non-resistant, brown, tan, pink, and gray, volcanic mudflow breccia containing matrix-supported angular clasts; mudflow clasts are mostly aphanitic and andesitic rock with sparse small phenocrysts of mostly pyroxene and plagioclase; includes lesser interbedded volcanoclastic conglomerate and tuffaceous sandstone, lava flows, and flow breccia; well exposed in a single mapped outcrop southeast of Second Mound where it rests on Isom Formation; additional Tda rubble—too small to map separately—was found worked into Qms, Qmso, and Tmb deposits at scattered locations in the Summit Mountain graben; deposited from clustered stratovolcanoes that formed most of the southern Marysville volcanic field (e.g., Callaghan, 1938; Anderson and Rowley, 1975; Rowley et al., 1979, 1998, 2002; Steven et al., 1979, 1990; Cunningham et al., 1983; Campbell et al., 1999); K-Ar dated at 21 to 26 Ma (Fleck et al., 1975), but some deposits predate the Wah Wah Springs Formation and therefore are 30 Ma or older; the most voluminous unit in the Marysville volcanic field; thickness varies from 0 to at least 200 feet (60 m).

unconformity

Ti Isom Formation, undivided (Late Oligocene) – Resistant, medium-gray to reddish-brown, crystal-poor, densely welded, trachydacitic ash-flow tuff, typically having distinctive rheomorphic features including flow folds, elongated vesicles, and flow breccias and thus informally described as a “tufflava” (Mackin, 1960; Cook, 1965, 1966; Anderson and Rowley, 1975, 2002); small (1–3 mm) phenocrysts constitute 10% to 15% or less of the rock and are mostly plagioclase (90%) and minor pyroxene and Fe-Ti oxides in a devitrified-glass groundmass; exhibits pronounced subhorizontal lamination or platiness, which Mackin (1960) called “lenticules”; most outcrops have a glass-like fracture habit and weather to grussy, ledgy slopes; the Isom Formation is about 26 to 27 Ma on the basis of many ⁴⁰Ar/³⁹Ar and K-Ar ages (Best et al., 1989b; Rowley et al., 1994a), and because it is locally interbedded with the ~26 Ma Buckskin Breccia (Anderson et al., 1987); Rivera et al. (in review) reported a U-Pb age on zircon of 26.26 ± 0.24 Ma for the Isom Formation in Summit Canyon; likely derived from the Indian Peak caldera complex at the Utah-Nevada border (Best et al., 1989a, 1989b); base not exposed, but is as much as 300 feet (90 m) thick in Summit Canyon; Biek et al. (2015) reported a thickness of about 350 feet (110 m) at Black Ledge about 8 miles (13 km) east of the Summit quadrangle.

unconformity

Brian Head Formation (Early Oligocene to Late Eocene) – The Brian Head Formation is the oldest widespread Tertiary volcanoclastic unit in the southern part of the Marysville volcanic field. See Sable and Maldonado (1997) and Biek et al. (2015) for a summary of the long and complicated nomenclatural history of the Brian Head Formation. The Brian Head Formation was deposited in low-relief fluvial, floodplain, and lacustrine environments in which large amounts of volcanic ash accumulated (Sable and Maldonado, 1997). Northern exposures include coarser deposits on the distal flanks of early stratovolcanoes of the Marysville volcanic field. The Brian Head Formation is 30 to 37 Ma, based on numerous isotopic and fossil ages (e.g., Fleck et al., 1975; Sable and Maldonado, 1997; Eaton et al., 1999a; Korth and Eaton, 2004; Biek et al., 2015; Malone et al., 2025).

Tbh Brian Head Formation, middle volcanoclastic unit – White, light-gray, and pale-yellow sandstone and limestone, with lesser amounts of varicolored mudstone, conglomerate, and multi-hued chalcedony; conglomerate clasts are quartzite, limestone, and chert with minor clasts of intermediate-composition volcanic rocks; chalcedony is various shades of white, gray, yellow, red, black, and brown, typically has a white weathering rind, is commonly highly brecciated and re-silicified, typically occurs in beds 1 to 3 feet (0.3–1 m) thick, is locally stained by manganese oxides, and likely produced by silicification of limestone beds (Maldonado, 1995; Sable and Maldonado, 1997; Schinkel, 2012); abundant bentonitic clay derived from weathered volcanic ash, weathers to strongly swelling soils; unit forms large landslide complexes; typically nonresistant, poorly exposed, and extensively covered by colluvium derived from overlying volcanic units; deposited in low-relief fluvial, floodplain, and lacustrine environments in which large amounts of volcanic ash accumulated (Sable and Maldonado, 1997); about 500 feet (150 m) thick at its type section on Brian Head Peak in the adjoining Brian Head quadrangle (Sable and Maldonado, 1997; Rowley et al., 2013; Biek et al., 2015) and probably of similar thickness in the Summit quadrangle, although less than 150 feet (45 m) is exposed.

unconformity

Claron Formation (Middle Eocene to Paleocene?) – Claron Formation strata are among the most visually striking rocks in the western U.S. and are prominently displayed at Cedar Breaks National Monument and Bryce Canyon National Park, among other places. In the Red Hills and Markagunt Plateau, the Claron Formation is divided into two informal members—an upper white member (which is itself divided into an uppermost mudstone interval, an upper limestone interval, a middle mudstone and sandstone interval, and a lower limestone interval) and the lower pink member (Biek et al., 2015). Claron strata were deposited in fluvial, floodplain, and lacustrine environments of an intermontane basin bounded by Laramide and Sevier uplifts; the pink member is almost entirely fluvial, and the white member is both lacustrine and fluvial (Goldstrand, 1990, 1991, 1992, 1994; Bown et al., 1997). Much of the pink member, and clastic parts of the white member, were greatly modified by bioturbation and pedogenic processes, creating a stacked series of paleosols (Mullett et al., 1988a, 1988b; Mullett, 1989; Mullett and Wells, 1990).

The age of the white member is well constrained as late Middle Eocene (Duchesnean Land Mammal Age), based on limiting U-Pb zircon ages for overlying Brian Head Formation (Rowley et al., 2013; Malone et al., 2025) and conglomerate at Boat Mesa (Biek et al., 2015; Malone et al., 2025), and by Late Eocene mammals and ostracods from what we now know to be part of basal Brian Head strata on the Markagunt Plateau (where it was originally misidentified as uppermost Claron Formation [Eaton et al., 2011, 2018]). Along Sweetwater Creek north of Bryce Canyon National Park, Eaton et al. (2018) recovered the Early Eocene (Wasatchian North American Land Mammal Age) rodents *Knightomys reginensis* and *K. minor* from their unit 7, roughly 900 feet (275 m) above their base of the formation in what may be Tcwm equivalent strata. Charophytes recovered from near the base of the lower pink member at Griffin Top, about 3 miles (5 km) north of Sweetwater Creek, suggest an Early Eocene (Ypresian) age for basal Claron strata (Sanjuan and Eaton, 2016). This suggests that the entire Claron Formation in the Bryce Canyon region is Early Eocene in age. However, throughout the western part of its outcrop belt the maximum age of the mostly nonfossiliferous pink member is poorly constrained as Early Eocene to Paleocene(?) (Goldstrand, 1990, 1994; Sanjuan and Eaton, 2016; Eaton et al., 2018; Biek et al., 2015).

- Tcwt Uppermost mudstone, siltstone, and sandstone unit of white member** (Late and Middle Eocene) – Varicolored and commonly mottled, pale-reddish-orange, reddish-brown, moderate-orange-pink, dark-yellowish-orange, and grayish-pink, calcareous mudstone and siltstone mapped in the southern Red Hills; locally includes minor fine-grained silty sandstone and micritic limestone; indistinguishable in lithology and color from the middle unit of the White Member (Tcwm) of the Claron Formation; forms a brightly colored slope on top of the upper limestone ledge of the white member; about 120 feet (37 m) thick in Winn Hollow.
- Tcwu Upper limestone unit of white member** (Eocene) – White, pale-yellowish-gray, pinkish-gray, and pale-orange micritic limestone, locally containing intraformational rip-up clasts mapped in the southern Red Hills; typically poorly bedded and knobby weathering; locally vuggy with calcite spar and commonly cut by calcite veinlets; resistant and so forms prominent ledges and flat ridge tops; upper conformable contact with Tcwt corresponds to a pronounced color change from white to pale-orange micritic limestone below to brightly colored reddish-orange mudstone and siltstone above; about 25 feet (8 m) thick.
- Tcwm Middle mudstone, siltstone, and sandstone unit of white member** (Eocene) – Varicolored calcareous mudstone and siltstone, and minor fine-grained calcareous sandstone mapped in the southern Red Hills; weathers to a poorly exposed slope; upper conformable contact corresponds to a pronounced color change from brightly colored reddish-orange mudstone and siltstone below to white to very pale-orange micritic limestone above; about 65 feet (20 m) thick.
- Tcwl Lower limestone unit of white member** (Eocene) – White, pale-yellowish-gray, and pale-orange micritic limestone mapped in the southern Red Hills; typically forms a cliff above the pink member (Tcp); upper conformable contact corresponds to a pronounced color change from white to pale-orange micritic limestone below to brightly colored reddish-orange mudstone and siltstone above; about 30 feet (10 m) thick.
- Tcp Pink member** (Eocene to Paleocene?) – Orangish-red, reddish-brown, and light-blue-gray limestone, mudstone, siltstone, sandstone, and conglomerate; locally mottled; oncolitic beds are common; limestone is poorly bedded, microcrystalline, and sandy, with 2% to 20% fine-grained quartz sand; locally argillaceous beds formed as calcic paleosols; fluvial and floodplain deposits greatly modified by bioturbation and pedogenic processes (Mullett et al., 1988a, 1988b; Mullett, 1989; Mullett and Wells, 1990); sandstone is thick-bedded, fine- to coarse-grained, calcareous, locally cross-bedded quartz arenite; mudstone is generally reddish-orange, silty, calcareous, contains calcareous nodules, and weathers to earthy, steep slopes between ledges of sandstone and limestone; pebbly conglomerate forms lenticular beds typically 5 to 15 feet (2–5 m) thick, containing rounded quartzite, limestone, and chert

pebbles and cobbles; upper conformable contact corresponds to a pronounced color change from reddish-orange mudstone and siltstone below to white to very pale orange, resistant, micritic limestone of the white member above; about 1000 feet (300 m) thick.

CRETACEOUS

Kgc Grand Castle Formation (Late Cretaceous?, Maastrichtian? to late Campanian?) – Light-gray and light-red massive conglomerate; clasts are well-rounded, pebble- to boulder-size quartzite, limestone, sandstone, and chert; typically cliff-forming; redefined and restricted by Biek et al. (2015) to only the upper of three informal members (resistant upper conglomerate, nonresistant middle sandstone, and resistant lower conglomerate) of the Grand Castle Formation of Goldstrand and Mullett (1997); deposited in a braided fluvial environment with paleoflow principally to the east and south-southeast, suggesting source areas in the Wah Wah, Blue Mountain, and Iron Springs thrust sheets of southwest Utah (Goldstrand and Mullett, 1997); Biek et al. (2015) recovered late Campanian to Maastrichtian pollen from Grand Castle strata south of Cedar Breaks National Monument; upper contact with reddish-orange Claron strata is poorly exposed on the Markagunt Plateau, but appears to be gradational and largely conformable in the Red Hills; thickness varies from about 150 feet (45 m) on the Markagunt Plateau to as much 350 feet (105 m) in the southern Red Hills.

unconformity

Kwcs, Kwcs?

Capping sandstone member of the Wahweap Formation (Late Cretaceous, middle Campanian) – White to very pale orange, locally iron stained, mostly medium-grained, trough cross-bedded quartz arenite; the sandstone “caps” the lower mudstone-rich members of the Wahweap Formation near its type section (Kaiparowits Basin) (Eaton, 1991) and much of the Markagunt Plateau; lower undivided mudstone members are generally missing in the Summit quadrangle and the capping sandstone member appears to sit conformably on the Drip Tank Member of the Straight Cliffs Formation; limited exposures in West Fork Braffits Creek include several feet of varicolored and mottled mudstone typical of lower undivided members of the Wahweap; named the Pardner Canyon Member by Beveridge et al. (2022) in the Kaiparowits Plateau; upper part of **Kwcs** contains abundant pebble stringers and conglomeratic beds with rounded quartzite, dolomite, chert, and limestone clasts; clasts are typically about 1 inch (2.5 cm) in diameter but as large as 2 to 3 inches (5–7.5 cm), and include common reddish-brown and purple quartzite clasts; quartz grains are typically well rounded and commonly frosted, recycled from Mesozoic eolianites (Pollock, 1999; Lawton et al., 2003; see also UGS and AtoZ, 2013); Goldstrand and Mullett (1997) and Lawton et al. (2003) showed that the member was deposited in a braided fluvial environment with a paleoflow direction principally to the east and south-southeast, suggesting source areas in Navajo Sandstone exposed in the upper plate of the Iron Springs thrust; contains carbonized or petrified plant debris, small mudstone rip-up clasts, iron concretions, and soft-sediment deformation features; typically poorly cemented, forming distinctive white, manzanita-covered slope-and-bench topography; queried in the southern Red Hills where yellow-weathering, easily erodible sandstone appears between underlying upper Iron Springs Formation and overlying Grand Castle strata; unusually iron-rich **Kwcs** concretionary zones in the southern Red Hills, and debris derived from them, appear to be the target of several shallow prospects in the area; Campanian to Santonian palynomorphs and a theropod dinosaur track discovered in **Kwcs** strata in the adjoining Parowan quadrangle confirm a Late Cretaceous age (Hunt et al., 2011; Biek et al., 2015); chronostratigraphic studies (based on new U-Pb zircon ages) by Beveridge et al. (2022) on the Kaiparowits Plateau placed the upper and lower boundaries of the Wahweap at about 77 and 82 Ma, respectively; typically about 200 feet (60 m) thick.

Iron Springs Formation (Late Cretaceous, Santonian or early Campanian to Cenomanian) – Mapped as upper and lower units in the Red Hills. Iron Springs strata were deposited principally in braided-stream and floodplain environments of a coastal plain (Johnson, 1984; Fillmore, 1991; Eaton et al., 2001; Milner et al., 2006) and is typically correlated to the Naturita Formation, Tropic Shale, and Straight Cliffs Formation (Eaton, 1999; Eaton et al., 2001). Late Cretaceous age is from Goldstrand (1994) and an ash that is 712 feet (217 m) below the top of the formation in Parowan Canyon, which yielded an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 83.0 ± 1.1 Ma (Eaton et al., 1999b). Lower Iron Springs strata (Kil) in the upper plate of the Iron Springs thrust may be associated with the maximum transgression of the Greenhorn Sea of late Cenomanian or early Turonian age (Eaton et al., 1997; Eaton, 1999; Biek et al., 2015). The entire formation is about 3500 to 4000 feet (1070–1220 m) thick in the Pine Valley Mountains southwest of the quadrangle (Cook, 1960).

- Kiu Upper unit** – Interbedded, ledge-forming, calcareous, cross-bedded, fine- to medium-grained sandstone and less-resistant, poorly exposed sandstone, siltstone, and mudstone present in the footwall of the Iron Springs thrust; variously colored grayish orange, pale yellowish orange, dark yellowish orange, white, pale reddish brown, and greenish gray; locally stained by iron-manganese oxides; Liesegang banding is common in the sandstone beds; sandstone beds range from quartz arenite to litharenite in composition (Fillmore, 1991; Goldstrand, 1992); weathers to repetitive, thick, tabular sandstone beds and thinner interbedded mudstone; upper contact with the Grand Castle Formation (redefined by Biek et al. [2015]) is locally difficult to map because of abundant Grand Castle-derived colluvium covering downslope units; Milner et al. (2006) reported on dinosaur tracks in upper Iron Springs strata near Parowan Gap, and also noted a diverse assemblage of plant fossils, bivalves, gastropods, turtles, fish, and trace fossils suggestive of late Santonian to early Campanian age; an incomplete section exposed in the Red Hills is about 600 feet (180 m) thick.
- Kil Lower unit** – Interbedded sandstone, siltstone, and mudstone similar to that of the upper unit (Kiu) but restricted to the upper plate of the Iron Springs thrust fault; a single incomplete thrust-fault bounded section that is partially concealed by the Red Hills lava flow and is less than 100 feet (30 m) thick.

Straight Cliffs Formation (Late Cretaceous, early Campanian to Turonian) – Peterson (1969) divided the Straight Cliffs Formation into four members in the Kaiparowits Basin, in descending stratigraphic order: the Drip Tank, John Henry, Smoky Hollow, and Tibbet Canyon Members. Several geologists mapped these members (separately or as lumped upper and lower Straight Cliffs strata) on the Paunsaugunt Plateau, including Tilton (1991, 2001a, 2001b), Doelling and Willis (1999), Sable and Hereford (2004), and Doelling (2008). Biek et al. (2015) described the difficulty encountered in early attempts to carry this nomenclature westward into the Markagunt Plateau. The Straight Cliffs Formation is an overall regressive sequence that formed during the last marine incursion of the Western Interior Seaway (e.g., Eaton et al., 2001; Moore and Straub, 2001; Tibert et al., 2003).

Cretaceous strata exposed along the western margin of the Markagunt Plateau from Fiddlers Canyon, in the adjoining Cedar City quadrangle, to Parowan are warped down to the northwest along the Cedar City-Parowan monocline (Threet, 1963; Anderson and Bucknam, 1979; Anderson and Mehnert, 1979; Anderson and Christenson, 1989). Early workers interpreted the monocline to be related to Basin and Range extension. Mapping by Biek et al. (2015) showed that Late Oligocene to Early Miocene regional ash-flow tuffs are not affected by monoclinal folding, and that the fold is likely related to Sevier-age deformation, a conclusion supported by this map. A newly discovered west-verging thrust fault—here called the Summit thrust fault—that extends from Strawberry Canyon northward to Summit Canyon, duplicates the upper parts of the Straight Cliffs Formation. The fault places gently northeast-dipping strata over a lower plate of steeply northwest-dipping rocks of the Cedar City-Parowan monocline, indicating a genetic link between the thrust faulting and the monocline. The Summit Canyon thrust fault likely formed during the late stages of folding that produced the Cedar City-Parowan monocline. Grand Castle and Claron formations exposed near the mouths of Summit and Red Creek Canyons are involved in monoclinal folding indicating that the fold largely developed after Claron strata were deposited. In contrast, the Iron Springs thrust fault exposed in Parowan Gap 8 miles (13 km) north-northwest of Summit Canyon, largely formed prior to deposition of the Grand Castle and Claron Formations with relatively minor displacements continuing into Claron time (Anderson and Dinter, 2010; Biek et al., 2015). Thus, the Cedar City-Parowan monocline and Summit thrust fault represent the Early Eocene eastward progression of Sevier deformation in this area.

Maldonado and Moore (1995) mapped a single thrust fault along the Cedar City-Parowan monocline near the east-central edge of the Parowan quadrangle that they interpreted to place Cretaceous rocks over the pink member of the Claron Formation. Maldonado et al. (1997) described additional thrust faults associated with the monocline in the Summit quadrangle that involve rocks as young as Early Miocene and named the faults the Parowan thrust fault zone. However, I favor the interpretation of Biek et al. (2015; p. 64), who argue that these features are best explained by high-angle normal faulting and landsliding.

- Ksd Drip Tank Member** (Late Cretaceous, early Campanian) – Massive, typically cliff-forming, white to light-gray, medium-grained sandstone and pebbly conglomerate; larger clasts (pebble to cobble) are subrounded to rounded, white, gray, and purplish-red quartzite, with local minor blue-gray Paleozoic limestone and black chert; sandy matrix consists of white to light-gray, subangular sand and silt; locally iron stained; forms a low, narrow ridge; formerly called the lower conglomerate member of the Grand Castle Formation (Eaton, 1991), but Biek et al. (2015) established the Drip Tank correlation used for this map; unit was deposited by east- and northeast-flowing braided streams (Tilton, 1991, 2001a, 2001b; Lawton et al., 2003); upper contact with the Wahweap Formation appears conformable and corresponds to the top of a white pebbly sandstone, above which is yellowish-brown, fine-grained sandstone of the capping sandstone member of the Wahweap; recent chronostratigraphic studies (based on new U-Pb zircon ages) on the Wahweap

Formation in the Kaiparowits Plateau by Beveridge et al. (2022) placed the lower boundary of the Wahweap with the Drip Tank Member at about 82 Ma, providing a minimum age for Drip Tank strata; thickness ranges widely from 50 to 150 feet (15–45 m).

unconformity

Ksjs John Henry and Smoky Hollow Members, undivided (Late Cretaceous, Santonian to Turonian) – Undivided in the Summit quadrangle, where underlying Smoky Hollow is commonly concealed by colluvium or landslides, and the Calico bed (Peterson, 1969) is either poorly developed or absent (Biek et al., 2015); Smoky Hollow strata are described below and John Henry strata are described separately; combined unit is about 1250 feet (380 m) thick near the southeastern corner of the map area and appears to thin to about 1000 feet (300 m) thick in the structurally complicated Hurricane Cliffs.

Smoky Hollow strata are slope-forming, brown and gray mudstone, shale, and interbedded yellowish-brown fine-grained sandstone; lower part contains a few thin coal beds, common carbonaceous shale, and several thin oyster coquina beds; upper unconformable contact typically corresponds to the base of the Calico bed—a stacked series of fluvial channel deposits of white to light-gray, fine- to medium-grained sandstone and conglomeratic sandstone—that is not recognized in the Summit quadrangle; deposited in fluvial and floodplain environments of a coastal plain (Eaton et al., 2001; Primm et al., 2018); Smoky Hollow strata are middle to late Turonian on the basis of a diverse assemblage of mollusks, benthic foraminifera, and ostracods from exposures in Cedar Canyon (Eaton et al., 2001; Tibert et al., 2003); Szwarc et al. (2015) reported a maximum depositional age for the member of 89.1 ± 6.35 Ma; probably about 250 feet (75 m) thick in the Summit quadrangle.

Ksj John Henry Member (Late Cretaceous, Santonian to late Turonian) – Yellowish- to reddish-brown, fine- to medium-grained, subarkosic sandstone and siltstone, and interbedded, locally mottled, gray, brown, and reddish-brown mudstone; forms ledgy slopes; sandstone is commonly bioturbated and locally stained by iron-manganese oxides; stacked or amalgamated sandstone beds make up most of the upper part of the unit; upper unconformable contact corresponds to a break in slope at the base of the Drip Tank Member; woody material and leaf impressions are locally abundant; Chentnik et al. (2015) identified four regressive-transgressive cycles within the member on the Kaiparowits Plateau and Benhallam et al. (2016) noted that the member provides an ~6-million-year record of coastal-plain to marginal-marine deposition; biotite from an ash bed about 800 feet (245 m) above the base of the member in Cedar Canyon yielded an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 86.72 ± 0.58 Ma (late Coniacian), and biotite from an ash bed 700 feet (213 m) below the top of the member in Parowan Canyon yielded an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 83.0 ± 1.1 Ma (early Campanian to late Santonian) (Eaton et al., 1999b); Szwarc et al. (2015) reported a maximum depositional age for the member of 82.8 ± 4.1 Ma; Eaton (2006) reported on mammal fossils in Cedar Canyon that suggest the lower part of the member there is late Turonian; thickness about 900 to 1000 feet (275–300 m) in the Summit quadrangle.

Kst Tibbet Canyon Member (Late Cretaceous, Turonian) – Yellowish-brown, medium- to thick-bedded, generally planar bedded, fine- to medium-grained quartzose sandstone and interbedded gray mudstone, carbonaceous shale, and thin to thick beds of oyster coquina; forms prominent cliffs along East Fork Braffits Creek canyon; upper conformable contact corresponds to a pronounced break in slope and is placed at the top of a coquina oyster bed and base of overlying thin coal and carbonaceous shale interval that caps the member; deposited during initial progradation of the Greenhorn Cycle in shoreface, beach, lagoonal, and estuarine environments adjacent to a coastal plain (Laurin and Sageman, 2001, 2007; Tibert et al., 2003); Szwarc et al. (2015) reported a maximum depositional age (U-Pb zircon) for the member of 94.3 ± 1.4 Ma; incomplete section is at least 450 feet (135 m) thick in the map area, but is typically 650 to 800 feet (200–245 m) thick on the west flank of the Markagunt Plateau (Biek et al., 2015).

ACKNOWLEDGMENTS

I thank Bob Biek (Utah Geological Survey [UGS], retired) for sharing his considerable knowledge of the local geology. Dave Hacker (Kent State University) provided valuable insight and map notes focused on the Marysvale gravity slide complex. Draft mapping made available to the UGS by Florian Maldonado and Edward Sable (USGS, retired) was a helpful resource. I thank Josh Dustin (UGS) for his excellent GIS and cartography work on the map. Reviews by Bob Biek, Stefan Kirby, Adam McKean, Stephanie Carney, and Mike Hylland (retired) of the UGS, greatly improved this map.

REFERENCES

- Anderson, J.J., Iivari, T.A., and Rowley, P.D., 1987, Geologic map of the Little Creek Peak quadrangle, Garfield and Iron Counties, Utah: Utah Geological and Mineral Survey Map 104, 11 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/M-104>.
- Anderson, J.J., and Rowley, P.D., 1975, Cenozoic stratigraphy of the southwestern High Plateaus of Utah, *in* Anderson, J.J., Rowley, P.D., Fleck, R.J., and Nairn, A.E.M., editors, Cenozoic geology of southwestern High Plateaus of Utah: Geological Society of America Special Paper 160, p. 1–52, <https://doi.org/10.1130/SPE160>.
- Anderson, J.J., and Rowley, P.D., 2002, The Oligocene Isom Formation, Utah and Nevada—a regional ash-flow tuff sheet containing fluidal features of a lava flow [abs.]: Geological Society of America Abstracts with Programs, v. 34, no. 4, p. 9.
- Anderson, L.P., and Dinter, D.A., 2010, Deformation and sedimentation in the southern Sevier Foreland, Red Hills, southwestern Utah, *in* Carney, S.M., Tabet, D.E., and Johnson, C.L., editors, Geology of south-central Utah: Utah Geological Association Publication 39, p. 338–366.
- Anderson, R.E., and Bucknam, R.C., 1979, Two areas of probable Holocene deformation in southwestern Utah: Tectonophysics, v. 52, p. 417–430, [https://doi.org/10.1016/0040-1951\(79\)90257-9](https://doi.org/10.1016/0040-1951(79)90257-9).
- Anderson, R.E., and Christenson, G.E., 1989, Quaternary faults, folds, and selected volcanic features in the Cedar City 1° x 2° quadrangle, Utah: Utah Geological and Mineral Survey Miscellaneous Publication 89-6, 29 p., scale 1:250,000, <https://doi.org/10.34191/MP-89-6>.
- Anderson, R.E., and Mehnert, H.H., 1979, Reinterpretation of the history of the Hurricane fault in Utah, *in* Newman, G.W., and Goode, H.D., editors, Basin and Range symposium: Rocky Mountain Association of Geologists and Utah Geological Association, p. 145–165.
- Ashland, F.X., 2003, Characteristics, causes, and implications of the 1998 Wasatch Front landslides, Utah: Utah Geological Survey Special Study 105, 49 p., <https://doi.org/10.34191/SS-105>.
- Averitt, P., 1962, Geology and coal resources of the Cedar Mountain quadrangle, Iron County, Utah: U.S. Geological Survey Professional Paper 389, 71 p., 3 plates, scale 1:24,000, <https://doi.org/10.3133/pp389>.
- Averitt, P., 1964, Upper Tertiary surficial deposits near Cedar City, Iron County, Utah: Geological Society of America Bulletin, v. 75, p. 37–44, [https://doi.org/10.1130/0016-7606\(1964\)75\[37:UTSDNC\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1964)75[37:UTSDNC]2.0.CO;2).
- Averitt, P., and Threet, R.L., 1973, Geologic map of the Cedar City quadrangle, Iron County, Utah: U.S. Geological Survey Geologic Quadrangle Map GQ-1120, scale 1:24,000, <https://doi.org/10.3133/gq1120>.
- Benhallam, W., Turner, A., Stright, L., and Johnson, C.L., 2016, Spatial analysis of channel-belt stacking patterns—metrics to discriminate between local and regional controls on deposition in the fluvial John Henry Member of the Straight Cliffs Formation, southern Utah, USA: Journal of Sedimentary Research, v. 86, p. 1310–1327, <http://dx.doi.org/10.2110/jsr.2016.77>.
- Best, M.G., Christiansen, E.H., and Blank, R.H., Jr., 1989a, Oligocene caldera complex and calc-alkaline tuffs and lavas of the Indian Peak volcanic field, Nevada and Utah: Geological Society of America Bulletin, v. 101, p. 1076–1090, [https://doi.org/10.1130/0016-7606\(1989\)101%3C1076:OCCACA%3E2.3.CO;2](https://doi.org/10.1130/0016-7606(1989)101%3C1076:OCCACA%3E2.3.CO;2).
- Best, M.G., Christiansen, E.H., Deino, A.L., Grommé, C.S., Hart, G.L., and Tingey, D.G., 2013, The 36–18 Ma Indian Peak-Caliente ignimbrite field and calderas, southeastern Great Basin, USA—multicyclic super-eruptions: Geosphere, v. 9, no. 4, p. 1–87, <https://doi.org/10.1130/GES00902.1>.
- Best, M.G., Christiansen, E.H., Deino, A.L., Gromme, C.S., McKee, E.H., and Noble, D.C., 1989b, Excursion 3A—Eocene through Miocene volcanism in the Great Basin of the western United States: New Mexico Bureau of Mines and Mineral Resources Memoir 47, p. 91–133, <https://doi.org/10.58799/M-47>.
- Best, M.G., and Grant, S.K., 1987, Stratigraphy of the volcanic Oligocene Needles Range Group in southwestern Utah and eastern Nevada: U.S. Geological Survey Professional Paper 1433A, p. 1–28, <https://doi.org/10.3133/pp1433AB>.
- Best, M.G., McKee, E.H., and Damon, P.E., 1980, Space-time-composition patterns of late Cenozoic mafic volcanism, southwestern Utah and adjoining areas: American Journal of Science, v. 280, p. 1035–1050, <https://doi.org/10.2475/ajs.280.10.1035>.
- Best, M.G., Scott, R.B., Rowley, P.D., Swadley, W.C., Anderson, R.E., Gromme, C.S., Harding, A.E., Deino, A.L., Christiansen, E.H., Tingey, D.G., and Sullivan, K.R., 1993, Oligocene-Miocene caldera complexes, ash-flow sheets, and tectonism in the central and southeastern Great Basin, *in* Lahren, M.M., Trexler, J.H., Jr., and Spinoso, C., editors, Crustal evolution of the Great Basin and Sierra Nevada: Field trip guide, Geological Society of America, Cordilleran and Rocky Mountain Sections meeting, p. 285–311.

- Beveridge, T.L., Roberts, E.M., Ramezani, J., Titus, A.L., Eaton, J.G., Irmis, R.B., and Sertich, J.J.W., 2022, Refined geochronology and revised stratigraphic nomenclature of the Upper Cretaceous Wahweap Formation, Utah, U.S.A. and the age of early Campanian vertebrates from southern Laramidia: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 591, <https://doi.org/10.1016/j.palaeo.2022.110876>.
- Biek, R.F., Rowley, P.D., Anderson, J.J., Maldonado, F., Moore, D.W., Hacker, D.B., Eaton, J.G., Hereford, R., Filkorn, H., and Matyjasik, B., 2015, Geologic map of the Panguitch 30' x 60' quadrangle, Garfield, Iron, and Kane Counties, Utah: Utah Geological Survey Map 270DM, 162 p., 3 plates, scale 1:62,500, <https://doi.org/10.34191/M-270dm>.
- Biek, R.F., Rowley, P.D., and Hacker, D.B., 2019, The gigantic Markagunt and Sevier gravity slides resulting from mid-Cenozoic catastrophic mega-scale failure of the Marysvale volcanic field, Utah, USA: *Geological Society of America Field Guide* 56, 121 p., <https://doi.org/10.1130/FLD056>.
- Biek, R.F., Rowley, P.D., and Hacker, D.B., 2020, Utah's ancient mega-landslides: Utah Geological Survey, Survey Notes, v. 52, no. 2, p. 1–3, <https://doi.org/10.34191/SNT-52-2>.
- Biek, R.F., Rowley, P.D., and Hacker, D.B., 2022, Utah's ancient mega-landslides—geology, discovery, and guide to Earth's largest terrestrial landslides: Utah Geological Survey Circular 132, 67 p., <https://doi.org/10.34191/C-132>.
- Blank, H.R., Jr., 1959, Geology of the Bull Valley district, Washington County, Utah: Seattle, University of Washington, Ph.D. dissertation, 177 p.
- Bown, T.M., Hasiotis, S.T., Genise, J.F., Maldonado, F., and Bowers, E.M., 1997, Trace fossils of *Hymenoptera* and other insects, and paleoenvironments of the Claron Formation (Paleocene and Eocene), southwestern Utah, in Maldonado, F., and Nealey, L.D., editors, *Geologic studies in the Basin and Range-Colorado Plateau transition in southeastern Nevada, southwestern Utah, and northwestern Arizona*, 1995: U.S. Geological Survey Bulletin 2153, p. 43–58, <https://doi.org/10.3133/b2153>.
- Braunagel, M.J., Griffith, W.A., Biek, R.F., Hacker, D.B., Rowley, P.D., Malone, D.H., Mayback, D., Rivera, T.A., Loffer, Z., and Smith, Z.D., 2023, Structural relationships across the Sevier gravity slide of southwest Utah and implications for catastrophic translation and emplacement processes of long runout landslides: *Geochemistry, Geophysics, Geosystems*, v. 24, p. 1–25, <https://doi.org/10.1029/2022GC010783>.
- Callaghan, E., 1938, Preliminary report on the alunite deposits of the Marysvale region, Utah: U.S. Geological Survey Bulletin 886-D, p. 91–131, <https://doi.org/10.3133/b886D>.
- Campbell, D.L., Steven, T.A., Cunningham, C.G., and Rowley, P.D., 1999, Aeromagnetic and gravity maps of the central Marysvale volcanic field, southwestern Utah: U.S. Geological Survey Miscellaneous Investigations Series Map I-2645-B, scale 1:100,000, <https://doi.org/10.3133/i2645B>.
- Chentnik, B.M., Johnson, C.L., Mulhern, J.S., and Stright, L., 2015, Valleys, estuaries, and lagoons—paleoenvironments and regressive-transgressive architecture of the Upper Cretaceous Straight Cliffs Formation, Utah, USA: *Journal of Sedimentary Research*, v. 85, p. 1166–1196, <http://dx.doi.org/10.2110/jsr.2015.70>.
- Cook, E.F., 1960, Geologic atlas of Utah, Washington County: Utah Geological and Mineralogical Survey Bulletin 70, 119 p., scale 1:125,000, <https://doi.org/10.34191/B-70>.
- Cook, E.F., 1965, Stratigraphy of Tertiary volcanic rocks in eastern Nevada: Nevada Bureau of Mines Report 11, 61 p.
- Cook, E.F., editor, 1966, Tufflavas and ignimbrites, a survey of Soviet studies: New York, American Elsevier Publishing Company, Inc., 212 p.
- Cornell, D., Butler, T., Holm, D., Hacker, D., and Spell, T., 2001, Stratigraphy and $^{40}\text{Ar}/^{39}\text{Ar}$ ages of volcanic rocks of the Pinto quadrangle, Colorado Plateau transition zone, SW Utah [abs.], in Erskine, M.C., Faulds, J.E., Bartley, J.M., and Rowley, P.D., editors, *The geologic transition, High Plateaus to Great Basin—a symposium and field guide (The Mackin Volume)*: Utah Geological Association and Pacific Section of the American Association of Petroleum Geologists, Utah Geological Association Publication 30, p. 420–421.
- Cunningham, C.G., Steven, T.A., Rowley, P.D., Glassgold, L.B., and Anderson, J.J., 1983, Geologic map of the Tushar Mountains and adjoining areas, Marysvale volcanic field, Utah: U.S. Geological Survey Miscellaneous Investigations Series Map I-1430-A, scale 1:50,000, <https://doi.org/10.3133/i1430A>.
- Doelling, H.H., 2008, Geologic map of the Kanab 30' x 60' quadrangle, Kane and Washington Counties, Utah, and Coconino and Mohave Counties, Arizona: Utah Geological Survey Miscellaneous Publication MP-08-2DM, scale 1:100,000, <https://doi.org/10.34191/MP-08-2dm>.

- Doelling, H.H., and Willis, G.C., 1999, Interim geologic map of the SE part of the Panguitch 30' x 60' quadrangle, Garfield and Kane Counties, Utah: Utah Geological Survey Open-File Report 367, 10 p., 1 plate, scale 1:100,000, <https://doi.org/10.34191/OFR-367>.
- Eaton, J.G., 1991, Biostratigraphic framework for the Upper Cretaceous rocks of the Kaiparowits Plateau, southern Utah, in Nations, J.D., and Eaton, J.G., editors, Stratigraphy, depositional environments, and sedimentary tectonics of the western margin, Cretaceous Western Interior Seaway: Geological Society of America Special Paper 260, p. 47–63, <https://doi.org/10.1130/SPE260-p47>.
- Eaton, J.G., 1999, Vertebrate paleontology of the Iron Springs Formation, Upper Cretaceous, southwestern Utah, in Gillette, D.D., editor, Vertebrate paleontology in Utah: Utah Geological Survey Miscellaneous Publication 99-1, p. 339–343, <https://doi.org/10.34191/MP-99-1>.
- Eaton, J.G., 2006, Late Cretaceous mammals from Cedar Canyon, southwestern Utah, in Lucas, S.G., and Sullivan, R.M., editors, Late Cretaceous vertebrates from the Western Interior: New Mexico Museum of Natural History and Science Bulletin 35, p. 373–402.
- Eaton, J.G., Hutchinson, J.H., Holroyd, P.A., Korth, W.W., and Goldstrand, P.M., 1999a, Vertebrates of the turtle basin local fauna, middle Eocene, Sevier Plateau, south-central Utah, in Gillette, D.D., editor, Vertebrate paleontology in Utah: Utah Geological Survey Miscellaneous Publication 99-1, p. 463–468, <https://doi.org/10.34191/MP-99-1>.
- Eaton, J.G., Kirkland, J.I., Hutchinson, J.H., Denton, R., O'Neill, R.C., and Parrish, J.M., 1997, Nonmarine extinction across the Cenomanian-Turonian boundary, southwestern Utah, with a comparison to the Cretaceous-Tertiary extinction event: Geological Society of America Bulletin, v. 109, p. 560–567, [https://doi.org/10.1130/0016-7606\(1997\)109%3C0560:NEA TCT%3E2.3.CO;2](https://doi.org/10.1130/0016-7606(1997)109%3C0560:NEA TCT%3E2.3.CO;2).
- Eaton, J.G., Korth, W.W., and Brinkman, D.B., 2018, Vertebrate fossils from the Claron Formation, Sweetwater Creek area, Garfield County, Utah: Rocky Mountain Geology, v. 53, p. 113–127, <https://doi.org/10.24872/rmgjournal.53.2.113>.
- Eaton, J.G., Laurin, J., Kirkland, J.I., Tibert, N.E., Leckie, R.M., Sageman, B.B., Goldstrand, P.M., Moore, D.W., Straub, A.W., Cobban, W.A., and Dalebout, J.D., 2001, Cretaceous and early Tertiary geology of Cedar and Parowan Canyons, western Markagunt Plateau, Utah, in Erskine, M.C., Faulds, J.E., Bartley, J.M., and Rowley, P.D., editors, The geologic transition, High Plateaus to Great Basin—a symposium and field guide (The Mackin Volume): Utah Geological Association and Pacific Section of the American Association of Petroleum Geologists, Utah Geological Association Publication 30, p. 337–363.
- Eaton, J.G., Maldonado, F., and McIntosh, W.C., 1999b, New radiometric dates from Upper Cretaceous rocks of the Markagunt Plateau, southwestern Utah, and their bearing on subsidence histories [abs.]: Geological Society of America Abstracts with Programs, v. 31, no. 4, p. A–11.
- Eaton, J.G., Tibert, N.E., and Biek, R.F., 2011, First mammals and ostracodes from the Paleogene Claron Formation, southwestern Utah [abs.]: Geological Society of America Abstracts with Programs, v. 43, no. 4, p. 77.
- Fillmore, R.P., 1991, Tectonic influence on sedimentation in the southern Sevier foreland, Iron Springs Formation (Upper Cretaceous), southwestern Utah, in Nations, J.D., and Eaton, J.G., editors, Stratigraphy, depositional environments, and sedimentary tectonics of the western margin, Cretaceous Western Interior Seaway: Geological Society of America Special Paper 260, p. 9–25.
- Fleck, R.J., Anderson, J.J., and Rowley, P.D., 1975, Chronology of mid-Tertiary volcanism in High Plateaus region of Utah, in Anderson, J.J., Rowley, P.D., Fleck, R.J., and Nairn, A.E.M., editors, Cenozoic geology of southwestern High Plateaus of Utah: Geological Society of America Special Paper 160, p. 53–62, <https://doi.org/10.1130/SPE160-p53>.
- Goldstrand, P.M., 1990, Stratigraphy and paleogeography of Late Cretaceous and Early Tertiary rocks of southwest Utah: Utah Geological Survey Miscellaneous Publication 90-2, 58 p, <https://doi.org/10.34191/MP-90-2>.
- Goldstrand, P.M., 1991, Tectonostratigraphy, petrology, and paleogeography of the Upper Cretaceous to Eocene rocks of southwest Utah: Reno, University of Nevada, Ph.D. dissertation, 205 p.
- Goldstrand, P.M., 1992, Evolution of Late Cretaceous and early Tertiary basins of southwest Utah based on clastic petrology: Journal of Sedimentary Petrology, v. 62, no. 3, p. 495–507, <https://doi.org/10.1306/D4267933-2B26-11D7-8648000102C1865D>.
- Goldstrand, P.M., 1994, Tectonic development of Upper Cretaceous to Eocene strata of southwestern Utah: Geological Society of America Bulletin, v. 106, p. 145–154, [https://doi.org/10.1130/0016-7606\(1994\)106%3C0145:TDOUCT%3E2.3.CO;2](https://doi.org/10.1130/0016-7606(1994)106%3C0145:TDOUCT%3E2.3.CO;2).
- Goldstrand, P.M., and Mullett, D.J., 1997, The Paleocene Grand Castle Formation—a new formation on the Markagunt Plateau of southwestern Utah, in Maldonado, F., and Nealey, L.D., editors, Geologic studies in the Basin and Range–Colorado Plateau transition zone in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1995: U.S. Geological Survey Bulletin 2153, p. 59–78, <https://doi.org/10.3133/b2153>.

- Holliday, M.E., Rivera, T., Jicha, B., Traylor, R.B., Biek, R.F., Braunagel, M.J., Griffith, W.A., Hacker, D.B., Malone, D.H., and Mayback, D.F., 2022, Emplacement age of the Markagunt gravity slide in southwestern Utah, USA: *Terra Nova*, 7 p., <https://doi.org/10.1111/ter.12630>.
- Hunt, G.J., Biek, R.F., DeBlieux, D.D., Madsen, S.K., Milner, A.R.C., Roberts, E.M., and Eaton, J.G., 2011, Dinosaur tracks confirm Late Cretaceous age for the lower Grand Castle Formation, Iron County Utah—stratigraphic and tectonic implications [abs.]: *Geological Society of America Abstracts with Programs*, v. 43, no. 4, p. 16.
- Hurlow, H.A., 2002, The geology of Cedar Valley, Iron County, Utah, and its relation to ground-water conditions: *Utah Geological Survey Special Study 103*, 74 p., 2 plates, <https://doi.org/10.34191/SS-103>.
- Johnson, B.T., 1984, Depositional environments of the Iron Springs Formation, Gunlock, Utah: *Brigham Young University Geology Studies*, v. 31, pt. 1, p. 29–46.
- Knudsen, T.R., 2014, Interim geologic map of the Enoch quadrangle, Iron County, Utah: *Utah Geological Survey Map 628*, 12 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/OFR-628>.
- Knudsen, T.R., 2024a, Interim geologic map of the Parowan quadrangle, Iron County, Utah: *Utah Geological Survey Open-File Report 764*, 21 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/OFR-764>.
- Knudsen, T.R., 2024b, Geologic map of the Cedar City 7.5' quadrangle, Iron County, Utah: *Utah Geological Survey Map 302DM*, 18 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/M-302DM>.
- Korth, W.W., and Eaton, J.G., 2004, Rodents and a marsupial (*Mammalia*) from the Duchesnean (Eocene) of the Sevier Plateau, Utah, in Dawson, M.R., and Lillegraven, J.A., editors, *Fanfare for an uncommon paleontologist—papers on vertebrate evolution in honor of Malcolm C. McKenna*: *Carnegie Museum of Natural History Bulletin*, no. 36, p. 109–119, [https://doi.org/10.2992/0145-9058\(2004\)36\[109:RAAMMF\]2.0.CO;2](https://doi.org/10.2992/0145-9058(2004)36[109:RAAMMF]2.0.CO;2).
- Laurin, J., and Sageman, B.B., 2001, Tectono–sedimentary evolution of the western margin of the Colorado Plateau during the latest Cenomanian and early Turonian, in Erskine, M.C., Faulds, J.E., Bartley, J.M., and Rowley, P.D., editors, *The geologic transition, High Plateaus to Great Basin—a symposium and field guide, The Mackin Volume*: *Utah Geological Association and Pacific Section of the American Association of Petroleum Geologists, Utah Geological Association Publication 30*, p. 57–74.
- Laurin, J., and Sageman, B.B., 2007, Cenomanian-Turonian coastal record in SW Utah, U.S.A.—orbital-scale transgressive-regressive events during oceanic anoxic event II: *Journal of Sedimentary Research*, v. 77, p. 731–756, <https://doi.org/10.2110/jsr.2007.076>.
- Lawton, T.F., Pollock, S.L., and Robinson, R.A.J., 2003, Integrating sandstone petrology and nonmarine sequence stratigraphy—application to the Late Cretaceous fluvial systems of southwestern Utah, U.S.A.: *Journal of Sedimentary Research*, v. 73, no. 3, p. 389–406, <https://doi.org/10.1306/100702730389>.
- LeBas, M.J., LeMaitre, R.W., Streckeisen, A., and Zanettin, B., 1986, A chemical classification of volcanic rocks based on the total alkali-silica diagram: *Journal of Petrology*, v. 27, p. 745–750, <https://doi.org/10.1093/petrology/27.3.745>.
- Mackin, J.H., 1960, Structural significance of Tertiary volcanic rocks in southwestern Utah: *American Journal of Science*, v. 258, no. 2, p. 81–131, <https://doi.org/10.2475/ajs.258.2.81>.
- Maldonado, F., 1995, Decoupling of mid-Tertiary rocks, Red Hills-western Markagunt Plateau, southwestern Utah, in Scott, R.B., and Swadley, W.C., editors, *Geologic studies in the Basin and Range-Colorado Plateau transition in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1992*: *U.S. Geological Survey Bulletin 2056*, p. 233–254, <https://doi.org/10.3133/b2056>.
- Maldonado, F., and Moore, R.C., 1995, Geologic map of the Parowan quadrangle, Iron County, Utah: *U.S. Geological Survey Geologic Quadrangle Map 1762*, 1 plate, scale 1:24,000, <https://doi.org/10.3133/gq1762>.
- Maldonado, F., Sable, E.G., and Nealey, D.L., 1997, Cenozoic low-angle faults, thrust faults, and anastomosing high-angle faults, western Markagunt Plateau, southwestern Utah, in Maldonado, F., and Nealey, L.D., editors, *Geologic studies in the Basin and Range-Colorado Plateau transition zone in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1995*: *U.S. Geological Survey Bulletin 2153*, p. 151–176, <https://doi.org/10.3133/b2153>.
- Maldonado, F., and Williams, V.S., 1993a, Geologic map of the Parowan Gap quadrangle, Iron County, Utah: *U.S. Geological Survey Geologic Quadrangle Map 1712*, 1 plate, scale 1:24,000, <https://doi.org/10.3133/gq1712>.
- Maldonado, F., and Williams, V.S., 1993b, Geologic map of the Paragonah quadrangle, Iron County, Utah: *U.S. Geological Survey Geologic Quadrangle Map 1713*, 1 plate, scale 1:24,000, <https://doi.org/10.3133/gq1713>.
- Malone, D., Stevens, G., Lesmann, S.R., Rivera, T., Biek, R., Braunagel, M., Griffith, W.A., Hacker, D., and Rowley, P., 2025, Claron basin provenance shift from Sevier-Laramide compression to Basin and Range extension—detrital zircon

- geochronology from the Eocene-Oligocene Brian Head Formation, Utah: *Geology of the Intermountain West*, v. 12, p. 155–168, <https://doi.org/10.31711/giw.v12.pp155-168>.
- Mayback, D.F., Braunagel, M.J., Malone, D.H., Griffith, W.A., Holliday, M.E., Rivera, T.A., Biek, R.F., Hacker, D.B., and Rowley, P.D., 2022, The concept of tectonic provenance—case study of the gigantic Markagunt gravity slide basal layer: *Terra Nova*, 9 p., <https://doi.org/10.1111/ter.12608>.
- Milner, A.R.C., Vice, G.S., Harris, J.D., and Lockley, M.G., 2006, Dinosaur tracks from the Upper Cretaceous Iron Springs Formation, Iron County, Utah, *in* Lucas, S.G., and Sullivan, R.M., editors, *Late Cretaceous vertebrates from the Western Interior: New Mexico Museum of Natural History and Science Bulletin* 35, p. 105–113.
- Moore, D.W., and Straub, A.W., 2001, Correlation of Upper Cretaceous and Paleogene(?) rocks beneath the Claron Formation, Crow Creek, western Markagunt Plateau, southwest Utah, *in* Erskine, M.C., Faulds, J.E., Bartley, J.M., and Rowley, P.D., editors, *The geologic transition, High Plateaus to Great Basin—a symposium and field guide (The Mackin Volume): Utah Geological Association and Pacific Section of the American Association of Petroleum Geologists, Utah Geological Association Publication* 30, p. 75–95.
- Mullett, D.J., 1989, Interpreting the early Tertiary Claron Formation of southern Utah [abs.]: *Geological Society of America Abstracts with Programs*, v. 21, no. 5, p. 120.
- Mullett, D.J., and Wells, N.A., 1990, Soil fabrics and horizontal cracking in the Paleogene Claron Formation of southern Utah [abs.]: *Geological Society of America Abstracts with Programs*, v. 22, no. 7, p. 335.
- Mullett, D.J., Wells, N.A., and Anderson, J.J., 1988a, Early Cenozoic deposition in the Cedar-Bryce depocenter—certainties, uncertainties, and comparisons with other Flagstaff-Green River basins [abs.]: *Geological Society of America Abstracts with Programs*, v. 20, no. 3, p. 217.
- Mullett, D.J., Wells, N.A., and Anderson, J.J., 1988b, Unusually intense pedogenic modification of the Paleocene-Eocene Claron Formation of southwestern Utah [abs.]: *Geological Society of America Abstracts with Programs*, v. 20, no. 5, p. 382.
- Nealey, L.D., Budahn, J.R., Maldonado, F., and Unruh, D.M., 1997, Geochemistry and petrogenesis of Quaternary basaltic rocks from the Red Hills and western Markagunt Plateau, southwestern Utah, *in* Maldonado, F., and Nealey, L.D., editors, *Geologic studies in the Basin and Range-Colorado Plateau transition in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1995: U.S. Geological Survey Bulletin* 2153-I, p. 177–198, <https://doi.org/10.3133/b2153>.
- Peterson, F., 1969, Four new members of the Upper Cretaceous Straight Cliffs Formation in southeastern Kaiparowits region, Kane County, Utah: *U.S. Geological Survey Bulletin* 1274-J, p. 1–28, <https://doi.org/10.3133/b1274J>.
- Pollock, S.L., 1999, Provenance, geometry, lithofacies, and age of the Upper Cretaceous Wahweap Formation, Cordilleran foreland basin, southern Utah: *Las Cruces, New Mexico State University*, 117 p.
- Primm, J.W., Johnson, C.L., and Stearns, M., 2018, Basin-axial progradation of a sediment supply driven distributive fluvial system in the Late Cretaceous southern Utah foreland: *Basin Research*, v. 30, no. 2, p. 249–278, <https://doi.org/10.1111/bre.12252>.
- Rivera, T., Holliday, M., Jicha, B., Malone, D.H., Braunagel, M.J., Franco, V.A.B., Biek, R.F., Griffith, W.A., and Hacker, D.B., 2025, Emplacement age of the Sevier gravity slide, Utah, USA: *Geochronology*, v. 7, p. 35–44, <https://doi.org/10.5194/gchron-7-35-2025>.
- Rivera, T., Malone, D.H., Biek, R.F., Lesmann, S.R., Stevens, G., Hacker, D.B., Braunagel, M.J., and Griffith, W.A., in review, The growth and collapse of a volcanic field—detrital zircon U-Pb geochronology and provenance of the Oligocene Bear Valley Formation, south-central Utah, USA: *The Sedimentary Record*.
- Rowley, P.D., 1975, Geologic map of the Enoch NE quadrangle, Iron County, Utah: *U.S. Geological Survey Geologic Quadrangle Map* GQ-1301, scale 1:24,000, <https://doi.org/10.3133/gq1301>.
- Rowley, P.D., 1976, Geologic map of the Enoch NW quadrangle, Iron County, Utah: *U.S. Geological Survey Geologic Quadrangle Map* GQ-1302, scale 1:24,000, <https://doi.org/10.3133/gq1302>.
- Rowley, P.D., Biek, R.F., Hacker, D.B., Vice, G.S., McDonald, R.E., Maxwell, D.J., Fasselin, R., Dustin, J., Cunningham, C.G., Steven, T.A., Anderson, J.J., Ekren, E.B., Machette, M.N., Wardlaw, B.R., Smith, Z.D., Kirby, S.M., Knudsen, T.R., Kleber, E.J., Hiscock, A.I., Malone, D.H., Rivera, T.A., and Jicha, B.R., in review, Geologic map of the Beaver 30' x 60' quadrangle, Beaver, Piute, Iron, and Garfield Counties, Utah: *Utah Geological Survey Map*, scale 1:62,500.
- Rowley, P.D., Biek, R.F., Sable, E.G., Boswell, J.T., Vice, G.S., Hatfield, S.C., Maxwell, D.J., and Anderson, J.J., 2013, Geologic map of the Brian Head quadrangle, Iron County, Utah: *Utah Geological Survey Map* 263DM, 38 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/M-263dm>.

- Rowley, P.D., Cunningham, C.G., Steven, T.A., Mehnert, H.H., and Naeser, C.W., 1998, Cenozoic igneous and tectonic setting of the Marysvale volcanic field, and its relation to other igneous centers in Utah and Nevada, *in* Friedman, J.D., and Huffman, A.C., Jr., coordinators, Laccolith complexes of southeastern Utah—time of emplacement and tectonic setting—workshop proceedings: U.S. Geological Survey Bulletin 2158, p. 167–202, <https://doi.org/10.3133/b2158>.
- Rowley, P.D., Cunningham, C.G., Steven, T.A., Workman, J.B., Anderson, J.J., and Theissen, K.M., 2002, Geologic map of the central Marysvale volcanic field, southwestern Utah: U.S. Geological Survey Geologic Investigations Series Map I-2645-A, scale 1:100,000, <https://doi.org/10.3133/i2645A>.
- Rowley, P.D., Mehnert, H.H., Naeser, C.W., Snee, L.W., Cunningham, C.G., Steven, T.A., Anderson, J.J., Sable, E.G., and Anderson, R.E., 1994a, Isotopic ages and stratigraphy of Cenozoic rocks of the Marysvale volcanic field and adjacent areas, west-central Utah: U.S. Geological Survey Bulletin 2071, 35 p., <https://doi.org/10.3133/b2071>.
- Rowley, P.D., Nealey, L.D., Unruh, D.M., Snee, L.W., Mehnert, H.H., Anderson, R.E., and Gromme, C.S., 1995, Stratigraphy of Miocene ash-flow tuffs in and near the Caliente caldera complex, southeastern Nevada and southwestern Utah, *in* Scott, R.B., and Swadley, W.C., editors, Geologic studies in the Basin and Range–Colorado Plateau transition in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1992: U.S. Geological Survey Bulletin 2056, p. 43–88, <https://doi.org/10.3133/b2056>.
- Rowley, P.D., Shroba, R.R., Simonds, F.W., Burke, K.J., Axen, G.J., and Olmore, S.D., 1994b, Geologic map of the Chief Mountain quadrangle, Lincoln County, Nevada: U.S. Geological Survey Geologic Quadrangle Map GQ-1731, scale 1:24,000, <https://doi.org/10.3133/ofr91135>.
- Rowley, P.D., Steven, T.A., Anderson, J.J., and Cunningham, C.G., 1979, Cenozoic stratigraphic and structural framework of southwestern Utah: U.S. Geological Survey Professional Paper 1149, 22 p., <https://doi.org/10.3133/pp1149>.
- Rowley, P.D., and Threet, R.L., 1976, Geologic map of the Enoch quadrangle, Iron County, Utah: U.S. Geological Survey Geologic Quadrangle Map GQ-1296, scale 1:24,000, <https://doi.org/10.3133/gq1296>.
- Rowley, P.D., Williams, V.S., Vice, G.S., Maxwell, D.J., Hacker, D.B., Snee, L.W., and Mackin, J.H., 2006, Interim geologic map of the Cedar City 30' x 60' quadrangle, Iron and Washington Counties, Utah: Utah Geological Survey Open-File Report 476DM, scale 1:100,000, <https://doi.org/10.34191/OFR-476dm>.
- Sable, E.G., and Hereford, R., 2004, Geologic map of the Kanab 30' x 60' quadrangle, Utah and Arizona: U.S. Geological Survey Geologic Investigations Series I-2655, scale 1:100,000, <https://doi.org/10.3133/i2655>.
- Sable, E.G., and Maldonado, F., 1997, The Brian Head Formation (revised) and selected Tertiary sedimentary rock units, Markagunt Plateau and adjacent areas, southwestern Utah, *in* Maldonado, F., and Nealey, L.D., editors, Geologic studies in the Basin and Range–Colorado Plateau transition zone in southeastern Nevada, southwestern Utah, and northwestern Arizona, 1995: U.S. Geological Survey Bulletin 2153, p. 7–26, <https://doi.org/10.3133/b2153>.
- Sanjuan, J., and Eaton, J.G., 2016, Charophyte flora from the Claron Formation (Aquarius Plateau, southwestern Utah)—biostratigraphic implications: *Micropaleontology*, v. 62, p. 323–330.
- Schinkel, T., 2012, Investigation of the origin of silicified layers within Palaeogene-aged volcanoclastic Brian Head Formation, southern Utah: New Britain, Connecticut, unpublished M.S. thesis, 100 p.
- Steven, T.A., Cunningham, C.G., Naeser, C.W., and Mehnert, H.H., 1979, Revised stratigraphy and radiometric ages of volcanic rocks and mineral deposits in the Marysvale area, west-central Utah: U.S. Geological Survey Bulletin 1469, 40 p., <https://doi.org/10.3133/b1469>.
- Steven, T.A., Morris, H.T., and Rowley, P.D., 1990, Geologic map of the Richfield 1° × 2° quadrangle, west-central Utah: U.S. Geological Survey Miscellaneous Investigations Series Map I-1901, scale 1:250,000, <https://doi.org/10.3133/i1901>.
- Stevens, G., Malone, D., Hacker, D., Braunagel, M., Lesmann, S.R., Biek, R.F., Griffith, W.A., and Rivera, T., 2023, Using zircon geochronology to constrain the age of the Black Mountains gravity slide, Marysvale Volcanic Field, Utah, USA [abs.]: Geological Society of America Abstracts with Programs, v. 55, no. 6, p. 9, <https://doi.org/10.1130/abs/2023AM-391831>.
- Szwarc, T.S., Johnson, C.L., Stright, L.E., and MacFarlane, C.M., 2015, Interactions between axial and transverse drainage systems in the Late Cretaceous Cordilleran foreland basin—evidence from detrital zircons in the Straight Cliffs Formation, southern Utah, USA: Geological Society of America Bulletin, v. 127, nos. 3 and 4, p. 372–392, <https://doi.org/10.1130/B31039.1>.
- Threet, R.L., 1963, Structure of the Colorado Plateau margin near Cedar City, Utah, *in* Heylman, E.B., editor, Guidebook to the geology of southwestern Utah—transition between the Basin-Range and Colorado Plateau provinces: Salt Lake City, Utah, Intermountain Association of Petroleum Geologists, Twelfth Annual Field Conference, p. 104–117.

- Tibert, N.E., Leckie, R.M., Eaton, J.G., Kirkland, J.I., Colin, J.P., Leithold, E.L., and McCormic, M.E., 2003, Recognition of relative sea-level change in Upper Cretaceous coal-bearing strata—a paleoecological approach using agglutinated foraminifera and ostracods to detect key stratigraphic surfaces, *in* Olson, H.C., and Leckie, R.M., editors, Micropaleontological proxies for sea-level change and stratigraphic discontinuities: Society of Economic Paleontologists and Mineralogists Special Publication no. 75, p. 263–299, <https://doi.org/10.2110/pec.03.75.0263>.
- Tilton, T.L., 1991, Upper Cretaceous stratigraphy of the southern Paunsaugunt Plateau, Kane County, Utah: Salt Lake City, University of Utah, Ph.D. dissertation, 2 plates, scale 1:24,000, 162 p.
- Tilton, T.L., 2001a, Geologic map of the Alton quadrangle, Kane County, Utah: Utah Geological Survey Miscellaneous Publication 01-4, 22 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/MP-01-4>.
- Tilton, T.L., 2001b, Geologic map of the Podunk Creek quadrangle, Kane County, Utah: Utah Geological Survey Miscellaneous Publication 01-3, 18 p., 2 plates, scale 1:24,000, <https://doi.org/10.34191/MP-01-3>.
- Utah Geological Survey and Apatite to Zircon, Inc. (UGS and AtoZ), 2013, U-Pb formation-age zircon geochronology results for the Brian Head, Bull Rush Peak, Casto Canyon, Cottonwood Mountain, Hatch, and Haycock Mountain quadrangles, Utah: Utah Geological Survey Open-File Report 621, variously paginated, <https://doi.org/10.34191/OFR-621>.
- Utah Geospatial Resource Center, 2020, Southern Utah lidar elevation data: Online, Utah Geospatial Resource Center, State Geographic Information Database, <https://gis.utah.gov/data/elevation-and-terrain/2020-lidar-southern-utah>.
- Williams, P.L., 1967, Stratigraphy and petrography of the Quichapa Group, southwestern Utah and southeastern Nevada: Seattle, University of Washington, Ph.D. dissertation, 141 p.