UTAH GEOLOGICAL MINERALOGICAL SURVEY

QUARTERLY REVIEW

Vol. 7, No. 2

Service to the State of Utah

May 1973

UGMS Reorganizes Under State Board

Effective July 1, the Utah Geological and Mineralogical Survey and its governing board will be transferred to the Utah Geological and Mineral Survey and a newly created board within the Department of Natural Resources.

The objectives of the reorganized State Survey, although much the same, are more clearly defined and enlarged to include study of geothermal resources, and geologic and topographic hazards that could affect the safety of, or cause economic loss to, the citizens of Utah.

Investigation of various mineral substances contained in State lands will continue, as will collection and distribution of reliable information regarding the mineral industry and mineral resources, topography and geology of the State.

The physical facilities will remain on the University of Utah campus (see *Quarterly Review*, November 1972, p. 3), but the functions, responsibilities and governing board once maintained by the University will be transferred to the Survey and its governing board.

The board shall be comprised of seven members appointed by the governor with the advice and consent of the senate: one knowledgeable in the field of geology as applied to civil engineering, four





Photomicrograph of exfoliation crack in thin section.

City and County Building

Weather Deteriorates Sandstone Face

building.

As part of decision-making plans preliminary to renovation of the exterior of the City and County Building in Salt Lake City, UGMS engineering geologist, Bruce N. Kaliser, conducted a study to evaluate the effects of weathering on the building's dimension stone.

Thirteen one-inch core samples were taken from the Kyune Sandstone face from the first floor level to the tower and analyzed using petrographic and binocular microscopes for mineralogic alteration, chemical decomposition, exfoliation planes and fractures.

According to Kaliser, all evidence points to the natural creation of exfoliation surfaces as the result of water penetrating and repeatedly freezing and thawing in the spaces between the sand grains. Exfoliation frequently is observed in the natural environment in Utah's deserts.

The greatest penetration, 40 mm or 1.6 inches, was indicated from a core on the west side of the building, probably the result of predominant winds aiding the penetration of the stone.

Construction of the historic building took three years and \$900,000 to build in the 1890's.¹ Today, after more than 79 years of use, it would cost \$15 million to build or more than \$3 million to restore, according to figures provided to the City and County commissions by the consulting architect on the project.

¹See UGMS' Special Studies 38, "Engineering Geology of the City and County Building, Salt Lake City, Utah," by B. N. Kaliser for more detail.



Phosphate ore from the Rex Peak area, stockpiled in the Brazier Canyon area near the Frank mine, is hauled to the Leefe plant about 12 miles north.



Stauffer Chemical Co.'s phosphate rock beneficiating plant, Leefe, Wyoming. Rail cars transport concentrate to the Garfield plant in Salt Lake County.

Overall Production Up

Phosphate Industry Grows in Utah Phosphate rock is mined from with open-pit works in the Rex

four general areas in the nation: Florida, North Carolina, Tennessee and the western states of Idaho, Wyoming, Montana and Utah.

According to U. S. Bureau of Mines figures, the U. S. produced about 26 million short tons of phosphate rock in 1965 and approximately 42.4 million short tons, valued at \$222.9 million, in 1972.

The increased production is a direct result of a growing demand for phosphate fertilizer; about 90 percent of Utah's phosphate production, which has increased by an average of more than 19 percent per year in recent years, goes to making phosphate fertilizer.

Utah's operations are controlled by Stauffer Chemical Co. with open-pit works in the Rex Peak area in the Crawford Mountains, Rich County, and 12 miles north of Vernal on the south flank of the Uinta Mountains.

Some 30 truckloads carrying approximately 65 tons of ore are hauled daily to a beneficiating plant at Leefe, Wyoming. Washer methods, put into effect last year, have reduced particulate matter and pollution considerably. The concentrate is then shipped by rail to the division plant at Garfield, Utah.

Stauffer recently announced plans to enlarge their phosphate mining and beneficiating operation at Vernal by some 40 percent. Production volume will increase to nearly 400,000 tons.

Survey Releases Latest Studies

Since November of last year, Utah Survey has published the following (available for purchase in person or by mail from Publications Office, 103 Utah Geological Survey Bldg., University of Utah 84112; add 10 percent for handling if ordering by mail):

Special Studies 41, "A Sulfur Isotopic Study of Uinta Basin Hydrocarbons," by R. L. Mauger, R. B. Kayser and J. W. Gwynn (\$1.50). Results of sulfur isotope analyses of tar from oilimpregnated sandstones, crude oils and solid hydrocarbons from the Uinta Basin (Tertiary), Utah, are given and the δS^{34} values are related to origin of the tars and the stratigraphy of the area.

Special Studies 42, "Geology and Diatremes of Desert Mountain, Utah," by D. C. Rees, M. P. Erickson and J. A. Whelan (\$1.50). The geomorphology, stratigraphy, geologic and regional structure and hydrothermal alteration of the Desert Mountain area in Juab County, Utah, are detailed. A geologic map in a back pocket accompanies the text.

Special Studies 43, "Geochemical Reconnaissance at Mercur, Utah," by G. W. Lenzi (\$1.50). Samples mainly from the Mercur, Sunshine and West Mercur areas were assayed for gold and silver and some sample splits not used in the commercial assay were analyzed by atomic-absorption spectrometry for copper and silver. Anomalously high values are located.

Earth Science Series No. 3, "The Earth Below the Hall of Justice, Salt Lake City, Utah," by Richard Van Horn (\$.10 postage). Interesting bits of information about Lake Bonneville and the whole Salt Lake City area were revealed in the excavation for the basement of the Hall of Justice. A geologic cross section and airphoto illustrate the explanation of how the underlying structure came to be.

Oil and Gas Field Studies 4 through 12, by P. R. Peterson: No. 4, "Salt Wash Field" (\$1.00); No. 5, "Bridger Lake Field" (\$1.50); No. 6, "Horseshoe Bend Field" (\$1.50); No. 7, "Upper Valley Field" (\$1.50); No. 8, "Castle Peak and Monument Butte Fields" (\$1.50); No. 9, "Fence Canyon Field" (\$1.00); No. 10, "Cedar Rim Area" (\$1.50); No. 11, "Flat Rock Area" (\$1.00); and No. 12, "Bluebell Field" (\$2.00).

A map depicting the structure of the field, stratigraphic data where significant to production, and a type log cover each field. Accompanying text details the field's discovery, development, reservoir data and production statistics, and describes its geology.

(continued on page 13)

Page 2

At Home With Geology

Water Drainage Damages Home Sites

by B. N. Kaliser UGMS Engineering Geologist

The surface water drainage of any prospective subdivision or single home site should be examined critically to evaluate hazardous or potentially damaging situations. A perennial stream on or bordering the site could flood periodically; a dry drainage or swale cutting the site could disgorge floodwaters and debris in the aftermath of a summer cloudburst.

The direction and stability of the drainage channels help define potential hazards as does a uniform or changing slope. The slope determines the velocity of the flow which, in turn, determines whether the sediment load increases or decreases. An increase could denote an erosion

condition, a decrease, a sedimentation or clean-up problem. Evidence of relatively recent material deposited on the site could signal possible recurrent sedimentation or clean-up problems in the future. Landscaping could be ruined and swimming pools filled with debris. Evidence of removal of material from the site could portend future erosion which can undermine pavement, landscaping and the house foundation itself. Water seeping out onto the surface or infiltrating into the subsurface also can have deleterious effects.

Using data about the site's drainage basin-its size and runoff characteristics, upstream and downstream-engineers can design storm sewers to accommodate

Summary Details Oil and Gas Trends

In a unique summary of worldwide onshore and offshore oil and gas and selected mineral resources, John P. Albers, associate chief geologist, USGS, Washington, D. C., analyzes salient data and shows historical trends and possible future directions of new developments.

Available as USGS Professional Paper 817 (\$2.10 at the USGS Inquiries Office, 125 S. State, Salt Lake City, Utah), the report highlights the following:

Annual production of oil in the United States rose from about 2,900 million barrels in 1960 to 4,091 million barrels in 1971. During the same period gas production rose from 14,945 billion cubic feet to 23,905 billion cubic feet.

Proved oil reserves for the world total about 634 billion barrels (511 onshore and 101 offshore plus an additional 22 billion for which onshore-offshore distribution data are not available); for the United States, 45.3 billion barrels (39.6 onshore and 5.7 offshore); and for the Soviet Union, about 75 billion barrels (less than 73.6 onshore and greater than 1.4 offshore). Estimates of total potential oil resources, onshore and offshore, for both the USA and USSR fall within the same broad category: between 100 billion barrels and 1,000 billion barrels.

Asia (including USSR) and North America are the dominant oil producing regions of the world today, and also have the largest reserves for the world of tomorrow. Asia alone, however, contains more than 70 percent of all the reserves of the 120 countries.

The major oil and gas production is shifting from North America to Africa and Asia.

Major gas resources within Asia are located in the USSR. The center of the world supply and production may shift from North America to the area of the USSR east of the Ural Mountains.

The recent trend of increasing production of oil throughout the world is evidenced by the "doubling time" (time required to double production, using 1960 as the base production period). The doubling time is about 2 years for Africa; 7 years for Asia; 2 years for Oceania; and an estimated 20 years for both North and South America. The relatively short doubling period and large production for Africa and Asia explain why the producing center of oil is rapidly shifting to those continents.



Sediment deposited over mobile home pad on Bonneville Terrace in Davis County. Note typical fan shape of deposit over break-in-slope.

flooding. The creation of impermeable surfaces, as in a new subdivision, however, aggravates the flooding hazard. Sediment and debris is carried farther down the drainage basin before being deposited in what might already be an urbanized area.

Streets become avenues of drainage and should complement the natural drainage system. Adequately designed culverts and bridges must be provided so that flood flows are not impeded. Impounding waters behind such structures can inundate homes and create a general hazard in itself.

Canyon Named To Honor Eardley

The U.S. Board on Geographic Names approved the name Eardley Canyon for a feature in Emery County, Utah.

Howard Ritzma, UGMS petroleum geologist, with the recommendations of several prominent geologists, proposed the name to honor Dr. Armand J. Eardley, a noted Utah geologist who passed away last November. Eardley descended into the canyon in 1930 while a member of a USGS field party and measured and described the older rocks in the deepest parts of the canyon.

Eardley Canyon, 3 miles long along the course of Straight Wash, lies 24 miles southwest of the city of Green River.

Utah Mineral Production-1971

compiled by Carlton H. Stowe UGMS Minerals Information Specialist

Mineral production and values in Utah for 1971, compiled by the U. S. Bureau of Mines, are given below. For a comparison with 1970 figures, see February 1972 *Quarterly Review*, p. 5.

Commodity		Value	Quantity	
BEAT	VER C	OUNTY		
Conner		6	6	
Pumice		W ¹	W	
Sand and gravel		6	6	
TOTAL	s	1,946,000		
BOXE	IDER	COUNTY		
DOAL		COUNT		
Lime		W	W	
Sand and gravel	e	445 000	635 000	* + 1
Stone		W	055,000 W	p. L.
TOTAL		W		
CAC	UF CO	MINTY		
	THE C	JUNII	and the second second	
Lime		W	W	
Stone (limestone quartzite)	\$	620,000	647,000	S. t.
TOTAI		W/	209,000	De te
IOIAL				
CARI	BON C	COUNTY		
Carbon dioxide		W	W	
Coal		W	3,608,000	s. t.
Natural gas		W	W	
Sand and gravel		W	W	
TOTAL		W		
DAGG	ETT (COUNTY		
Natural gas		W	W	
Petroleum		W	6,000	
Sand and gravel		W	W	
Stone		W	W	
TOTAL		W		
DAV	VIS CO	UNTY		
Sand and gravel	\$	875,000	1,193,000	s. t.
TOTAL	s	875,000		
DUCH	ESNE	COUNTY		
Natural gas		W	W	
Petroleum		W	2,984,000	
Sand and gravel		W	W	
Stone		W	W	
TUTAL		W		
EME	RY C	OUNTY		
Coal		W	836.000	s.t.
Natural gas		W		
Petroleum		W	5,000	
Sand and gravel		W	W	
Uranium		W	W	
Vanadium		W	W	
TOTAL	\$	6,150,000		

GARFI	ELD	COUNTY	5.012-21	
Petroleum		w	1.948.000	
Sand and gravel		W	W	
Uranium		W	W	
Vanadium		W	W	
TOTAL	\$	6,045,000		
GRA	ND C	OUNTY		
Copper		6	6	
Natural gas		W	9.1	8 BCF ³
Petroleum		W	116,000	
Potassium salts		W		
Sand and gravel		W	W	
Uronium		W	W	
Vanadium		w	w	
TOTAL		2 710 000	1000	
IUIAL	3	2,710,000		
IKU	NCO	UNII	117	
Pumice		W	W	
Sand and gravel		W	w	
Stone		W	W	
TOTAL	\$ 1	2,049,000		
JUA	B CO	UNTY		
Clays		W	W	
Copper		6	6	
Fluorspar		W	W	
Sand and gravel	s	31.000	47.000	s. t.
Silver	1.00	6	6	
Stone		W	W	
Zinc				
TOTAL	\$	884,000		
KAN	NE CO	UNTY		
Coal		w	12 000	
Pumice		Ŵ	12,000 W	Se L.
Sand and gravel	,	W	31.000	s. t.
TOTAL	\$	140,000		
MILLA	RDC	OUNTY		
Descriting one contrates		w	W	
Pumice		W	W	
Sand and gravel		w	w	
TOTAL		W		
MORC	IAN (OUNTY		
Coment masoner and portland		w	W	
Sand and gravel	e	36 000	21 000	. +
Stone		W	W	0. 6.
TOTAL		W		
PITT	F CO	UNTY		
Clove		W	W	
Copper		W	W	
Gold	s	13.778	334	t. 0.4
Lead		208,739	756	s. t.
Sand and gravel		29,000	51,000	s. t.
Silver		151,234	97,823	t. o.
ZIIIC		313,371	973	S. T.
TOTAL	\$	715,000		
RIC	H CO	UNTY		
Phosphate rock		W	w	
Sand and gravel		W	W	
TOTAL				

(continu	ued fro	om page 4)		
SALT	LAKE	COUNTY		
Cement, portland		W	W	
Copper	\$2	66,399,068	256,153	s. t.
Gold		12,757,141	309,264	t. o.
Lead		4,788,489	17,350	s. t.
Lime		W	W	
Molybdenum		W	W	
Salt Sand and group!		2 296 000	2 014 000	
Sand and graver		5,200,000	3,014,000	5. 6.
Stone		W	3,331,023 W	1. 0.
Zinc		9 737	3 135 283	s.t.
TOTAL	¢3	18 010 000	5,150,200	
SAN I	TIAN A	COUNTY		
Gamma	Unit	6		
Notural con		W	7	DOE
Natural gas		W	. (. (0 BCF
Petroleum		W	11 485 000	
Sand and gravel		w	W	
Silver		6	6	
Uranium		W	W	
Vanadium		W	W	
TOTAL	\$:	54,562,000		
SANP	ETE C	COUNTY		
G			W	
Clays		W	W	
Sand and gravel		W	W	
TOTAL		W		
SEVI	ER CO	OUNTY		
Clavs		w	w	
Coal		W	158,000	s. t.
Gypsum		W	W	
Salt		W	W	
Sand and gravel		W	W	
TOTAL	\$.	2,324,000		
SUMI	MIT C	OUNTY		
Classe		w	W	
Coal		W	12 000	. +
Conner	\$	12 948	12,000	s.t.
Gold		6,146	149	t. o.
Lead		247,848	898	s. t.
Natural gas		W	W	
Petroleum		W	841,000	
Sand and gravel		W	W	
Silver		110,357	71,382	t. o.
Stone		W 000 077	W	
Zinc		282,877	879	s. t.
TOTAL	\$	5,747,000		
TOOL	ELE C	OUNTY		
Conner		6	6	
Gold		6	6	
Lead		6	6	
Lime		W	W	
Potassium salts		W	W	
Salt		W	W	
Sand and gravel		W	W	
Silver		6	6	
Stone (limestone, dolomite,		070 000	887	
marble)	•	970,000	W	
Vinc		6	6	
TOTAL	\$	8.911.000		
TOTAL	ATT	0,711,000		
UINT	AHC	UUNTY		
Asphalt and related bitumens		W	W	
Gilsonite		W	W	DOT
Natural gas		W	17.6	BCF
Natural gas liquids		W	W	

Petroleum		W	6,244,000	
Phosphate rock		W	W	
Sand and gravel		W	W	
Suver		W	W	
TOTAL.	\$	30,790,000		
	UTAH C	OUNTY		
Clay		W	W	
Gold		W	W	
Lead		W	W	
Lime		W	W	
Manganiferous ore		W	112	1. t. ⁵
Sand and gravel	\$	644,000	813,000	s. t.
Stone		W	W	
Zinc		W	W	
TOTAL	\$	9,099,000		
W.	ASATCH	COUNTY		
Copper	2	1.445.600	1.390	s. t.
Gold	*	2.411.352	58,457	t. o.
Lead		1 321 998	4.790	s t
Sand and gravel		W	W	0
Silver		891 521	576 663	to
Stone		W	W	
Zinc		878 094	2.727	s.t.
TOTAL		010,021	-,	
TOTAL		-		
WAS	HINGTO	ON COUNTY		
Petroleum		W	W	
Pumice		W	W	
Sand and gravel Stone	\$	72,000 W	36,000 W	s. t.
TOTAL		_		
		COUNTY		
	WAINE	COUNTI		
Sand and gravel Uranium	\$	28,000 W	103,000 W	s. t.
TOTAL	\$	28,000		
	VEBER C	OUNTY		
-	TEDER C			
Clays		W	W	
Magnesium compounds		W	W	
Potassium salts		W	W	
Salt		W	W	
Sodium sulfate	\$	657,000 W	611,000 W	S. T.
TOTAL	\$	5,286,000		
Undistributed ¹⁺⁶				
Copper	\$	6,131,216	5,896	s. t.
Gemstones		90,000		
Gold		32,671	792	t. o.
Lead		3,995,348	14,475	s. t.
Sand and gravel		3,707,000	3,063,000	s. t.
Silver		1,850,535	1,196,984	t. o.
Stone		3,947,000	2,266,000	s. t.
Zinc		3,666,179	11,385	s. t.
Undisclosed values		35,094,051		
TOTAL	\$	58,514,000		
GRAND TOTAL	\$5	525,694,000		
$^{1}W = Withheld to$	avoid	disclosing	individual com	pany

confidential data (by USBM); values are included in county totals. County totals that have been withheld to avoid disclosing individual company confidential data are included with "undistributed." 2 s.t = short tons

³MCF = million cubic feet; BCF = billion cubic feet

⁴t. o. = troy ounces

⁵l. t. = long tons

⁶Production of Beaver, Grand, Juab, San Juan and Tooele counties combined to avoid disclosing individual company confidential data.

Utah Geology In Print 1972

A list of papers appearing in 1972 which pertain to the geology and mineral industry in Utah has been prepared and printed below.

The staff of the University of Utah Engineering and Physical Sciences Library, under the direction of Edith Rich, generously provided the *Quarterly Review* staff with the list of papers. Sylvia Goeltz, UGMS, compiled the subject index.

The papers are listed alphabetically by author and by subject.

The following sources were used to provide information:

Abstracts of North American Geology.

Applied Science and Technology Index.

Engineering Index Monthly 1972.

Geological Society of America, Bibliography and Index of Geology.

Miscellaneous.

University of Utah and Brigham Young University theses.

U. S. Geological Survey Publications, 1972.

Utah Geological and Mineralogical Survey Publications, 1972.

- AMBJAH, R., 1972, An apparatus to determine water dispersion through sand: M. S. thesis, Univ. of Utah.
- ANDERSEN, D. W., 1972, Stratigraphy of the Duchesne River Formation (Eocene-Oligocene?), northern Uinta Basin, northeastern Utah: M. S. thesis, Univ. of Utah; *also* (and M. D. Picard), Utah Geol. and Mineralog. Survey Bull. 97, 29 p.
- ANDERSEN, D. W. and M. D. Picard, 1972, Secondary and post-depositional redbeds of Duchesne River Formation (Eocene-Oligocene?), Uinta Basin, Utah (abs.): Am. Assoc. Petrol. Geol. Bull., v. 56, no. 3, p. 601.
- ANDERSON, J. M., 1972, A study of the possible correlation of strain, tilt and earthquakes: M. S. thesis, Univ. of Utah.
- ARMSTRONG, R. L., 1972, Low-angle (denudation) faults, hinterland of the Sevier orogenic belt, eastern Nevada and western Utah: Geol. Soc. Am. Bull., v. 83, no. 6, p. 1729-1754.
- AUSTIN, L. H. and G. V. Skogerboe, 1970, Hydrologic inventory of the Uintah study unit: Utah Div. Water Res. Lab., Prog. Rept., no. 40-5, 182 p.
- BAGHOOMIAN, O., 1972, Excess pore pressure and *in situ* measurement of shear strength gain in clays: Ph. D. thesis, Univ. of Utah.
- BAILEY, G. B., 1971, Wall rock alteration as a guide to lode deposit ore in the Park City district, Utah: M. S. thesis, Univ. of Iowa.

- BAKER, A. A., 1972a, Geologic map of NE part of Spanish Fork Peak quadrangle, Utah: U. S. Geol. Survey open-file rept.
- 1972b, Geologic map of the Bridal Falls quadrangle, Utah: U. S. Geol. Survey Geol. Quad. Map GQ-998.
- BEUTNER, E. C., 1972, Curvature of the frontal Sevier orogenic belt (abs.): Geol. Soc. Am. Abs., v. 4, no. 7, p. 451.
- BJORKLUND, L. J. and L. J. McGreevy, 1971, Ground-water resources of Cache Valley, Utah and Idaho: Utah Dept. Nat. Res. Tech. Publ., no. 36, 72 p.
- BLAKEY, R. C., 1972, Stratigraphy and depositional environments of Moenkopi Formation in southeastern Utah (abs.): Am. Assoc. Petrol. Geol. Bull., v. 56, no. 3, p. 604.
- BOLKE, E. L. and D. Price, 1972, Hydrologic reconnaissance of the Blue Creek Valley area, Box Elder County, Utah: Utah Dept. Nat. Res. Tech. Publ., no. 37, 38 p.
- BOLKE, E. L. and K. M. Waddell, 1972, Groundwater conditions in the East Shore area, Box Elder, Davis and Weber counties, Utah, 1960-1969: Utah Dept. Nat. Res. Tech. Publ., no. 35, 53 p.
- BOWERS, W. E., 1972, The Canaan Peak, Pine Hollow, and Wasatch formations in the Table Cliff region, Garfield County, Utah: U. S. Geol. Survey Bull. 1331-B, 39 p.
- BRIMHALL, W. H., 1972, Recent history of Utah Lake as reflected in its sediments-a first report: Brigham Young Univ. Geol. Studies, v. 19, pt. 2, p. 121-126.
- BURCHFIEL, B. C. and C. W. Hickcox, 1972, Structural development of central Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 55-66.
- CADIGAN, R. A., 1971a, Geochemical distribution of some metals in the Moenkopi Formation and related strata, Colorado Plateau region: U. S. Geol. Survey Bull. 1344.
- 1971b, Petrology of the Triassic Moenkopi Formation and related strata in the Colorado Plateau region, with a section on stratigraphy by J. H. Stewart: U. S. Geol. Survey Prof. Paper 692.
- CADIGAN, R. A. and D. E. Stuart-Alexander, 1972, Geochemical factor analysis of intrusion breccia and reconstituted rocks of Mule Ear diatreme, San Juan County, Utah: U. S. Geol. Survey Prof. Paper 800-B, p. B125-B135.
- CARPENTER, H. C., H. W. Sohns and G. U. Dinneen, 1970, Oil shale research related to proposed nuclear projects: Sym. on Eng.

with Nuclear Explosives, January 14-16, 1970, v. 2, p. 1364-1375; *also* Engineering Index, v. 1, no. 1, p. 257.

- CASE, J. E. and H. R. Joesting, 1972, Principal facts for gravity stations in the central Colorado Plateau, Arizona, Colorado, New Mexico and Utah: U. S. Dept. Commerce, Natl. Tech. Inf. Serv. Rept. PB2-11306.
- CASHION, W. B. and J. R. Donnell, 1972, Chart showing correlation of selected key units in the organic-rich sequence of the Green River Formation, Piceance Creek Basin, Colorado, and Uinta Basin, Utah: U.S. Geol. Survey Oil Gas Inv. Chart OC-65.
- CHATFIELD, J., 1972, Case history of Red Wash field, Uintah County, Utah, in Stratigraphic oil and gas fields-classification, exploration methods and case histories: Am. Assoc. Petrol. Geol. Mem., no. 16, p. 342-353 (Soc. Explor. Geophys., Spec. Publ., no. 10, 1972).
- CLARKSON, J. F., W. L. Roberts and A. L. Lingard, 1971, Hidalgoite from Gold Hill, Utah: Min. Rec., v. 2, no. 5, p. 212-213.
- COMPTON, R. R., 1972, Geologic map of the Yost quadrangle, Box Elder County, Utah, and Cassia County, Idaho: U.S. Geol. Survey Map I-672.
- CONDIE, K. C. and C. K. Barsky, 1972, Origin of Quaternary basalts from the Black Rock Desert region, Utah: Geol. Soc. Am. Bull. 83, p. 333-352; *also* Applied Science Tech. Index, v. 60, no. 8, p. 171.
- COOK, K. L. and M. W. Major, 1971, Secular strain measurements near the Wasatch fault, Utah (abs.), *in* Symposium on distribution of stresses within the earth, slow deformation, mechanism of earthquakes, pt. 5: Int. Union Geod. Geophys., 15th Gen. Assem., Abs., Moscow, p. 35.
- COOK, K. L. and J. M. Anderson, 1972, Correlation of earthquakes with changes in strain and tilt near the Wasatch fault, Utah (abs.): Geol. Soc. Am. Abs., Cordilleran Sec., 68th Ann. Mtg., v. 4, no. 3, p. 139-140.
- COOK, K. L. and J. W. Berg, Jr., 1972, Principal facts for gravity stations along the central and southern Wasatch Front, Utah: U.S. Dept. Commerce, Natl. Tech. Inf. Serv., PB-206-675, 32 p.
- COOK, K. L. and J. R. Montgomery, 1972, East-west transverse structural trends in eastern Basin and Range Province as indicated by gravity data (abs.): Geol. Soc. Am. Abs., North-Central Sec., 6th Ann. Mtg., v. 4, no. 5, p. 315.
- COOPER, H. F., Jr. and S. E. Blouin, 1971, Dynamic *in situ* rock properties from buried high explosive arrays, *in* Dynamic rock mechanics: Sym. Rock Mech., Proc., no. 12, p. 45-70.
- COVINGTON, H. R., 1972a, Map showing length of freeze-free season in the Salina

(continued from page 6)

quadrangle, Utah: U.S. Geol. Survey Misc. Geol. Inv. Map I-591-E.

_____ 1972b, Map showing springs in the Salina quadrangle, Utah: U.S. Geol. Survey Misc. Geol. Inv. Map I-591-G.

_____1972c, Topographic relief map of the Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-C.

COVINGTON, H. R. and P. L. Williams, 1972a, Map showing normal annual and monthly precipitation in the Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-D.

_____1972b, Surface water map of the Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-F.

- CRITTENDEN, M. D., Jr., 1972a, Geologic map of the Browns Hole quadrangle, Utah: U. S. Geol. Survey Map GQ-968.
- _____1972b, Willard thrust and the Cache allochthon, Utah: Geol. Soc. Am. Bull., v. 83, no. 9, p. 2871-2880.
- CROCKETT, F. J., 1971, A preliminary study of the use of the oxygen isotope ratios as an exploration tool in the Park City district, Utah: M. S. thesis, Univ. of Utah.
- CROSBY, G. W., 1972, Dual origin of Basin and Range faults, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 67-73.
- CULBERTSON, W. C., 1972, Road log from Rock Springs, Wyoming, to Manila, Utah: Mtn. Geol., pt. I, v. 9, no. 2-3, p. 103-109.
- CUMMINGS, K. F., 1972, Oil and gas exploration activity hingeline play, central Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 91-92.
- DAVIDSON, D. F., 1971, Some aspects of geochemistry and mineralogy of Bear Lake sediments, Utah-Idaho: M. S. thesis, Utah State Univ.
- DECKER, E. R., 1972, Heat flow and radioactivity in Colorado, New Mexico, Texas, Utah, and Wyoming (abs.): Eos (Am. Geophys. Union, Trans.), v. 53, no. 4, p. 516-517.
- DE VOTO, R. H. and others, 1972, Tertiary and Cretaceous energy resources of the southern Rocky Mountains, *in* Energy and mineral resources of the southern Rocky Mountains: Mtn. Geol., v. 9, no. 2-3, p. 79-134 (AAPG Field Trip 2).
- DOELLING, H. H., 1972a, Coal in the Sevier-Sanpete region, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 81-89.
- _____1972b, Tertiary strata, Sevier-Sanpete region, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 41-53.

- DOELLING, H. H. and R. L. Graham, 1972a, Southwestern Utah coal fields-Alton, Kaiparowits Plateau and Kolob-Harmony: Utah Geol. and Mineralog. Survey. Mono. Series, no. 1, 333 p.
- _____1972b, Eastern and northern Utah coal fields-Vernal, Henry Mountains, Sego, La Sal-San Juan, Tabby Mountain, Coalville, Henrys Fork, Goose Creek and Lost Creek: Utah Geol. and Mineralog. Survey Mono. Series, no. 2, 409 p.
- DOYURAN, V., 1972, Geology and groundwater resources of Ogden Valley, Utah: Ph. D. thesis, Univ. of Utah.
- DUNRUD, C. R. and B. K. Barnes, 1972, Engineering geologic map of the Geneva mine area, Carbon and Emery counties, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-704.
- EARDLEY, A. J., 1972, Willard thrust and the Cache allochthon, Utah-discussion: Geol. Soc. Am. Bull., v. 83, no. 9, p. 2881.
- EPPICH, G. K., 1972, Aeromagnetic survey of south-central Utah: M.S. thesis, Univ. of Utah.
- ERB, E. E., Jr., 1971, Geology of the volcanic rocks north of Salina Canyon, Salina, Utah: M. S. thesis, Ohio State Univ.
- FRANCIS, G. G., 1972, Stratigraphy and environmental analysis of the Swan Peak Formation and Eureka Quartzite, northern Utah: M. S. thesis, Utah State Univ.
- FURNISS, B. and W. D. Tidwell, 1972, Cycadeoidales from the Cedar Mountain Formation near Moab, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 377.
- GALLOWAY, C. L., 1971, Structural geology of eastern part of the Smithfield quadrangle, Utah: M. S. thesis, Utah State Univ.
- GOEKE, G., 1970, Eine kambrische trilobitenfauna im Millard Co., Utah, USA (A Cambrian trilobite fauna from Millard County, Utah): Aufschluss, v. 21, no. 7-8, p. 273-275.
- GORDON, M., Jr., 1971, Primitive squid gladii from the Permian of Utah: U.S. Geol. Survey Prof. Paper 750-C, p. C34-C38.
- GOSE, W. A. and C. E. Helsley, 1972, Paleomagnetic and rock-magnetic studies of the Permian Cutler and Elephant Canyon formations in Utah: Jour. Geophys. Res., v. 77, no. 8, p. 1534-1548.
- GROMMÉ, C. S., E. H. McKee and M. C. Blake, Jr., 1972, Paleomagnetic correlations and potassium-argon dating of middle Tertiary ash-flow sheets in the eastern Great Basin, Nevada and Utah: Geol. Soc. Am. Bull., v. 83, no. 6, p. 1619-1638.
- GUTSCHICK, R. C. and J. Rodriguez, 1972, Paleosymbionts of the Louisiana Limestone (upper Devonian) and two Cordilleran correlatives (abs.): Geol. Soc. Am. Abs.,

Northeastern Sec., 7th Ann Mtg., v. 4, no. 1, p. 18-19.

- HACKMAN, R. I. and P. L. Williams, 1972, Map showing drainage basins and historic cloudburst floods in the Salina quadrangle: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-N.
- HALE, L. A., 1972, Depositional history of the Ferron Formation, central Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 29-40.
- HAMMOND, D. R., 1971, A petrofabric study of the Tintic Quartzite (lower and middle Cambrian) in the area of the Willard thrust, northern Utah: M. S. thesis, Univ. of Utah.
- HAMMOND, D. R. and W. T. Parry, 1972, Willard thrust-direction of thrusting from quartz deformation lamellae measurements: Geol. Soc. Am. Bull. 83, p. 1103-1106; also Applied Science Tech. Index, v. 60, no. 10, p. 85.
- HANSEN, W. R., 1972, Geology of the Flaming Gorge quadrangle, Utah-Wyoming: U.S. Geol. Survey Geol. Quad. Map GQ-75.
- HARR, N. J., 1972, Phosphatic rocks, in Geologic atlas of the Rocky Mountain region: Rocky Mtn. Assoc. Geol., p. 322-323.
- HAWS, F. W., R. W. Jeppson and A. L. Huber, 1970, Hydrologic inventory of the Weber River study unit: Utah Div. Water Res. Lab., Prog. Rept., no. 40-6, 131 p.
- HAYNES, D. D. and others, 1972, Geology, structure and uranium deposits, Cortez quadrangle, Colorado and Utah: U. S. Geol. Survey Map I-629.
- HELMSTAEDT, H., O. L. Anderson and A. T. Gavasci, 1972, Petrofabric studies of eclogite, spinel-websterite and spinellherzolite xenoliths from kimberlite-bearing breccia pipes in southeastern Utah and northeastern Arizona: Jour. Geophys. Res., v. 77, no. 23, p. 4350-4365.
- HEWITT, W. P., C. H. Stowe and R. R. Stromberg, 1972, Utah's geothermal resourceslocation, potential, and administrative agencies, *in* Geothermal overviews of the western United States: Geotherm. Res. Coun., Davis, Calif., 7 p.
- HINTZE, L. F., 1972, Lower Paleozoic strata in central Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 7-11.
- HITE, R. J., 1972, Shelf carbonate sedimentation controlled by salinity in the Paradox. Basin, southeast Utah, *in* Environments of sandstone, carbonate and evaporite deposition: Mtn. Geol., v. 9, no. 2-3, p. 329-344.
- HITE, R. J., F. W. Cater and J. A. Liming, 1972, Pennsylvanian rocks and salt anticlines, Paradox Basin, Utah and Colorado, *in*

(continued from page 7)

Geologic atlas of the Rocky Mountain region: Rocky Mtn. Assoc. Geol., p. 133-142.

- HITE, R. J. and others, 1972, Carbonate and evaporite facies of the Paradox Basin, *in* Environments of sandstone, carbonate and evaporite deposition: Mtn. Geol., v. 9, no. 2-3, p. 299-328.
- HOGG, N. C., 1972a, High potassium calcalkaline volcanic rocks from the eastern Great Basin (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 382.
- _____1972b, Shoshonitic lavas in west central Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 2, p. 133-184.
- HOGGAN, R. D., 1971, Paleoecology of the Devonian Guilmette Formation in western Utah and east central Nevada: Ph. D. thesis, Brigham Young Univ.
- HOHL, A. H., 1970, Periglacial features and related surficial deposits of Bull Creek Basin, Henry Mountains, Utah: Ph. D. thesis, Johns Hopkins Univ; *also* Dissert. Abs., v. 33B, no. 5, p. 2153-B.
- HOOD, J. W., 1971, Hydrologic reconnaissance of Hansel Valley and northern Rozel Flat, Box Elder County, Utah: Utah. Dept. Nat. Res. Tech. Publ., no. 33, 25 p.
- _____1972, Hydrologic reconnaissance of the Promontory Mountains area, Box Elder County, Utah: Utah. Dept. Nat. Res. Tech. Publ., no. 38, 35 p.
- HOWARD, J. D., 1972, Trace fossils as criteria for recognizing shorelines in stratigraphic record, *in* Recognition of ancient sedimentary environments: Soc. Econ. Paleon. Mineral., Spec. Publ., no. 16, p. 215-225.
- HOWES, R. C., 1972, Geology of the Wildcat Hills, Utah: M. S. thesis, Utah State Univ.
- HURLEY, W. D., Jr., 1972, Slope stability-Echo Canyon conglomerate: Ph. D. thesis, Univ. of Utah.
- HYATT, M. L. and others, 1969, Hydrologic inventory of the Utah Lake drainage area: Utah. Div. Water Res. Lab., Prog. Rept., no. 40-3, 138 p.
- IRWIN, C. D., 1971, Stratigraphic analysis of upper Permian and lower Triassic strata in southern Utah: Am. Assoc. Petrol. Geol. Bull., v. 55, no. 11, p. 1976-2007; also Engineering Index, v. 10, no. 4, p. 237.
- JENSEN, L. E., 1972, Effect of stress gradients on fracture of rock: M. S. thesis, Univ. of Utah.
- JOHN, E. C., 1972, Petrology and petrography of the intrusive igneous rocks of the Levan area, Juab County, Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 97-107.

- JOHNSON, A. H., 1972, Paleomagnetic results from the lower Jurassic (abs.): Eos (Am. Geophys. Union, Trans.), v. 53, no. 4, p. 357.
- JONES, A. M., 1972, Study of internal structures of fine-grained clastic rocks by Xradiography: Brigham Young Univ. Geol. Studies, v. 19, pt. 1, p. 131-158.
- JONES, L. M. and G. Faure, 1972, Strontium isotope geochemistry of Great Salt Lake, Utah: Geol. Soc. Am. Bull., v. 83, no. 6, p. 1875-1879.
- KALISER, B. N., 1972, Environmental geology of Bear Lake area, Rich County, Utah: Utah Geol. and Mineralog. Survey Bull. 96, 32 p.
- KIDSON, E. J., 1971, Palynology and paleoecology of the Buck Tongue of the Mancos Shale (upper Cretaceous) from east central Utah and western Colorado: Ph. D. thesis, Mich. State Univ.; also Dissert. Abs., v. 32B, no. 9, p. 5257-B.
- KURYVIAL, R. J., 1971, Behavior of some minor elements in coexisting plagioclase, orthoclase and biotite from three intrusive bodies (Tertiary) located in the central Wasatch Range (Utah): Ph. D. thesis, Miami Univ.
- LA PRAY, B. A., 1972, Bibliography of U.S. Geological Survey Water-Resources Reports for Utah: Utah Dept. Nat. Res. Inf. Bull., no. 22, 53 p.
- LAWYER, G. F., 1972, Sedimentary features and paleoenvironment of the Dakota Sandstone (early upper Cretaceous) near Hanksville, Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 2, p. 89-120.
- MAIONE, S. J., 1972, Stratigraphy of the Frontier Sandstone Member of the Mancos Shale (upper Cretaceous) on the south flank of the eastern Uinta Mountains, Utah and Colorado: M. S. thesis, Colo. Sch. Mines.
- MAUGER, R. L., 1972, A sulfur isotope study of bituminous sands from the Uinta Basin, Utah (abs.): Int. Geol. Cong. Abs.-Cong. Geol. Int., Resumes, no. 24, p. 168-169.
- MAY, F. E. and A. Traverse, 1971, Palynology of the Dakota Sandstone, Willis Creek Canyon, Paunsaugunt Plateau near Bryce Canyon National Park, southern Utah (abs.): Am. Assoc. Strat. Palynol., Ann. Mtg., Abs. Paper, no. 4, p. 14.
- MC CARTER, M. K., 1972, A correlation of strength and dynamic properties of some clastic sedimentary rocks: Ph. D. thesis, Univ. of Utah.
- MC GREEVY, L. J., 1972, Specificconductance survey of the Malad River, Utah and Idaho: U.S. Geol. Survey Prof. Paper 800-C, p. C239-C241.
- MEDLYN, D. A. and W. D. Tidwell, 1972, Petrified woods from the Morrison Formation near Clay Point, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 394.

- MELTON, R. A., 1972, Paleoecology and paleoenvironments of the Upper Honaker Trail Formation near Moab, Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 2, p. 45-88.
- MERRILL, R. B., J. K. Robertson and P. J. Wyllie, 1972, Dehydration reaction of titanoclinohumite; reconnaissance to 30 kilobars: Earth Planet. Sci. Letter., v. 14, no. 2, p. 259-262.
- MERRILL, R. C., 1972, Geology of the Mill Fork area, Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 1, p. 65-88.
- MIKULICH, M. J. and R. B. Smith, 1972, Subsurface geology of the Great Salt Lake, Utah, from seismic reflection and aeromagnetic surveys (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 395.
- MILLER, J. F., 1971, Conodont evolution and biostratigraphy of the upper Cambrian and lowest Ordovician (Texas, Oklahoma, Nevada and Utah): Ph. D. thesis, Univ. of Wisc., Madison.
- MOLENAAR, C. M., 1972, Petroleum and natural gas-the Paradox Basin, *in* Geologic atlas of the Rocky Mountain region: Rocky Mtn. Assoc. Geol., p. 282-284.
- MONSALVE, O. A., 1972, Geology of the San Arroyo gas field, Grand County, Utah: M. S. thesis, Univ. of Utah.
- MOORE, C. A., 1972, Factors which may affect occurrence of gas in San Juan and Uinta basins, Rocky Mountains (abs.): Am. Assoc. Petrol. Geol. Bull., v. 56, no. 3, p. 640-641.
- MORLEY, L. A., 1972, Electro-osmotic stabilization of mine minerals: Ph. D. thesis, Univ. of Utah.
- MUELLER, G. T. J. and others, 1972, A crustal low-velocity zone in Utah (abs.): Geol. Soc. Am. Abs., Cordilleran Sec., 68th Ann. Mtg., v. 4, no. 3, p. 204.
- MUELLER, S. and M. Landisman, 1971, An example of the unified method of interpretation for crustal seismic data: R. Aston. Soc., Geophys. Jour., v. 23, no. 4, p. 365-371.
- MULLENS, T. E., 1971, Revised correlation of some carboniferous rock units in the Causey Dam quadrangle, Weber County, Utah: U. S. Geol. Survey Prof. Paper 750-D, p. D105-D107.
- NAESER, C. W., 1971, Geochronology of the Navajo-Hopi diatremes, Four Corners area: Jour. Geophys. Res., v. 76, p. 4978-4985.
- NELSON, M. E., 1971, Stratigraphy and paleontology of Norwood Tuff and Fowkes Formation, northwestern Utah and southwestern Wyoming (abs.): Dissert. Abs. Int., v. 32, no. 6, p 3541B.
 - ____1972, Age and stratigraphic relations of the Fowkes Formation and Norwood Tuff,

(continued from page 8)

southwestern Wyoming and northeastern Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 397-398.

- NEWELL, R. A., 1971, Geology and geochemistry of the northern Drum Mountains, Juab County, Utah: M. S. thesis, Colo. Sch. Mines.
- NEWMAN, G. J., 1972, Conodonts and biostratigraphy of the lower Mississippian in western Utah and eastern Nevada (abs.): Geol. Soc. Am. Abs., North Central Sec., 6th Ann. Mtg., v. 4, no. 5, p. 340.
- OLMORE, S. D., 1972, Pillow lavas of the Chinle Formation in the southern Wah Wah Mountains, Beaver County, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 400.
- OLSEN, D. R., 1972, Igneous petrology of White Pine Canyon, Wasatch Mountains, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 400-401.
- OSTERWALD, F. W. and others, 1971, Field instrumentation studies of earth tremors and their geologic environments in central Utah mining areas: U. S. Geol. Survey Prof. Paper 693, 20 p.
- O'SULLIVAN, R. B. and others, 1972, Stratigraphy of the Cretaceous rocks and the Tertiary Ojo Alamo Sandstone, Navajo and Hopi Indian reservations, Arizona, New Mexico, and Utah: U. S. Geol. Survey Prof. Paper 521-E.
- PABST, A. and C. Milton, 1972, Leucosphenite and its occurrence in the Green River Formation of Utah and Wyoming: Am. Min., v. 57, no. 11-12, p. 1801-1822.
- PAUL, C. R. C., 1972, Cheirocystella antiqua gen. et sp. nov. from the lower Ordovician of western Utah, and its bearing on the evolution of the Cheirocrinidae (Rhombifera: Glyptocystitida): Brigham Young Univ. Geol. Studies, v. 19, pt. 1, p. 15-63.
- PETERSON, D. L., 1972, Principal facts for gravity stations in the San Francisco district and adjoining areas, Utah: Natl. Tech. Inf. Serv. Rept. PB2-10681.
- PETERSON, P. R., 1972a, Pariette Bench field: Utah Geol. and Mineralog. Survey Oil and Gas Field Studies, no. 1.
- _____1972b, Agate field area: Utah Geol. and Mineralog. Survey Oil and Gas Field Studies, no. 2.
- _____1972c, Grassy Trail Creek field: Utah Geol. and Mineralog. Survey Oil and Gas Field Studies, no. 3.
- PETERSON, P. R. and H. R. Ritzma, 1972, Oil-impregnated sandstone, Thistle area, Utah County, Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 93-95.

- PICARD, M. D. and L. R. High, Jr., 1972a, Criteria for recognizing lacustrine rocks, *in* Recognition of ancient sedimentary environments: Soc. Econ. Paleon. and Min., Spec. Publ. 16, p. 108-145.
- 1972b, Paleoenvironmental reconstructions in an area of rapid facies change, Parachute Creek Member of Green River Formation (Eocene), Uinta Basin, Utah: Geol. Soc. Am. Bull., v. 83, no. 9, p. 2689-2708.
- PINNELL, M. L., 1972, Geology of the Thistle quadrangle, Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 1, p. 89-130.
- PRATT, H. R., W. S. Brown and W. F. Brace, 1971, *In situ* determination of strength properties in a quartz diorite rock mass, *in* Dynamic rock mechanics: Sym. Rock Mech., Proc., no. 12, p. 27-43.
- PRICE, D., 1972a, Map showing general availability of ground water in the Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-M.
- 1972b, Map showing general chemical quality of ground water in the Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-K.
- RAUP, O. B., 1972a, Bromine distribution in some halite rocks of the Paradox Member, Hermosa Formation, in Utah, *in* Environments of sandstone, carbonate and evaporite deposition: Mtn. Geol., v. 9, no. 2-3, p. 345-354.
- _____1972b, Origin of rare-earths in a marine evaporite mineral (with discussion), *in* Geology of saline deposits-Geologie des depots salins (incl. Fr. summ.): Unesco Earth Sci. Ser., no. 7, p. 103-109.
- RIGBY, J. K. and A. J. Swensen, 1972, Pennsylvanian sponges from the Oquirrh Group of central Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 2, p. 127-132.
- RITZMA, H. R., 1971, Faulting on the north flank of the Uinta Mountains, Utah and Colorado, *in* Symposium on Wyoming tectonics and their economic significance: Wyo. Geol. Assoc. Guidebook, no. 23, p. 145-150.
- 1972a, Petroleum and natural gas-the Uinta Basin, in Geologic atlas of the Rocky Mountain region: Rocky Mtn. Assoc. Geol., p. 276-278.
- _____1972b, Six Utah "hingeline" wells, in Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 75-80.
- ROBISON, R. A., 1972, Hypostoma of agnostid trilobites: Lethaia, v. 5, no. 2, p. 239-248.
- ROBISON, R. A. and L. F. Hintze, 1972, An early Cambrian trilobite faunule from Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 1, p. 3-13.

- ROEHLER, H. W., 1972, Geologic map of the Red Creek Ranch quadrangle, Wyoming, Utah and Colorado: U. S. Geol. Survey Map. GQ-1001.
- ROWLEY, P. D., 1972, Cenozoic history of the Black Mountains, southwestern Utah (abs.): Geol. Soc. Am. Abs., with Prog., v. 4, no. 6, p. 405.
- ROWLEY, P. D. and J. H. Anderson, 1972, Cenozoic normal fault structures of the southwestern high plateaus and adjacent Great Basin, Utah (abs.): Geol. Soc. Am. Abs., with Prog., v. 4, no. 6, p. 406.
- SAVAGE, W. Z. and G. M. Sowers, 1972, Criteria for separating stable and unstable folding-III, Stable bending (abs.): Eos (Am. Geophys. Union, Trans.), v. 53, no. 4, p. 523-524.
- SAXON, F. C., 1972, Water-resource evaluation of Morgan Valley, Morgan County, Utah: M. S. thesis, Univ. of Utah.
- SAYEED, U. A., 1972, A note on the Kern Mountains plutonic complex, White Pine County, Nevada and Juab County, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 407.
- SCHULINGKAMP, W. J., 1972, Petrology of the Ordovician Swan Peak Formation, southeastern Idaho and north central Utah: M. S. thesis, Utah State Univ.
- SHAWE, D. R., 1972, Reconnaissance geology and mineral potential of Thomas, Keg and Desert calderas, central Juab County, Utah: U. S. Geol. Survey Prof. Paper 800-B, p. B67-B77.
- SHEEHAN, P. M., 1971, Silurian Brachiopoda, community ecology and stratigraphic geology in western Utah and eastern Nevada, with a section on late Ordovician stratigraphy: Ph. D. thesis, Univ. of Calif., Berkeley.
- SHUEY, R. T., 1972, Paleomagnetic and paleoclimatic implications of a deep core from Pleistocene Lake Bonneville (abs.): Eos (Am. Geophys. Union, Trans.), v. 53, no. 4, p. 356.
- SMITH, J. W., L. G. Trudell and W. A. Robb, 1972, Oil yields and characteristics of Green River Formation oil shales at Wasco EX-1, Uintah County, Utah: U. S. Bur. Mines Rept. Inv. 7693.
- SMITH, R. B. and others, 1972, Source mechanisms of microearthquakes associated with underground mines in Utah (abs.): Geol. Soc. Am. Abs., Cordilleran Sec., 68th Ann. Mtg., v. 4, no. 3, p. 240.
- SMITH, R. K., 1972, The mineralogy and petrology of the contact metamorphic aureole around the Alta stock, Utah: Ph. D. thesis, Univ. of Iowa; *also* Dissert. Abs., v. 32B, no. 12, p. 7119-B.
- SORENSEN, M. L. and M. D. Crittenden, 1972, Preliminary geologic map of part of the

(continued from page 9)

Wasatch Range near North Ogden, Utah: U. S. Geol. Survey Misc. Field Studies Map MF-428.

- STAATZ, M. H., 1972, Geologic map of the Dugway Proving Ground SW quadrangle, Tooele County, Utah: U. S. Geol. Survey Map GQ-992.
- STANLEY, W. D., 1972, Geophysical study of unconsolidated sediments and basin structure in Cache Valley, Utah and Idaho: Geol. Soc. Am. Bull., v. 83, no. 6, p. 1817-1830.
- STEINER, M. B. and C. E. Helsley, 1972, Jurassic polar movement relative to North America: Jour. Geophys. Res., v. 77, no. 26, p. 4981-4993.
- STEWART, J. H. and others, 1972, Stratigraphy and origin of the Chinle Formation and related upper Triassic strata in the Colorado Plateau region: U. S. Geol. Survey Prof. Paper 690.
- STOKES, W. L., 1972, Stratigraphic problems of the Triassic and Jurassic sedimentary rocks of central Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 21-28.
- STOWE, C. H., 1972a, Energy and the environment-a potential crisis in resources supply: Utah Geol. and Mineralog. Survey Circular 54, 18 p.

(compiler), 1972b, Oil and gas production in Utah to 1970: Utah Geol. and Mineralog. Survey. Bull. 94, 179 p.

- STUART-ALEXANDER, D. E., E. M. Shoemaker and H. J. Moore, 1972, Geologic map of the Mule Ear diatreme, San Juan County, Utah: U. S. Geol. Survey Map I-674.
- SUMSION, C. T. and others, 1972, Groundwater conditions in Utah, spring of 1972: Utah Dept. Nat. Res., Div. Water Res. Coop. Inv. Rept., no. 10, 73 p.
- TANIS, J. I., 1972, Isostasy and crustal thickness in Utah and adjacent areas as revealed by gravity profiles (abs.): Dissert. Abs. Int., v. 32, no. 7, p. 4017B.
- TAYLOR, M. E., 1971, The biostratigraphy and systematics of trilobites (Arthropoda; Tribolita) from the upper Cambrian (upper Franconian-Tempealeauan stages) in the Central Coast Basin, Nevada and Utah: Ph. D. thesis, Univ. of Calif., Berkeley.
- TERRELL, F. M., 1972a, Lateral facies distribution and paleoecology of part of the Honaker Trail Formation (Pennsylvanian) near Potash, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 416.
 - _____1972b, Lateral facies and paleoecology of Permian Elephant Canyon Formation, Grand County, Utah: Brigham Young Univ. Geol. Studies, v. 19, pt. 2, p. 3-44.
- THOMAS, H. E. and T. F. Sterner, 1972, Hotfilm flow logging as applied to *in situ* oil

shale and mining experiments: Soc. Prof. Well Log Anal. Ann. Logging Sym., Trans., no. 13, paper L, 14 p.

- THOMPSON, W. D., 1971, Stratigraphy of black shale facies of Green River Formation (Eocene), Uinta Basin, Utah: M. S. thesis, Univ. of Utah.
- TIDWELL, W. D. and N. Hebbert, 1972, Tempskya from the Cedar Mountain Formation near Moab, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 417-418.
- TIDWELL, W. D., G. Thayn and F. Terrell, 1972, New upper Pennsylvanian fossil plant locality from the Honaker Trail Formation near Moab, Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 417.
- TOLLEY, M. A., 1971, Hydrothermal alteration of ash-flow tuffs in the Indian Peak district of southwestern Utah: M. S. thesis, Univ. of Mo. at Rolla.
- TOOKER, E. W., 1971, Regional structural controls of ore deposits, Bingham mining district, Utah, U. S. A.: Mining Geol. Spec. Issue 3, p. 76-81 (Soc. Mining Geol. Japan, Proc., IMA-IAGOD Mtgs., 1970, IAGOD Vol.).
- TRENT, R. H., 1972, The feasibility of mining a low grade copper deposit in a remote area: M. S. thesis, Univ. of Utah.
- U. S. GEOLOGICAL SURVEY, 1971, Aeromagnetic map of part of west central Utah: U. S. Geol. Survey open-file rept.
- U. S. GEOLOGICAL SURVEY, U. S. BUREAU OF LAND MANAGEMENT AND THE U. S. BUREAU OF MINES, 1971, Base maps of areas underlain by Green River Formation in the eastern part of the Uinta Basin, Utah and Colorado, the southern part of the Green River Basin, Wyoming and Utah, and the Piceance Creek Basin, Colorado: U. S. Geol. Survey open-file rept.
- VAN DE GRAAFF, F. R., 1972, Fluvial-deltaic facies of the Castlegate Sandstone (Cretaceous), east central Utah: Jour. Sed. Petrol., v. 42, no. 3, p. 558-571.
- VAN HORN, R., 1972a, Landslide and associated deposits of the Sugar House quadrangle, Salt Lake County, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-766-D.
- _____1972b, Map showing relative ages of faults in the Sugar House quadrangle, Salt Lake County, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-766-B.
- _____1972c, Relative slope stability map of the Sugar House quadrangle, Salt Lake County, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-766-E.
- ____1972d, Slope map of the Sugar House quadrangle, Salt Lake County, Utah: U.S. Geol. Survey Misc. Geol. Inv. Map I-766-C.
- _____1972e, Surficial geologic map of the Sugar House quadrangle, Salt Lake County,

Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-766-A.

- _____1972f, The earth below the Hall of Justice, Salt Lake City, Utah: Utah Geol. and Mineralog. Survey Earth Sci. Ser., no. 3.
- VAN WEST, F. P., 1972, Green River oil shale, in Geologic atlas of the Rocky Mountain region: Rocky Mtn. Assoc. Geol., p. 287-289.
- VARNEY, P. J., 1972, Depositional environment of the Mineral Fork Formation (Precambrian), Wasatch Mountains, Utah: M. S. thesis, Univ. of Utah.
- WADDELL, K. M. and D. Price, 1972, Quality of surface water in the Bear River Basin, Utah, Wyoming, and Idaho: U. S. Geol. Survey Hydrol. Inv. Atlas HA-417.
- WALLACE, C. A., 1972, A basin analysis of the upper Precambrian Uinta Mountain Group, Utah: Ph. D. thesis, Univ. of Calif., Santa Barbara; also Dissert. Abs., v. 33B, no. 5, p. 2158-B.
- WELSH, J. E., 1972, Upper Paleozoic stratigraphy, in Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 13-20.
- WENGERD, S. A., 1971, Exploration for petroleum along wrench faults-the Paradox Basin model (abs.), in Exploration todayenergy tomorrow (SEG Ann. Int. Mtg., 41st; program and abs.): Soc. Explor. Geophys., p. 56-57.
- WHEELOCK, T. G. B., 1971, Jurassic ophiuroid echinoderms from Utah and Wyoming: M. S. thesis, Univ. of Wyo.
- WHELAN, J. A., 1972a, Ochenius bar theory of saline deposition supported by quantitative data, Great Salt Lake, Utah (abs.): Int. Geol. Cong. Abs.-Cong. Geol. Int., Resumes, no. 24, p. 327-328.
- _____1972b, Ultimate degree of freshening of south arm, Great Salt Lake, Utah, as result of fill construction (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 421.
- WHELAN, J. A. and N. Stauffer, 1972, Preliminary report on possible solutions to "fill effect" causing dilution of south arm brines and concentration of north arm brines, Great Salt Lake, Utah: Utah Geol. and Mineralog. Survey Spec. Studies 40, 12 p.
- WHITE, W. B. and J. J. Van Gundy, 1971, Geological reconnaissance of Timpanogos Cave, Utah (abs.): Natl. Speleol. Soc. Bull., v. 33, no. 4, p. 147-148.
- WILLIAMS, J. S., 1972, Fossil distribution across the middle-upper Devonian boundary, Portage and Logan canyons, north central Utah (abs.): Geol. Soc. Am. Abs., Rocky Mtn. Sec., 25th Ann. Mtg., v. 4, no. 6, p. 421-422.

(continued from page 10)

- WILLIAMS, P. L., 1972a, Map showing relative ease of excavation in the Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-J.
- _____1972b, Map showing types of bedrock and surficial deposits, Salina quadrangle, Utah: U. S. Geol. Survey Misc. Geol. Inv. Map I-591-H.
- WILLIAMSON, C. R., 1972, Carbonate petrology of the Green River Formation (Eocene), Uinta Basin, Utah and Colorado: M. S. thesis, Univ. of Utah.
- WILSON, J. D. S. and A. J. Sinclair, 1969, Q-mode factor analysis applied to mineral exploration data, *in* Decision-making in mineral exploration-II, Proceedings of a symposium: British Columbia Univ., Ext. Dept. Eng. Program, p. 244-257.
- WINKLER, P. L., 1972, Source mechanisms of earthquakes associated with coal mines in eastern Utah: M. S. thesis, Univ. of Utah.
- WOODFILL, R. D., 1972, A geologic and petrographic investigation of a northern part of the Keetley volcanic field, Summit and Wasatch counties, Utah: Ph. D. thesis, Purdue Univ.; *also* Dissert. Abs., v. 33B, no. 2, p. 787-B.
- WOODWARD, L. A., 1972a, Upper Precambrian strata of the eastern Great Basin as potential host rocks for mineralization: Econ. Geol., v. 67, no. 5, p. 677-681.
- _____1972b, Upper Precambrian stratigraphy of central Utah, *in* Plateau-Basin and Range transition zone, central Utah: Utah Geol. Assoc. Publ. 2, p. 1-5.
- YANCY, T. E., 1971, Biostratigraphy, paleoecology and paleontology of the Arcturus Group (Permian; eastern Nevada and western Utah): Ph. D. thesis, Univ. of Calif., Berkeley.
- YOUNG, D. S., 1972, Stress analysis in openpit mines: Ph. D. thesis, Univ. of Utah.

SUBJECT INDEX

ABSOLUTE AGE

Navajo-Hopi diatremes, Four Corners area: Naeser, 1971.

AREAL GEOLOGY

Drum Mountains: Newell, 1971.

- Juab County, calderas: Shawe, 1972.
- Keetley volcanic field, Summit and Wasatch counties: Woodfill, 1972.
- Mill Fork area: R. C. Merrill, 1972.
- Ogden Valley: Doyuran, 1972.
- Road log, Rock Springs to Manila: Culbertson, 1972.
- Thistle quadrangle: Pinnell, 1972.

Wildcat Hills: Howes, 1972.

ECONOMIC GEOLOGY

Bingham mining district, controls: Tooker, 1971.

QUARTERLY REVIEW

- Brines, Great Salt Lake: Whelan and Stauffer, 1972.
- Coal, eastern and northern Utah: Doelling and Graham, 1972b.
- Coal, Sevier-Sanpete region: Doelling, 1972a.
- Coal, southwestern Utah: Doelling and Graham, 1972a.
- Feasibility study, low grade copper deposit: Trent, 1972.
- Geothermal resources, Utah's: Hewitt, Stowe and Stomberg, 1972.
- Great Basin, Precambrian strata: Woodward, 1972a.
- Hot-film flow logging, oil shale: Thomas and Sterner, 1972.
- Moenkopi Formation, Colorado Plateau: Cadigan, 1971a.
- Natural gas, San Juan and Uinta basins: Moore, 1972.
- Oil shale, nuclear projects: Carpenter, Sohns and Dinneen, 1970.
- Oil shales, Green River Formation: Smith, Trudell and Robb, 1972.
- Organic rich deposits, regional: Van West, 1972.
- Organic-rich sequence, Green River Formation: Cashion and Donnell, 1972.
- Paradox Basin, exploration: Wengerd, 1971.
- Petroleum and natural gas, hingeline play: Cummings, 1972.
- Petroleum and natural gas, hingeline wells: Ritzma, 1972b.
- Petroleum and natural gas, Paradox Basin: Molenaar, 1972.
- Petroleum and natural gas, production: Stowe, 1972b.
- Petroleum and natural gas, Thistle area: Peterson and Ritzma, 1972.
- Petroleum and natural gas, Uinta Basin: Ritzma, 1972a.
- Phosphatic rocks, Rocky Mountain region: Harr, 1972.
- Red Wash field, Uintah County: Chatfield, 1972.
- Rock alteration, Park City district: Bailey, 1971.
- Rocky Mountains, energy resources: DeVoto and others, 1972.
- San Arroyo gas field, Grand County: Monsalve, 1972.

ENGINEERING GEOLOGY

- Bear Lake area, Rich County: Kaliser, 1972.
- Clastic sedimentary rocks, strength and dynamic properties: McCarter, 1972.
- Coal mines, earthquakes: Winkler, 1972.

- Echo Canyon conglomerate, slope stability: Hurley, 1972.
- Microearthquakes, underground mines: Smith and others, 1972.
- Open-pit mines, stress analysis: Young, 1972.
- Quartz diorite rock, strength properties: Pratt, Brown and Brace, 1971.
- Rock properties, dynamic rock mechanics: Cooper and Blouin, 1971.
- Stress gradients, fracture of rock: Jensen, 1972.

GENERAL

- Bibliography, USGS water-resources reports: LaPray, 1972.
- Crisis, resource supply: Stowe, 1972a.

GEOCHEMISTRY

Bear Lake sediments: Davidson, 1971.

- Drum Mountains, Juab County: Newell, 1971.
- Great Salt Lake, Ochenius bar theory: Whelan, 1972a.
- Great Salt Lake, south arm: Whelan, 1972b.
- Great Salt Lake, strontium isotope: Jones and Faure, 1972.
- Marine evaporite mineral, rare-earths: Raup, 1972b.
- Minor elements, plagioclase, orthoclase and biotite: Kuryvial, 1971.
- Mule Ear diatreme, San Juan County: Cadigan and Stuart-Alexander, 1972.
- Paradox Member, Hermosa Formation: Raup, 1972a.
- Park City district, oxygen isotope ratios: Crockett, 1971.
- Park City district, Q-mode factor analysis: Wilson and Sinclair, 1969.

Uinta Basin, bituminous sands: Mauger, 1972.

GEOMORPHOLOGY

- Bull Creek Basin, Henry Mountains: Hohl, 1970.
- Timpanogos Cave, geological reconnaissance: White and Van Gundy, 1971.

GEOPHYSICAL SURVEY

- Central Colorado Plateau, gravity stations: Case and Joesting, 1972.
- Earth tremors, field instrumentation studies: Osterwald and others, 1971.
- Gravity stations, Wasatch Front: Cook and Berg, 1972.
- San Francisco district, gravity stations: D. L. Peterson, 1972.
- South central Utah: Eppich, 1972.
- Wasatch fault, secular strain measurements: Cook and Major, 1971.

(continued on next page)

and the second second

(continued from page 11)

Wasatch fault, strain and tilt: Cook and Anderson, 1972.

GEOPHYSICAL SURVEYS

Page 12

- Basin and Range province, structural trends: Cook and Montgomery, 1972.
- Cache Valley: Stanley, 1972.
- Crustal seismic data: Mueller and Landisman, 1971.
- Crustal zone, low-velocity: Mueller and others, 1972.
- Great Salt Lake, subsurface geology: Mikulich and Smith, 1972.

Heat flow and radioactivity: Decker, 1972.

Utah, regional: Tanis, 1972.

HYDROLOGY

- Bear River Basin, surface water: Waddell and Price, 1972.
- Blue Creek Valley, Box Elder County: Bolke and Price, 1972.
- Cache Valley, ground water resources: Bjorklund and McGreevy, 1971.
- East Shore area, Box Elder, Davis and Weber counties: Bolke and Waddell, 1972.
- Ground water, Utah: Sumsion and others, 1972.
- Ground water resources, Ogden Valley: Doyuran, 1972.
- Hansel Valley and Rozel Flat, Box Elder County: Hood, 1971.
- Malad River, Udy Hot Springs: McGreevy, 1972.
- Morgan Valley, Morgan County: Saxon, 1972.
- Promontory Mountains, Box Elder County: Hood, 1972.
- Uintah study unit, hydrologic inventory: Austin and Skogerboe, 1970.
- Utah Lake drainage area: Hyatt and others, 1969.

Water dispersion, apparatus: Ambjah, 1972.

Weber River: Haws, Jeppson and Huber, 1970.

MAPS, AEROMAGNETIC

West central Utah: USGS, 1971.

MAPS, ENGINEERING

- Sugar House quadrangle, faults ages: Van Horn, 1972b.
- Sugar House quadrangle, Salt Lake County: Van Horn, 1972a.
- Sugar House quadrangle, slope map: Van Horn, 1972d.
- Sugar House quadrangle, slope stability: Van Horn, 1972c.

MAPS, ENGINEERING GEOLOGIC

Geneva mine area, Carbon and Emery counties: Dunrud and Barnes, 1972.

MAPS, GENERAL

Green River Formation, base maps: USGS, BLM and Bureau of Mines, 1971.

QUARTERLY REVIEW

- Salina quadrangle, bedrock and surficial deposits: P. L. Williams, 1972b.
- Salina quadrangle, drainage basins and cloudburst floods: Hackman, 1972.
- Salina quadrangle, excavation: P. L. Williams, 1972a.
- Salina quadrangle, freeze-free season: Covington, 1972a.
- Salina quadrangle, ground water: Price, 1972b.
- Salina quadrangle, ground water availability: Price, 1972a.
- Salina quadrangle, precipitation: Covington and Williams, 1972a.
- Salina quadrangle, springs: Covington, 1972b.
- Salina quadrangle, surface water: Covington and Williams, 1972b.
- Salina quadrangle, topographic relief map: Covington, 1972c.

MAPS, GEOLOGIC

- Bridal Falls quadrangle: Baker, 1972b.
- Browns Hole quadrangle: Crittenden, 1972a.
- Cortez quadrangle, miscellaneous geologic investigations: Haynes and others, 1972.
- Dugway Proving Grounds, Tooele County: Staatz, 1972.
- Flaming Gorge quadrangle: Hansen, 1972.
- Mule Ear diatreme, San Juan County: Stuart-Alexander, Shoemaker and Moore, 1972.
- Red Creek Ranch quadrangle: Roehler, 1972.
- Spanish Fork Peak quadrangle: Baker, 1972a.
- Sugar House quadrangle, surficial geologic: Van Horn, 1972e.
- Wasatch Range, North Ogden: Sorensen and Crittenden, 1972.
- Yost quadrangle, Box Elder County: Compton, 1972.

MAPS, OIL AND GAS

- Agate field area: P. R. Peterson, 1972b.
- Grassy Trail Creek field: P. R. Peterson, 1972c.
- Pariette Bench field: P. R. Peterson, 1972a.

MINERALOGY

- Clays, pressure and shear strength: Baghoomian, 1972.
- Leucosphenite, Green River Formation: Pabst and Milton, 1972.
- Mine minerals, electro-osmotic stabilization: Morley, 1972.
- Titanoclinohumite, Moses Rock dike: Merrill, Robertson and Wyllie, 1972.

PALEOMAGNETIC

Ash-flow sheets, Great Basin: Gromme, McKee and Blake, 1972.

May 1973

Cutler and Elephant Canyon formations, Permian: Gose and Helsley, 1972.

Jurassic: Johnson, 1972.

- Jurassic, eastern Utah: Steiner and Helsley, 1972.
- Lake Bonneville, deep core: Shuey, 1972.
- Tertiary ash-flow sheets, Great Basin: Gromme, McKee and Blake, 1972.

PALEONTOLOGY

- Arcturus Group, Permian: Yancy, 1971.
- Brachiopoda, Silurian: Sheehan, 1971.
- Cedar Mountain Formation, Cycadeoidales: Furniss and Tidwell, 1972.
- Cheirocystella antiqua, Ordovician: Paul, 1972.
- Conodont, Cambrian and Ordovician: Miller, 1971.
- Conodonts, Mississippian: Newman, 1972.
- Devonian Guilmette Formation, western Utah: Hoggan, 1971.
- Echinoderms, Jurassic: Wheelock, 1971.
- Fossil distribution, Devonian boundary: J.S. Williams, 1972.
- Honaker Trail Formation, fossil plant locality: Tidwell, Thayn and Terrell, 1972.
- Honaker Trail Formation, paleoecology and paleoenvironments: Melton, 1972.
- Norwood Tuff and Fowkes Formation, northwestern Utah: Nelson, 1971.
- Paleosymbionts, Louisiana Limestone: Gutschick and Rodriguez, 1972.
- Palynology, Dakota Sandstone: May and Traverse, 1971.
- Palynology and paleoecology, Buck Tongue, Mancos Shale: Kidson, 1971.
- Pennsylvanian sponges, Oquirth group: Rigby and Swensen, 1972.
- Squid gladii, Permian: Gordon, 1971.
- Tempskya, Cedar Mountain Formation: Tidwell and Hebbert, 1972.
- Trilobite faunule, Cambrian: Robison and Hintze, 1972.
- Trilobites, Cambrian: Robison and Hintze, 1972; Taylor, 1971.
- Trilobites, Cambrian, Millard County: Goeke, 1970.

PETROLOGY

- Alta stock, contact metamorphic: R. K. Smith, 1972.
- Black Rock Desert region, Quaternary basalts: Condie and Barsky, 1972.
- Breccia pipes, southeastern Utah: Helmsteadt, Anderson and Gavasci, 1972.

May 1973

QUARTERLY REVIEW

(continued from page 12)

- Green River Formation, carbonate: Williamson, 1972.
- Indian Peak district, hydrothermal alteration: Tolley, 1971.
- Keetley volcanic field, Summit and Wasatch counties: Woodfill, 1972.
- Kern Mountains, plutonic complex, Juab County: Sayeed, 1972.
- Levan area, Juab County: John, 1972.
- Moenkopi Formation, Triassic: Cadigan, 1971b.
- Petrified woods, Morrison Formation: Medlyn and Tidwell, 1972.
- Pillow lavas, Chinle Formation, Beaver County: Olmore, 1972.
- Shoshonitic lavas, west central Utah: Hogg, 1972b.
- Swan Peak Formation, north central Utah: Schulingkamp, 1972.
- Volcanic rocks, Salina Canyon: Erb, 1971.
- White Pine Canyon, Wasatch Mountains: Olsen, 1972.

SEDIMENTARY PETROLOGY

Bear Lake sediments: Davidson, 1971.

- Castlegate Sandstone, east central Utah: Van de Graaff, 1972.
- Dakota Sandstone, Upper Cretaceous: Lawyer, 1972.
- Duchesne River Formation, Uinta Basin: Andersen, 1972.
- Elephant Canyon Formation, Grand County: Terrell, 1972b.
- Honaker Trail Formation, Pennsylvanian: Terrell, 1972a.
- Lacustrine and fluvial sandstone, criteria: Picard and High, 1972a.

SURVEY RELEASES (continued from page 2)

Map 31, "Geology of the Escalante-Boulder Area," by C. Carew McFall (\$2.00). Single multicolored sheet showing the geology of the Escalante-Boulder area.

Map 33, "Oil-Impregnated Rock Deposits of Utah," compiled by Howard R. Ritzma (\$1.50, two sheets). Oil-impregnated rock deposits are located and accompanying text and tables summarize the geology, economic significance and sulfur content of oils extracted from the deposits; updates preliminary Map 25 published in 1968.

- Mineral Fork Formation, Wasatch Mountains: Varney, 1972.
- Paleoenvironmental reconstruction, Parachute Creek Member: Picard and High, 1972b.
- Paradox Basin: Hite, 1972.
- Pennsylvanian, evaporite and carbonate cycles: Hite and others, 1972.
- Trace fossils, stratigraphic record: Howard, 1972.

SEDIMENTATION

Ferron Formation, central Utah: Hale, 1972.

Utah Lake, recent history: Brimhall, 1972.

STRATIGRAPHY

- Black Shale facies, Green River Formation: Thompson, 1971.
- Carboniferous rock, Causey Dam quadrangle: Mullens, 1971.
- Cenozoic, Black Mountains: Rowley, 1972.
- Chinle Formation: Stewart and others, 1972.
- Cretaceous and Tertiary, Ojo Alamo Sandstone: O'Sullivan and others, 1972.
- Duchesne River Formation, northern Uinta Basin: Andersen, 1972.
- Duchesne River Formation, Uinta Basin: Andersen and Picard, 1972.
- Fowkes Formation and Norwood Tuff: Nelson, 1972.
- Frontier Sandstone Member, Mancos Shale: Maione, 1971.
- Green River Formation, Piceance Creek Basin: Cashion and Donnell, 1972.
- Moenkopi Formation, southeastern Utah: Blakey, 1972.
- Norwood Tuff and Fowkes Formation, northwestern Utah: Nelson, 1971.

Monograph Series No. 2, "Eastern and Northern Utah Coal Fields: Vernal, Henry Mountains, La Sal-San Juan, Sego, Tabby Mountain, Goose Creek, Lost Creek, Coalville and Henrys Fork," by H. H. Doelling and R. L. Graham (\$20.00). The second of three in a series to research records, compile data and complete mapping of Utah coal.

The volume is divided into coal fields, areas and quadrangles. Geographic data, geology, mines

"Civilization exists by geological consentsubject to change without notice." Will Durant

- Paleozoic, central Utah: Welsh, 1972.
- Paleozoic strata, central Utah: Hintze, 1972.
- Paradox Basin, Hermosa Group: Hite, 1972.
- Permian and Triassic strata, southern Utah: Irwin, 1971.
- Pine Hollow and Wasatch formations, Garfield County: Bowers, 1972.
- Precambrian, central Utah: Woodward, 1972b.
- Swan Peak Formation and Eureka Quartzite, northern Utah: Francis, 1972.
- Tertiary strata, Sevier-Sanpete region: Doelling, 1972b.
- Triassic and Jurassic, central Utah: Stokes, 1972.
- Uinta Mountain Group, Precambrian: Wallace, 1972.

STRUCTURAL GEOLOGY

- Basin and Range faults, origin: Crosby, 1972.
- Central Utah, structural development: Burchfiel and Hickcox, 1972.
- Great Basin, Cenozoic faults: Rowley and Anderson, 1972.
- Sevier orogenic belt: Beutner, 1972.
- Sevier orogenic belt, western Utah: Armstrong, 1972.
- Smithfield quadrangle, eastern part: Galloway, 1971.
- Stable and unstable folding, criteria: Savage and Sowers, 1972.
- Tintic Quartzite, Willard thrust: Hammond, 1971.

Uinta Mountains, faulting: Ritzma, 1971.

- Willard thrust and Cache allochthon: Crittenden, 1972b; Eardley, 1972.
- Willard thrust, direction: Hammond and Parry, 1972.

and development, land ownership and control, coal quality and reserves are treated.

Multicolored maps, and numerous other figures and tables illustrate the 411-page edition.

Monograph Series No. 1, "Southwestern Utah Coal Fields: Alton, Kaiparowits Plateau and Kolob-Harmony," is also available and sells for \$17.00; No. 3, "Central Utah Coal Fields: Wasatch Plateau, Book Cliffs, Sevier-Sanpete and Emery" plus a summary of Utah's total coal picture will be released in the near future and will sell for \$20.00. The set can be purchased for \$50.00.

May 1973

SUMMER FIELD WORK IN UTAH

The geologists who plan to work in Utah during the 1973 field season are listed below. The reference numbers in the left column correspond as far as possible with the location numbers on the accompanying map.

- 1 Anderson, J. J. Kent State Univ.
- 2 Anderson, O. L. and Gordon Jacoby Lake Powell Research
- 3 Averitt, Paul and R. L. Gee Threet qua USGS
- 4 Biesinger, J. C. Utah State Univ.
- 5 Bilbey, Sue Ann Utah State Univ.
- 6 Bircher, J. E. Utah State Univ.
- 7 Blau, J. G. Utah State Univ.
- 8 Chappell, J. C. Utah State Univ.
- 9 Compton, Robert Stanford Univ.
- 10 Crittenden, M. D., Jr. USGS
- 11 Dawson, M. R. Carnegie Museum
- 12 Doelling, H. H. UGMS
- 13 Fuller, R. H. Utah State Univ.
- 14 Gardiner, L. L. Utah State Univ.
- 15 Goeltz, N. Sylvia UGMS
- 16 Gray, W. E. Utah State Univ.
- 17 Hardy, C. T. Utah State Univ.
- 18 Hickcox, C. W. Univ. of Florida
- 19 Jones, R. L. Utah State Univ.
- 20 Kaliser, B. N. UGMS
- 21 Kohler, J. F. Utah State Univ.
- 22 Mecham, Brent Utah State Univ.

- Geology of southwest High Plateaus and Black Mountains, Piute, Garfield, Iron and Beaver Counties.
- Geohydrology of Kane, Garfield and San Juan Counties.
- L. Geologic mapping of the Cedar City quadrangle, Iron County.

Mineral assemblages in core samples and their geochemical significance, Bear Lake, Utah-Idaho.

Geochemistry and petrography of the Morrison Formation, Dinosaur Quarry quadrangle, Uintah County.

Stratigraphy and environmental analysis of the Worm Creek Member of the St. Charles Formation, north central Utah.

Geology of southern part of James Peak quadrangle, Cache County.

- Mineralization in the Bear River Range, Utah-Idaho.
- General geologic mapping, south Grouse Creek Mountains, Box Elder County.
- Detailed mapping of Willard Peak area, Ogden Canyon to Brigham City.
- Upper Eocene mammals, Uinta Basin.
- (a) Box Elder County minerals inventory;
 (b) Garfield County minerals inventory;
 (c) Kane County minerals inventory;
 (d) East Sego coal field mapping;
 (e) Paradox Basin potash evaluation study.
- Chemical exchange at the sediment-water interface, Bear Lake, Utah-Idaho.
- Environmental analysis of the Upper Cambrian Nounan Formation, Bear River Range, and Wellsville Mountains, north central Utah.
 - Environmental geological atlas, portion of the Wasatch Front area.
 - Structural geology of the south part of Clarkston Mountain, Malad Range, Utah-Idaho.
 - (a) Structural geology of James Peak quadrangle, Cache County; (b) Structural geology of Malad and Bannock Ranges, Utah-Idaho.
 - Pavant Range allochthon, central Utah.
 - Clay-water systems and cation-exchange equilibria in three forest soils, Bear River Range, Cache County.
 - (a) Hydrogeologic study, Salt Lake City; (b) Slope stability and geology, Davis County, and Crandall Canyon, Summit County; (c) Gypsiferous residual soils, St. George area;
 (e) Hydrogeologic evaluation, Bonneville Salt Flats.
 - Morphology, sediment distribution and geochemistry of relict lake-margin deposits in the Rozel Flats area, north Utah.
- Petrography and geochemistry of the Fish Haven and Laketown Dolomites, north central Utah and southeast Idaho.



- 23 Mendenhall, A. J. Utah State Univ.
- 24 Oaks, R. Q., Jr., and R. R. Alexander Utah State Univ.
- 25 Peterson, Carol UGMS
- 26 Ritzma, H. R. UGMS
- 27 Rowley, P. D. USGS
- 28 Summers, P. L. Utah State Univ.
- 29 Todd, V. R. USGS
- 30 Van Horn, Richard USGS
- 31 Weiss, M. P. Northern Ill. Univ.
- 32 Whelan, J. A. and Lee Perry UGMS
- 33 Yeates, E. A. Univ. of Utah
- 34 Zeller, H. D., Fred Peterson and W. E. Bowers USGS

- Structural geology of east part of Richmond and west part of Naomi Peak quadrangles, Utah-Idaho.
- Depositional environments and stratigraphy of Middle and Upper Cambrian units of north Utah and south Idaho.
- Geothermal investigation of Thermal and Roosevelt Hot Springs.
- (a) Fracture and lineament studies, NE Uinta Basin;
 (b) Informational core drilling in tar sands;
 (c) Structure mapping project;
 (d) Stratigraphic study in Circle Cliffs area;
 (e) Oil and gas field study, Boundary Butte area.
- Geology of the Iron Springs Mining District, Iron County.
- Surficial geology of intermontane valleys, Mount Pisgah quadrangle, Cache County.
- Stratigraphy and structure of Upper Paleozoic, Mesozoic and Tertiary rocks, Matlin Mountains, Box Elder County.
- Detailed geologic ground mapping, Salt Lake City.
- Geology of Flagstaff Formation, Sanpete County and vicinity.
- (a) Alteration studies of the Rocky, Star and Indian Peak Ranges; (b) Fluorite studies of Wah Wah and Shauntie Hills.
- The Intermountain seismic belt and quake hazards along the Wasatch Fault, east of Salt Lake City.
- Coal, geology and stratigraphic studies, Kaiparowits Plateau, Kane and Garfield Counties.

Readiness Reduces Disaster Loss

Although earthquake losses can be reduced substantially only by adequate and enforced building codes, zoning provisions and vigorous community programs designed to strengthen disaster preparedness, an individual can lessen the dangers to himself and his family by learning what to do in the event of an earthquake.

Actual movement of the ground in an earthquake is seldom the direct cause of death or injury; most casualties result from falling objects and debris. Landslides and huge seismic sea waves generated by earthquakes also can cause great damage.

Precautionary actions *before* an earthquake can reduce the dangers:

Citizens should support local safe building codes with efficient inspection and firm enforcement;

School building programs should be supported which provide for the strengthening of old, weak buildings or their replacement with earthquake-resistant structures;

Homeowners or tenants should bolt down or provide other strong support for water heaters and other gas appliances since fire damage can result from broken gas lines and appliance connections;

Sites for construction should be selected and engineered to reduce the hazard of damage from an earthquake;

Occasional home earthquake drills can provide knowledge to avoid injury and panic during an earthquake;

A flashlight and a battery powered transistor radio should be ready for use at all times, and

Immunizations should be kept up to date for all family members.

During an earthquake:

If indoors, watch for falling objects. Stay away from windows, mirrors and chimneys. If shaking is severe, get under a table, desk or bed. Usually, it is best not to run outside;

If in a high-rise office building, get under a desk. Do not dash for exits, since stairways may be broken and jammed with people;



Above: Dissolution of gypsum salts and resultant collapse of ground surface, St. George area. Right: Close-up of "miniature karsts."



<u>HOLE-IN-ONE</u> Gypsum Dissolves Out, Surface Sinks In

Gypsum, a salt occurring naturally in the substrata, dissolves readily and can cause the ground surface to subside.

The resulting problems are extensive in the St. George area, southeast Utah, because precipitation and runoff waters introduced into the substrata are absorbed easily into the mainly silty and sandy residual soils, playing havoc with the included gypsum cement.

According to Bruce N. Kaliser, Utah Survey engineering

If in a crowded store, do not rush for a doorway. Others may have the same idea, and

If outside, avoid high buildings, walls, power poles and other objects which could fall. Do not run through the streets.

After an earthquake:

Do not attempt to move seriously injured persons unless they are in immediate danger of further injury;

Do not touch downed power lines or objects touched by downed wires. Immediately clean up spilled medicines, drugs and other potentially harmful materials. Emergency water may be obtained from water heaters, toilet tanks, melted ice cubes and canned vegetables or fruits;

Telephones should not be used except for emergencies;

Do not go sightseeing, particularly in beach and waterfront areas where seismic waves could strike, and geologist, this has been one of the wettest water years of recordmore than 4.61 inches of precipitation in the first six months (October through March), or 188 percent of the normal for that period-which has aggravated the situation.

He describes the surface manifestation of the subsidence as a "miniature karst," and is studying its relevance to homeowners.

Be prepared to respond to requests for help from police, fire-fighting, civil defense and relief organizations.

Accurate prediction of the time and place of earthquakes cannot be made yet. They will continue to harass mankind, occurring most frequently in areas where they have been relatively common in the past.

The Utah Highway Department will use 7 million tons of tailings from Kennecott Copper Corp. flotation concentrators at Magna and Arthur to construct a 5-mile interstate section west of Salt Lake Valley; total project cost is \$13.5 million. The tailings will be transported moist to reduce dust.

UGMS REORGANIZES (continued from page 1)

representatives of various segments of the mineral industry throughout the State (such as solid fuels, metals and hydrocarbons), one reflecting the economic or scientific interests of the mineral industry in the State and one interested in the goals of the Survey from the public at large. The director of the Division of State Lands will be an *ex officio* member without voting privileges.

Future directors of the Survey will be appointed by the board with the concurrence of the executive director of the Department of Natural Resources. He will be the executive and administrative head of the Survey and will be officially designated as the State geologist.

The move was made to consolidate administrative offices of the Survey and to provide more effective representation before the State legislature.

Earthquake Epicenters

General earthquake epicenters in or near Utah for November and December 1972, with dates of occurrence and approximate

UTAH GEOLOGICAL AND MINERALOGICAL SURVEY 103 Utah Geological Survey Building

THE UNIVERSITY OF UTAH SALT LAKE CITY, UTAH 84112

Address correction requested

magnitudes, are listed below. Unless otherwise indicated, localities are in Utah.

Magnituda

	Durchard
November	-0.1
2 Near Pineview	1.8
3 Near Cedar City	<2.0
7 Heber Valley	2.5
8 Heber Valley	2.8
8 South end of Utah Lake	1.8
9 Near Logan	2.0
9 South Utah-Nevada border	<2.0
12 Heber Valley	1.7
13 Near Mount Pleasant	2.0
13 Near Mount Pleasant	2.0
14 Near Kimball Junction	2.1
15 Near Woodruff	3.0
15 Near Emery	3.1
16 Cedar City (felt)	3.3
17 South Utah-Nevada border	2.3
17 South Utah-Nevada border	2.2
17 South Utah-Nevada border	3.2
22 Near Enhraim	22
22 Near Kimball Junction	17
22 Near Kindan Junction	
23 South Otan-Nevada bolder	
24 North of Montpeller, Idano	
20 Near Hamlin Vallay, south	
Litch Newsda border	25
	5.5
29 Near Hamin Valley, south	
Utah-Nevada border	4.0
30 North of Vernal	2.5
30 Heber Valley	<2.0
P 1	
December	
1 Near Richmond	2.1
1 Near Waterpocket Fold	2.0
5 Near Rangley, Colorado	2.3
6 Southeast of Logan	2.6
6 Southeast of Logan	2.2
6 Southeast of Logan	2.0
8 Near Cedar City	2.0
8 Southeast of Thermopolis,	
Wyoming (felt)	4.2

9 South of Castle Dale 2.8

9 Near Bear Lake 3.4

9 South of Nephi 2.5

10 North of LaSal 2.0
11 North of Bear Lake in Idaho 3.7
11 North of Bear Lake in Idaho 2.5
15 Near Bear Lake 2.0
17 Near Cedar City 2.0
19 Near Heber City 1.6
19 Near Heber City 1.1
23 South of Sunnyside 2.1
24 Near Heber City 3.0
29 Near south end of Great Salt Lake 2.5
29 Near south end of Great Salt Lake 2.2
29 Near south end of Great Salt Lake 2.2
30 Near south end of Great Salt Lake 2.4
31 Near Cedar City 2.0
31 Near Levan 2.0

These earthquakes were recorded by the University of Utah seismograph stations under the direction of Kenneth L. Cook. All locations and magnitudes are preliminary determinations; the final determinations will be printed in the University of Utah Seismological Bulletin, issued quarterly

QUARTERLY REVIEW
State of Utah Calvin L. Rampton Governor
Jniversity of Utah Alfred C. Emery President
College of Mines & Mineral
Industries Alex G. Oblad Acting Dean
Jtah Geological & Mineralogical Survey
UTAH GEOLOGICAL AND
MINERALOGICAL SURVEY
103 Utah Geological Survey Building University of Utah Salt Lake City, Utah
84112

Nonprofit Org. U. S. Postage Paid Permit No. 1529 Salt Lake City, Utah